

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Massachusetts High School Survey Risk Behaviors and Sex of Sexual Partners Report

#### Unintentional Injuries/Violence

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN9: Percentage of students who never or rarely wore a seat belt when riding in a car driven by someone else	11.4 (9.3-14.0)	1,226	13.8 (9.4-19.9)	142			●
QN10: Percentage of students who rode one or more times during the past 30 days in a car or other vehicle driven by someone who had been drinking alcohol	24.3 (21.6-27.2)	1,222	30.0 (23.2-37.7)	142			●
QN11: Among students who drove a car or other vehicle during the past 30 days, the percentage who drove when they had been drinking alcohol one or more times during the past 30 days	9.5 (7.1-12.5)	653	18.5 (10.1-31.3)	63			●
QN12: Among students who drove a car or other vehicle during the past 30 days, the percentage who texted or e-mailed while driving on one or more of the past 30 days	41.7 (36.2-47.5)	663	46.2 (30.6-62.7)	66			●
QN13: Percentage of students who carried a weapon such as a gun, knife, or club on one or more of the past 30 days	15.9 (13.4-18.8)	1,178	23.7 (17.1-31.8)	137		●	
QN14: Percentage of students who carried a gun on one or more of the past 30 days	3.7 (2.6-5.2)	1,216	7.8 (4.0-14.5)	140			●
QN15: Percentage of students who carried a weapon such as a gun, knife, or club on school property on one or more of the past 30 days	4.0 (2.7-5.8)	1,219	8.1 (4.7-13.6)	138			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Massachusetts High School Survey Risk Behaviors and Sex of Sexual Partners Report

#### Unintentional Injuries/Violence

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN16: Percentage of students who did not go to school on one or more of the past 30 days because they felt they would be unsafe at school or on their way to or from school	4.3 (3.3-5.7)	1,218	13.0 (8.4-19.7)	140		●	
QN17: Percentage of students who had been threatened or injured with a weapon such as a gun, knife, or club on school property one or more times during the past 12 months	5.1 (4.2-6.2)	1,226	14.5 (10.1-20.4)	142		●	
QN18: Percentage of students who were in a physical fight one or more times during the past 12 months	26.3 (23.9-28.8)	1,212	39.7 (30.0-50.2)	135		●	
QN19: Percentage of students who were in a physical fight one or more times during the past 12 months in which they were injured and had to be treated by a doctor or nurse	2.8 (2.2-3.6)	1,215	5.6 (2.6-11.6)	137			●
QN20: Percentage of students who were in a physical fight on school property one or more times during the past 12 months	6.2 (4.9-7.8)	1,216	11.9 (5.9-22.4)	137			●
QN24: Percentage of students who had ever been bullied on school property during the past 12 months	17.9 (15.5-20.5)	1,221	40.4 (31.2-50.4)	137		●	
QN25: Percentage of students who had ever been electronically bullied during the past 12 months	16.6 (14.4-19.0)	1,222	29.8 (20.9-40.6)	138		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Massachusetts High School Survey Risk Behaviors and Sex of Sexual Partners Report

#### Unintentional Injuries/Violence

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN26: Percentage of students who felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	24.2 (21.1-27.7)	1,221	45.8 (36.7-55.2)	140		●	
QN27: Percentage of students who seriously considered attempting suicide during the past 12 months	13.1 (11.6-14.7)	1,215	37.6 (29.8-46.1)	140		●	
QN28: Percentage of students who made a plan about how they would attempt suicide during the past 12 months	12.0 (10.0-14.2)	1,215	29.0 (22.1-37.0)	138		●	
QN29: Percentage of students who actually attempted suicide one or more times during the past 12 months	5.4 (4.0-7.4)	1,105	19.4 (12.7-28.6)	130		●	
QN30: Percentage of students who made a suicide attempt during the past 12 months that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse	1.4 (0.9-2.1)	1,098	9.6 (4.8-18.3)	129		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Massachusetts High School Survey Risk Behaviors and Sex of Sexual Partners Report

#### Tobacco Use

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN31: Percentage of students who ever tried cigarette smoking, even one or two puffs	45.7 (42.4-48.9)	1,210	60.1 (50.6-69.0)	135		●	
QN32: Percentage of students who smoked a whole cigarette for the first time before age 13 years	6.6 (5.2-8.5)	1,213	19.9 (12.3-30.6)	141		●	
QN33: Percentage of students who smoked cigarettes on one or more of the past 30 days	16.5 (14.6-18.6)	1,210	32.9 (26.2-40.4)	139		●	
QNFRCIG: Percentage of students who smoked cigarettes on 20 or more of the past 30 days	5.4 (4.1-7.2)	1,210	10.6 (6.4-17.1)	139			●
QN36: Percentage of students who smoked cigarettes on school property on one or more of the past 30 days	4.3 (3.0-6.1)	1,217	12.2 (6.9-20.7)	140		●	
QN39: Percentage of students who used chewing tobacco, snuff, or dip on one or more of the past 30 days	7.9 (5.8-10.8)	1,222	8.7 (4.7-15.5)	141			●
QN40: Percentage of students who smoked cigars, cigarillos, or little cigars on one or more of the past 30 days	16.6 (13.9-19.7)	1,223	20.3 (14.3-28.0)	140			●
QNANYTOB: Percentage of students who smoked cigarettes or cigars or used chewing tobacco, snuff, or dip on one or more of the past 30 days	27.1 (24.7-29.7)	1,206	39.2 (31.3-47.6)	137		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Massachusetts High School Survey Risk Behaviors and Sex of Sexual Partners Report

#### Alcohol/Other Drug Use

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN41: Percentage of students who had at least one drink of alcohol on one or more days during their life	82.5 (79.5-85.1)	1,209	82.6 (76.1-87.6)	139			●
QN43: Percentage of students who had at least one drink of alcohol on one or more of the past 30 days	53.4 (49.6-57.1)	1,197	54.3 (45.5-62.9)	135			●
QN44: Percentage of students who had five or more drinks of alcohol in a row, that is, within a couple of hours, on one or more of the past 30 days	30.4 (27.1-34.0)	1,206	34.1 (25.7-43.8)	136			●
QN47: Percentage of students who used marijuana one or more times during their life	63.0 (60.0-65.9)	1,204	68.9 (59.6-76.9)	135			●
QN48: Percentage of students who tried marijuana for the first time before age 13 years	9.5 (7.7-11.6)	1,205	24.4 (15.6-36.1)	135		●	
QN49: Percentage of students who used marijuana one or more times during the past 30 days	38.8 (35.9-41.8)	1,216	49.0 (39.6-58.4)	138		●	
QN50: Percentage of students who used any form of cocaine, including powder, crack, or freebase one or more times during their life	4.9 (3.3-7.3)	1,213	18.9 (11.7-29.2)	141		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Massachusetts High School Survey  
Risk Behaviors and Sex of Sexual Partners Report

Alcohol/Other Drug Use

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN53: Percentage of students who used methamphetamines one or more times during their life	2.2 (1.3-3.7)	1,220	10.1 (5.5-17.8)	142		●	
QN54: Percentage of students who used ecstasy one or more times during their life	7.3 (5.7-9.3)	1,226	16.6 (10.6-25.0)	141		●	
QN55: Percentage of students who took steroid pills or shots without a doctor's prescription one or more times during their life	2.0 (1.4-2.9)	1,225	7.7 (4.7-12.4)	141		●	
QN57: Percentage of students who used a needle to inject any illegal drug into their body one or more times during their life	0.7 (0.3-1.4)	1,222	8.0 (4.0-15.3)	140		●	
QN58: Percentage of students who were offered, sold, or given an illegal drug by someone on school property during the past 12 months	27.8 (24.7-31.1)	1,223	43.2 (35.4-51.4)	141		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Massachusetts High School Survey Risk Behaviors and Sex of Sexual Partners Report

#### Sexual Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN59: Percentage of students who ever had sexual intercourse	72.3 (67.4-76.7)	1,139	79.1 (67.2-87.4)	126			●
QN60: Percentage of students who had sexual intercourse for the first time before age 13 years	5.2 (4.1-6.6)	1,131	10.7 (5.0-21.5)	127			●
QN61: Percentage of students who had sexual intercourse with four or more people during their life	16.7 (14.6-19.0)	1,133	26.5 (17.9-37.3)	127		●	
QN62: Percentage of students who had sexual intercourse with one or more people during the past three months	52.4 (48.0-56.9)	1,137	56.4 (46.2-66.0)	127			●
QN63: Among students who had sexual intercourse during the past three months, the percentage who drank alcohol or used drugs before last sexual intercourse	22.9 (19.3-26.9)	592	24.4 (16.1-35.1)	70			●
QN64: Among students who had sexual intercourse during the past three months, the percentage who did not use a condom during last sexual intercourse	38.7 (33.7-43.9)	584	73.0 (59.3-83.4)	69		●	
QN65: Among students who had sexual intercourse during the past three months, the percentage who did not use birth control pills to prevent pregnancy before last sexual intercourse	73.6 (69.0-77.7)	578	63.9 (52.1-74.2)	67			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Massachusetts High School Survey Risk Behaviors and Sex of Sexual Partners Report

#### Sexual Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QNOTHH: Among students who had sexual intercourse during the past three months, the percentage who did not use an IUD or implant or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	92.1 (88.6-94.6)	578	96.0 (87.4-98.8)	67			●
QNOTHHPL: Among students who had sexual intercourse during the past three months, the percentage who did not use birth control pills, an IUD or implant, or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	65.7 (60.9-70.2)	578	59.8 (48.4-70.3)	67			●
QNQUALBC: Among students who had sexual intercourse during the past three months, the percentage who did not use both a condom during last sexual intercourse and birth control pills, an IUD or implant, or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	87.3 (84.0-89.9)	574	94.8 (89.1-97.6)	67		●	
QNBCNONE: Among students who had sexual intercourse during the past three months, the percentage who used no method of birth control to prevent pregnancy before last sexual intercourse	7.9 (6.1-10.3)	578	24.5 (15.4-36.7)	67		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Massachusetts High School Survey Risk Behaviors and Sex of Sexual Partners Report

#### Dietary Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>‡</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>‡</sup>	No Difference <sup>‡</sup>
	% 95% CI*	N	% 95% CI	N			
QNOWT: Percentage of students who were overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex) <sup>§</sup>	13.2 (11.0-15.7)	1,181	14.8 (9.6-22.2)	136			●
QNOBESE: Percentage of students who were obese (i.e., at or above the 95th percentile for body mass index, by age and sex) <sup>§</sup>	9.1 (7.4-11.1)	1,181	11.5 (6.6-19.4)	136			●
QN66: Percentage of students who described themselves as slightly or very overweight	26.6 (23.9-29.5)	1,219	41.2 (34.0-48.8)	143		●	
QN68: Percentage of students who went without eating for 24 hours or more to lose weight or to keep from gaining weight during the past 30 days	10.5 (8.8-12.4)	1,216	26.5 (18.5-36.4)	140		●	
QN69: Percentage of students who took any diet pills, powders, or liquids without a doctor's advice to lose weight or to keep from gaining weight during the past 30 days	4.1 (3.1-5.3)	1,214	5.9 (3.3-10.2)	140			●
QN70: Percentage of students who vomited or took laxatives to lose weight or to keep from gaining weight during the past 30 days	3.8 (3.0-4.8)	1,200	15.5 (10.3-22.7)	141		●	
QN73: Percentage of students who did not eat green salad during the past seven days	33.9 (30.2-37.7)	1,214	37.6 (29.7-46.3)	140			●

\*Confidence interval.

<sup>‡</sup>Based on t-test analyses, p<0.05.

<sup>§</sup>Based on reference data from the 2000 CDC Growth Charts.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Massachusetts High School Survey Risk Behaviors and Sex of Sexual Partners Report

#### Dietary Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN74: Percentage of students who did not eat potatoes during the past seven days	27.1 (24.3-30.2)	1,207	37.1 (28.4-46.8)	141		●	
QN77: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day during the past seven days	16.2 (14.0-18.8)	1,209	29.3 (21.3-38.7)	141		●	
QNSODA0: Percentage of students who drank a can, bottle, or glass of soda or pop during the past seven days	75.1 (72.6-77.4)	1,209	78.2 (70.5-84.3)	141			●
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day during the past seven days	10.4 (8.6-12.5)	1,209	21.8 (15.5-29.8)	141		●	
QNSODA3: Percentage of students who drank a can, bottle, or glass of soda or pop three or more times per day during the past seven days	5.3 (4.0-6.9)	1,209	12.1 (7.8-18.3)	141		●	
QN78: Percentage of students who did not drink milk or who drank less than three glasses per day of milk during the past seven days	88.9 (86.9-90.6)	1,201	92.3 (85.6-96.1)	141			●
QNMILK0: Percentage of students who drank 0 glasses per day of milk during the past seven days	17.0 (14.2-20.2)	1,201	28.2 (19.6-38.7)	141		●	
QNMILK1: Percentage of students who drank less than one glass per day of milk during the past seven days	58.6 (54.8-62.4)	1,201	69.2 (59.3-77.6)	141			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Massachusetts High School Survey  
Risk Behaviors and Sex of Sexual Partners Report

Dietary Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI <sup>*</sup>	N	% 95% CI	N			
QNMILK2: Percentage of students who drank less than two glasses per day of milk during the past seven days	74.8 (71.8-77.5)	1,201	81.4 (72.9-87.8)	141			●
QN79: Percentage of students who did not eat breakfast on all of the past seven days	63.6 (59.8-67.2)	1,202	73.4 (65.4-80.1)	141		●	
QNNOBKFT: Percentage of students who ate breakfast on none of the past seven days	11.9 (9.5-14.8)	1,202	17.4 (10.6-27.3)	141			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Massachusetts High School Survey  
Risk Behaviors and Sex of Sexual Partners Report

Physical Activity

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN80: Percentage of students who were physically active for a total of at least 60 minutes per day less than five of the past seven days	52.2 (48.2-56.1)	1,201	60.4 (51.5-68.7)	141			●
QNPA0DAY: Percentage of students who were physically active for a total of at least 60 minutes per day on 0 of the past seven days	11.4 (9.9-13.2)	1,201	23.1 (17.9-29.4)	141		●	
QNPA7DAY: Percentage of students who were physically active for a total of at least 60 minutes per day on less than seven of the past seven days	73.5 (70.3-76.5)	1,201	79.5 (71.2-85.9)	141			●
QN81: Percentage of students who watched three or more hours per day of TV on an average school day	25.2 (22.6-28.0)	1,203	28.4 (20.3-38.3)	139			●
QN82: Percentage of students who played video or computer games or used a computer for something that was not school work three or more hours per day on an average school day	38.0 (34.5-41.6)	1,206	50.8 (39.5-62.1)	139		●	
QN83: Percentage of students who did not attend physical education (PE) classes in an average week when they were in school	46.9 (39.3-54.6)	1,185	46.8 (37.4-56.5)	140			●
QNDLYPE: Percentage of students who attended physical education (PE) classes less than daily in an average week when they were in school	82.7 (76.6-87.4)	1,185	84.8 (76.2-90.6)	140			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Massachusetts High School Survey  
Risk Behaviors and Sex of Sexual Partners Report

Physical Activity

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI <sup>*</sup>	N	% 95% CI	N			
QN84: Percentage of students who did not play on sports teams during the past 12 months	37.1 (33.2-41.2)	1,199	51.2 (43.3-59.2)	140		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Massachusetts High School Survey  
Risk Behaviors and Sex of Sexual Partners Report

Other

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>‡</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>‡</sup>	No Difference <sup>‡</sup>
	% 95% CI <sup>*</sup>	N	% 95% CI	N			
QN85: Percentage of students who were not sure if they had ever been taught or who had not ever been taught in school about AIDS or HIV infection	11.7 (9.1-14.9)	1,225	15.5 (10.4-22.6)	141			●

\*Confidence interval.

<sup>‡</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Massachusetts High School Survey Risk Behaviors and Sex of Sexual Partners Report

#### Site-Added

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN87: Percentage of students who would not describe their grades in school as mostly A's or mostly B's during the past 12 months	31.9 (27.4-36.7)	1,216	39.3 (29.8-49.6)	143			●
QN88: Percentage of students who have lived in the United States less than one year	0.3 (0.1-0.9)	1,219	2.3 (0.8-6.8)	141			●
QN89: Percentage of students who typically do not sleep at home with their parents or guardians at night	4.3 (3.1-5.9)	1,222	9.8 (5.3-17.5)	142			●
QN91: Percentage of students who are transgender	1.2 (0.7-2.1)	1,221	9.0 (5.3-14.9)	141		●	
QN92: Percentage of students who have any long-term learning disabilities	7.7 (5.8-10.0)	1,221	9.7 (6.3-14.7)	141			●
QN93: Percentage of students who have physical disabilities or long-term health problems	11.1 (9.5-13.0)	1,222	14.7 (9.9-21.3)	140			●
QN94: Percentage of students who disagree, strongly disagree, or are not sure that their teachers really care about them and give them encouragement and support	32.8 (29.0-36.8)	1,220	43.1 (33.4-53.4)	141		●	
QN95: Percentage of students who cannot talk with or are not sure if they can talk with at least one of their parents or other adult family members about things that are important to them	17.2 (14.9-19.7)	1,208	31.2 (24.6-38.7)	139		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Massachusetts High School Survey Risk Behaviors and Sex of Sexual Partners Report

#### Site-Added

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN96: Among students who drove a car or other vehicle during the past 30 days, the percentage who talked on a cell phone while driving on one or more of the past 30 days	49.5 (42.1-56.9)	707	46.6 (34.1-59.5)	71			●
QN97: Percentage of students who have been a member of a gang during the past 12 months	6.8 (5.3-8.7)	1,224	17.8 (11.4-26.7)	140		●	
QN98: Percentage of students who have ever been hurt physically by a date or someone they were going out with	10.8 (9.1-12.7)	1,222	28.1 (18.7-39.9)	140		●	
QN99: Percentage of students who have ever had sexual contact against their will	10.2 (8.6-12.0)	1,216	33.5 (24.8-43.5)	139		●	
QN100: Percentage of students who did something to purposely hurt themselves without wanting to die, such as cutting or burning themselves on purpose, one or more times during the past 12 months	15.0 (12.6-17.8)	1,223	42.7 (33.8-52.0)	142		●	
QN101: Percentage of students who had at least one drink of alcohol on school property on one or more of the past 30 days	3.6 (2.6-4.9)	1,225	16.5 (10.3-25.2)	142		●	
QN102: Percentage of students who used marijuana on school property one or more times during the past 30 days	5.7 (4.5-7.3)	1,224	19.5 (11.6-30.8)	142		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Massachusetts High School Survey Risk Behaviors and Sex of Sexual Partners Report

#### Site-Added

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN103: Percentage of students who have used heroin (also called smack, junk, or diesel) one or more times during their life	1.2 (0.7-2.1)	1,224	9.4 (4.9-17.6)	142		●	
QN104: Percentage of students who sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high one or more times during the past 30 days	2.5 (1.7-3.7)	1,225	13.6 (8.1-21.9)	141		●	
QN105: Percentage of students who have not ever been taught or are not sure if they have ever been taught in school about how to use condoms	40.5 (33.4-47.9)	1,223	38.2 (28.7-48.7)	139			●
QN106: Percentage of students who did not talk at all with their parents or other adults in their family about sexuality or ways to prevent HIV infection, other sexually transmitted diseases (STDs), or pregnancy during the past 12 months	51.8 (48.7-55.0)	1,221	47.7 (39.3-56.2)	138			●
QN108: Percentage of students who have been pregnant or gotten someone pregnant one or more times	4.1 (3.0-5.7)	1,221	11.5 (6.1-20.7)	143		●	
QN109: Percentage of students who have not ever been tested or are not sure if they have ever been tested for HIV, the virus that causes AIDS	84.8 (81.6-87.4)	1,218	73.5 (62.7-82.1)	141	●		
QN110: Percentage of students who have not ever been or are not sure if they have ever been tested for other sexually transmitted diseases (STDs) such as genital herpes, chlamydia, syphilis, or genital warts	82.4 (79.7-84.7)	1,221	69.9 (56.9-80.4)	142	●		

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Massachusetts High School Survey Risk Behaviors and Sex of Sexual Partners Report

#### Site-Added

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN111: Percentage of students who have ever been told by a doctor or nurse that they had HIV infection or any other sexually transmitted disease (STD)	1.8 (1.2-2.7)	1,218	5.1 (2.2-11.2)	142			●
QN112: Percentage of students who did not eat fruit or drink 100% fruit juices during the past 7 days	8.4 (6.7-10.5)	1,213	14.9 (9.2-23.1)	141			●
QN113: Percentage of students who did not eat other vegetables such as carrots, peas, or broccoli during the past 7 days	13.2 (10.7-16.1)	1,211	17.3 (11.3-25.7)	141			●
QN114: Percentage of students who drank a can, bottle, or glass of a sports drink such as Gatorade or PowerAde one or more times per day during the past seven days	11.6 (9.8-13.7)	1,197	12.8 (8.3-19.4)	141			●
QN115: Percentage of students who had not been taught or are not sure if they had ever been taught in school about how to choose foods that will keep you healthy and how to eat a balanced, nutritious diet	19.5 (16.8-22.5)	1,189	20.1 (14.8-26.7)	140			●
QN116: Percentage of students who are eligible to get a free or reduced-price lunch at school	34.5 (28.5-41.0)	1,195	42.8 (34.6-51.4)	137		●	
QN117: Percentage of students who have not ever been taught in school or are not sure if they have ever been taught in school how to follow a personal fitness plan, including setting fitness goals for themselves and keeping track of their progress	36.6 (31.2-42.4)	1,197	46.2 (37.4-55.3)	139		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Massachusetts High School Survey  
Risk Behaviors and Sex of Sexual Partners Report

Site-Added

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI <sup>*</sup>	N	% 95% CI	N			
QN118: Percentage of students who did not do volunteer work, community service, or help people outside of their home without getting paid during an average month when they are in school	48.5 (44.2-52.7)	1,205	54.6 (45.6-63.3)	139			●

<sup>\*</sup>Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.