

Yearly Operational Plan (YOP)
Transmission Line Vegetation Management
Massachusetts Municipal Wholesale
Electric Company

2012

Prepared By:

Michael DiMauro
Environmental Engineer
Engineering Services Department
327 Moody Street
Ludlow, Massachusetts 01056
(413) 308-1243
mdimauro@mmwec.org

Yearly Operational Plan

The purpose of 333 CMR 11.00, Rights of Way Management, is to promote the implementation of integrated pest management techniques and to establish standards, requirements, and procedures necessary to minimize the risk of unreasonable adverse effects on human health and the environment associated with the use of herbicides to maintain streets, road, sidewalks and paths. These regulations establish procedures that guarantee ample opportunity for public and municipal agency review and input on the right-of-way maintenance plans.

A yearly operational plan (YOP) must be submitted to the Department of Agricultural Resources every year herbicides are intended for use to maintain rights-of-way. The YOP provides a detailed program for vegetation management for the year. A five year Vegetation Management Plan (VMP) covering the period from 2011 – 2015 was approved Department in 2011 and is available for review at the office of the Massachusetts Municipal Wholesale Electric Company (MMWEC), Board of Health, Conservation Commission, and Ludlow Selectman's Office.

Upon receipt of this YOP, the Department publishes a notice in the Environmental Monitor. MMWEC must provide a copy of the proposed YOP and Environmental Monitor notice to the Board of Health, Conservation Commission, and the Selectman's Office for the Towns of Ludlow, Hancock and Lanesborough in which the proposed herbicide treatment is proposed. The Department allows a 45-day comment period on the proposed YOP beginning with the publication of the notice and receipt of the YOP and Environmental Monitor notice by the Town.

Public notification of herbicide application to the rights of way is made at least 21 days in advance of the treatment by a separate notice. Notice is made to the Department of Agricultural Resources, Selectman's Office, Board of Health, and the Conservation Commissions in the Towns of Ludlow, Hancock and Lanesborough.

Any comments on this YOP should be made to the person designated herein as the person supervising the YOP or the person performing the treatment.

Individual Supervising YOP

Name and Title: Michael DiMauro, Environmental Engineer
Department: Engineering Services
Address: MMWEC, 327 Moody Street, Ludlow, MA
Telephone: (413) 308-1243
Signature: _____

Contractor Performing Herbicide Treatment

A licensed herbicide applicator under contract to MMWEC will perform the herbicide treatment. Applicators are certified by the Massachusetts Department of Agricultural Resources in the applicator category:

Certified Applicator: [David O'Brien](#)
Certified Applicator License Number: [22047](#)
Company or Department: [Lewis Tree Service, Inc.](#)
Address: [300 Lucius Gordon Drive,
West Henrietta, NY 14586](#)
Telephone Number: [413- 237-9870](#)
Email: Dobrien@lewistree.com

Yearly Operational Plan

The following information is provided as details of the YOP for the MMWEC Transmission Line Rights of Way in accordance with the requirements of 333 CMR 11.06 (2):

Herbicides Proposed

The herbicides proposed for use in calendar year 2012 are:

<u>Herbicide</u>	<u>EPA Reg</u>	<u>Rate</u>
Rodeo Concentrate	62719-324	3%
Escort XP	352-439	2oz/100 gal
 <u>Carrier/Adjuvant</u>		
Aquafac		1%

Complete information on these products is attached, including the Material Safety Data Sheets for the product. The herbicide fact sheet for the above listed herbicides are attached to and made part of this YOP.

Herbicide Application Techniques and Alternative Control Procedures

The herbicide(s) will be applied in accordance with the instructions in the attached manufacturer's information and as described below. Alternative control procedures, applicable at the designated "No Spray Zones" will consist of hand cutting, mowing, or selective trimming as necessary.

Selected Treatment Method for this YOP

The selected method of herbicide treatment for this YOP is the Selective Foliar method.

Description: A herbicide mixture is applied to fully developed leaves, stems, needles or blades of individual, target species. The herbicide concentrate is usually mixed or diluted with water and applied as a uniform spray over the plant's foliage. Two types of equipment for foliar treatments are used: back pack and vehicle mounted. Both treatments use low pressure, below 60 pounds per square inch (psi) at the nozzle, for applications. This technique is generally the most economical and effective method, particularly in medium and high brush density situations and to control noxious and poisonous vegetation that presents a hazard to inspection and maintenance crews. The application period usually extends from early June through the beginning of leaf abscission in the fall when not restricted by regulations.

Backpack Foliar Techniques utilize hand-operated pumps or motorized, backpack sprayers. The motorized, backpack sprayer produces an air current that delivers the herbicide mixture in small droplets from the portable three to five gallon spray tank to the target vegetation. Both techniques only require the applicator to dampen or lightly wet the target leaf area not to the point of runoff. This minimizes the amount of excess herbicide drip from target species onto desirable ground cover. Low volume applications also eliminate the need to bring heavy equipment on the ROW for the transportation of large quantities of herbicide solution.

Vehicle Mounted Techniques generally utilizes a 100-500 gallon hydraulic sprayer mounted on a truck, tractor or tracked vehicle equipped with hand-held spray guns. The herbicide mixture is directed at specific vegetation or broadcast for uniform coverage. Specially designed showerhead type nozzles reduce spray volumes and limit droplet fines thus reducing the potential for spray drift off-target. These nozzles deliver effective spray coverage at relatively low spray pressures of sixty psi and less. This technique is capable of delivering uniform, penetrating spray coverage to dense, tall, target vegetation such as Phragmites and Japanese Knotweed. It is particularly useful for sites where total weed control or pre-emergent herbicide applications are required for fire safety, such as at pumping stations and valve sites or for initial treatment of invasive plants.

All equipment used for vegetation management programs must be maintained in good working condition, and should be of adequate design and ability to produce the professional quality of work that the Town requires. Because the Town recognizes the vast variety and performance of herbicide application equipment, dictating how that equipment should be calibrated to deliver precise amounts of herbicide to effectively control a host of vegetation conditions is literally impossible. Therefore, the Town insists that the contractor provide the most appropriate application equipment, calibrated to effectively and legally control target vegetation.

Both the Contractor and MMWEC are responsible to insure that vegetation management activities are conducted in a professional, safe, efficient manner, with special attention directed towards minimal environmental impact. The contractor is qualified, licensed and certified to apply herbicides. "Qualified" means those personnel who have been trained to recognize and identify target and non-target vegetation and are knowledgeable in the safe and proper use of both mechanical and chemical vegetation management techniques. All personnel applying herbicides in Massachusetts must be licensed in the Commonwealth and must work under the on-site supervision of a certified applicator. All contract personnel will also follow all label instructions regarding Personal Protective Equipment (PPE).

MMWEC will rely on the independent contractor listed in the YOP for vegetation management applications and requires, in a contractual agreement, that contractors comply with all applicable federal and state laws and regulations. These include, but are not limited to, applicable OSHA, FIFRA and DOT regulations, 333 CMR 1-15.00,

Rights-of-Way Management, Chapter 132B, Chapter 85 of the Acts of 2000 (see Appendix IV) and 321 CMR 10.00 as managed by NHESP.

Herbicides will only be applied in a safe and judicious manner, in compliance with all-applicable State and Federal pesticide regulations.

Applicators will at all times exercise good judgment and common sense during herbicide treatment activities, and will immediately cease operations if adverse conditions or other circumstances warrant.

Herbicides will NOT be applied during the following adverse weather conditions:

- A. During high wind velocity, per 333 CMR 11.03
- B. Foliar applications during periods of dense fog, or moderate to heavy rainfall
- C. Foliar applications of volatile herbicides during periods of high temperatures (90 plus degrees Fahrenheit) and low humidity
- D. Cut Stump applications when deep snow (i.e. 6" plus or ice frozen on stem or stump) prevents adequate coverage of target plants to facilitate acceptable control

The contractors or a representative of MMWEC must complete daily vegetation management reports that include:

- A. Date, name and address of vegetation management contractor(s)
- B. Identification of site or work area
- C. List of crew members
- D. Type of equipment and hours used, both mechanical and chemical
- E. Method of application and description of target vegetation
- F. Amount, concentration, product name of herbicide(s), adjuvants, and dilutants (EPA registration numbers must be on file)
- G. Weather conditions
- H. Notation of any unusual conditions or incidents, including public inquiries
- I. Recording and/or verification of sensitive areas on ROW maps

Identification of Target Vegetation

The target vegetation for this YOP will be all non-desirable, tall growing species of vegetation that could reduce line clearances below required minimum distance for safe and reliable electric transmission.

Vegetation management crews will exercise care to insure that low-growing desirable vegetation and other non-target organisms are not unreasonably affected by the application of herbicides.

Description of Methods Used to Flag or Otherwise Designate Sensitive Areas

The sensitive areas detailed herein are easily recognizable in the field as described and will be marked with an orange painted line in the street.

For the 2012 YOP, it is the intention of MMWEC to spray target species where permitted along the entire 5.2 mile transmission line in Ludlow and the 23 kV collector circuit for the Berkshire Wind Power Project in the Towns of Hancock and Lanesborough, Ma.. The YOP maps identify the Sensitive Areas where “No Spray Zones” have been established, all of which are clearly identified by structures, houses, or other field methods.

With the assistance of the contractor, sensitive areas will be identified and marked along the ROW prior to any herbicide application.

On the MMWEC transmission line in Ludlow, Ma, potential sensitive habitats were identified by the NHESP in 2 locations.

The first area was located between stations 30+25 and T-line Structure #5. The area was previously mapped by NHESP to contain the Climbing Fern (*Lygodium palmatum*). This area was surveyed in accordance with NHESP requirements and protocols in the early fall of 2011 by a certified biologist and it was determined that there were no occurrences of the species in question. However, this area was not sprayed due to scheduling and the timing of the study. This area will not be sprayed in 2012 unless approved by NHESP.

The second area is located between T-Line Structures 1 and 8. This area was previously mapped by NHESP to contain a species of special concern, the Blue Spotted Salamander (*Ambystoma laterale*). The BMPs as described in the ROW vegetation management plan for Vernal Pool Habitat shall be followed in this area.

Sensitive Area Restrictions

333 CMR 11.04

CONTROL STRATEGIES FOR SENSITIVE AREAS

Sensitive Area	Minimum Buffer Zone (feet)	Control Method	Restriction Code
Public Ground Water Supplies	400'	Mechanical Only	None
Primary Recharge Area	Designated buffer zone or 1/2 mile radius	Mechanical, Recommended Herbicides*	1
Public Surface Water Supplies (Class A & Class B)	100'	Mechanical Only	None
	100'-400'	Recommended Herbicides	1
Tributary to Class A Water Source, within 400' upstream of water source	100'	Mechanical Only	None
	100'-400'	Recommended Herbicides	1
Tributary to Class A Water Source, greater than 400' upstream of water source	10'	Mechanical Only	None
	10'-200'	Recommended Herbicides	1
Class B Drinking Water Intake, within 400' upstream of intake	100'	Mechanical Only	None
	100'-200'	Recommended Herbicides	1
Private Drinking Water Supplies	50'	Mechanical Only	None
	50'-100'	Recommended Herbicides	2
Surface Waters	10'	Mechanical Only	None
	10'-100'	Recommended Herbicides	2
Rivers	10' from mean annual high water line	Mechanical Only	None
	10'-200'	Recommended Herbicides	2
Wetlands	100' (treatment in wetlands permitted up to 10' of standing water)* [†]	Low-pressure Foliar, CST, Basal Recommended Herbicides	1
Habitated Areas	100' (for high-pressure foliar only)	Recommended Herbicides	2
Agricultural Area (Crops, Fruits, Pastures)	100' (for high-pressure foliar only)	Recommended Herbicides	2
Certified Vernal Pools	10'	Mechanical Only	None
Certified Vernal Pool Habitat	10'-outer boundary of habitat	As recommended by NHESP in their permit process, no treatment without written permission	
Priority Habitat	As recommended by NHESP in their permit process, no treatment without written permission		

Restriction Code "1": A minimum of twenty-four months shall elapse between applications

Code "2": A minimum of twelve months shall elapse between applications

*Massachusetts recommended herbicides for sensitive sites

[†]Per "The Wetlands Impact Study," 10/1995 (see Appendix III)

Procedures and Locations for Handling, Mixing and Loading of Herbicide Concentrates

The herbicide will be mixed in a controlled environment at prior to entering the right of way. Should mixing become necessary while the contractor is in the right of way, all mixing shall be performed on level ground in the access road away from wetlands and other sensitive areas.

Although it is expected that all the mixed herbicide will be used, any remaining will be safely removed by the contractor at the end of each days work and stored by the contractor in accordance with manufacturer's instructions. The absorbent product "Speedi-Dri" or equivalent will be available for use at the locations of application. If there is a leak in the hose, the pump will be immediately shutoff and removed from the work area in a suitable container for repair. Equipment used will be washed in accordance with all applicable regulations at the contractors place of business.

Herbicides will be handled and applied only in accordance with the label instructions. Contractors will strictly adhere to all mandated safety precautions directed towards the public, the applicator and the environment.

Remedial Plan to Address Spills and Related Accidents

All mixing and loading of herbicides will be conducted at the central facility where the herbicides are stored. Only the amount of herbicide necessary to carry out the vegetation control, based on the monitoring results, will ensure that there will be no waste and minimize potential problems. The vehicle(s) carrying out the spray operations will be equipped with a bag of absorbent, activated charcoal, leak-proof containers, a broom and a shovel in case of minor spills. A clipboard log of the herbicides on the vehicle will be kept on the vehicle. Herbicide labels and fact sheets should be carried on-site by the applicator.

As soon as any spill is observed, immediate action will be taken to contain the spill and protect the spill area. The cause of the spill must be identified and secured. Spill containment will be accomplished by covering the spill with Speedi-Dri or other absorptive material or, for large spills, building clay or soil dikes to impede spill progress. Until completely clean, protection of the spill area will be accomplished by placing barriers, flagging or crewmembers at strategic locations. If a fire is involved, care will be taken to avoid breathing fumes from any burning chemicals.

Minor spills will be remedied by soaking up the spill with Speedi-Dri or other adsorptive material and placing it in leak proof containers, removed from the site and disposed of properly. Dry herbicides, such as granulars, will be swept up or shoveled up directly in leak proof containers for proper disposal. All contaminated soil will be placed in leak proof containers, removed from the site and disposed of properly. Activated charcoal will be incorporated into the soil at the spill location at a rate of seven pounds per thousand square feet to inactivate any herbicide residue. Any minor spill will be reported to the Pesticide Bureau.

Major spills will be handled in a similar manner as minor spills, except in cases where the spill cannot be contained and/or removed by the crew. MassDEP will be contacted when there is a spill of a regulated quantity of material.

In the event of a spill, information on safety precautions and clean up procedures may be gathered from the following sources:

Emergency Contacts

Ludlow

Ludlow DPW 413-583-5625

Ludlow Fire Department 413-583-8332

Ludlow Police Department 413-583-8305

Hancock

Hancock DPW 413-738-5225

Hancock Fire Department 413-738-5446

Hancock Police Department 413-738-5225

Lanesborough

Lanesborough DPW 413-442-1167

Lanesborough Fire Department 413-443-2321

Lanesborough Police Department 413-443-4107

National and State

Bureau Environmental Health Assessment 617-624-5757

Protection Incident Response Unit 888-304-1133

EPA Pesticide Hotline 800-858-7378

ChemTrec (800) 424-9300 Chemical Industry
Emergency Response
System


Mass. Pesticide Bureau (617) 626-1781 For all Pesticide Spills,
Fire, and Related
Accidents

Mass. DEP Western Regional Office (413) 784-1100
888-304-1133 For Emergencies Involving
Reportable Quantities of
Hazardous Materials

BASF Company (800) 454-2673 For Medical Emergencies
Involving Arsenal Herbicide

DowAgroSciences	(800) 992-5994	For Emergencies Involving Dow Products
Du Pont	(800) 441-3637	For Medical Emergencies Involving Du Pont Products
Novartis	(800) 267-6351	For Emergencies Involving Vanquish
Mass. Poison Information Center	(800) 682-9211	For Medical Emergencies Involving Pesticides

YOP
MAPS
LUDLOW



Massachusetts Municipal Wholesale Electric Company


R.W.BECK and ASSOCIATES

ENGINEERS AND CONSULTANTS

STONY BROOK ENERGY CENTER

MMWEC PHASE 1 PROJECT

345 KV TRANSMISSION LINE No. 334



DRAWING NUMBER	TITLE	CONTRACT NUMBER
	345 KV TRANSMISSION LINE NO. 334	T6
2296-G-201	VICINITY MAP AND INDEX	o
2296-C-201	PLAN AND PROFILE (SHEET 1)	o
2296-C-202	PLAN AND PROFILE (SHEET 2)	o
2296-C-203	PLAN AND PROFILE (SHEET 3)	o
2296-C-204	PLAN AND PROFILE (SHEET 4)	o
2296-C-205	PLAN AND PROFILE (SHEET 5)	o
2296-C-206	PLAN AND PROFILE (SHEET 6)	o
2296-C-207	PLAN AND PROFILE (SHEET 7)	o
2296-C-208	PLAN AND PROFILE (SHEET 8)	o
2296-C-209	PLAN AND PROFILE (SHEET 9)	o
2296-C-210	PLAN AND PROFILE (SHEET 10)	o
2296-C-211	CLEARING PLANS & SECTIONS	o
2296-C-212	ACCESS ROAD PLANS & SECTIONS	o
2296-S-201	POLE SETTING & FOUNDATION DETAILS	o
2296-E-201	STRUCTURE WT-1	o
2296-E-202	STRUCTURE WA-1	o
2296-E-203	STRUCTURE WA-2	o
2296-E-204	STRUCTURE WA-3	o
2296-E-205	STRUCTURE WDE-1	o
2296-E-206	STRUCTURE ST-1	o
2296-E-207	LINE AND GUY ATTACHMENT DETAILS	o

o DENOTES CONTRACT DRAWING

No.	Date	By	Chd	Revision
6	5/11	JPH	MD	ADDED RANDALL RD. AND NASH HILL RD.
5	3/93	YL	YL	ADDED BRIDGE RD.
4	3/93	YL	JP	CONVERTED TO CAD FROM R.W.BECK DRG
3	7/90/91	SLK	SLK	CONFORMING TO CONSTRUCTION RECORDS.
2	1-4-89	CJS	RLD	REVISED INDEX
1	1/15/88	CJS	RLD	TS-TRANSMISSION LINE CONSTRUCTION CONTRACT
0	1/31/88	CJS	RLD	TS-CLEARING CONTRACT

REVISIONS



STONY BROOK ENERGY CENTER
MMWEC PHASE 1 PROJECT

R.W.BECK and ASSOCIATES

ANALYTICAL AND CONSULTING ENGINEERS

Seattle, Washington Denver, Colorado

345KV. TRANSMISSION LINE NO.334

VICINITY MAP AND INDEX

Des.	RD	Eng.	RLD	Appr.	Check.	Drawn.	Scale.	Sheet.	Drawn.
Des.	C/S	Eng.	RLD	Appr.	Check.	Drawn.	Scale.	Sheet.	Drawn.
File.	MC-6244-ET2-8A	Date.	1-31-80	2296-G-201.4					




MMWEC
 Massachusetts Municipal Wholesale Electric Company
 A PUBLIC CORPORATION


STONY BROOK ENERGY CENTER

MMWEC PHASE I PROJECT


FIGURE 3.4-3
 TOPOGRAPHIC MAP SHOWING THE SBEC SITE BOUNDARY AND TRANSMISSION CORRIDOR



VICINITY MAP



MMWEC
5.2 MILE, 345 KV TRANSMISSION LINE
LUDLOW, MA.



MERIDIAN OF THE WESTOVER COORDINATE SYSTEM WHICH EQUALS THAT OF THE MASSACHUSETTS COORDINATE SYSTEM MAINLAND ZONE

NOTES

1. FIELD VERIFY LOCATION OFF 345 KV. TAKEOFF STRUCTURE AND P.I. NO.1
2. REFERENCE STRUCTURE DETAIL DRAWINGS FOR PLACEMENT OF O.H.G.W. AND CONDUCTOR DAMPERS.
3. THESE DRAWINGS ARE EXACT COPIES OF R.W. BECK ORIGINALS. R.W. BECK USED 742' SAG TEMPLATE FOR ALL STRUCTURES. SEE MMWEC SKETCH SK-451.

CONTINENTAL FIELD SERVICE CORPORATION
ELMSFORD, NEW YORK

No.	Date	By	Chg	Revision
6	5/11	JPH	MD	ADDED WETLAND TEXT
5	2/27/84	YL	JP	CONVERTED TO CAD, ADDED NOTE 3
4	7/10/84	SLK	SHK	CONFORMING TO CONSTRUCTION RECORDS.
3	6-4-83	CJS	RLD	CHANGED COORDINATES ADDED STRUCTURE NUMBERS NOTE 2 AND DAMPERS
2	2/4/83	WB	RLD	ADDED ANCHORS AT P.I. NO.1
1	2/18/83	CJS	RLD	T6-TRANSMISSION LINE CONSTRUCTION CONTRACT
0	1/21/83	CJS	RLD	T5-CLEARING CONTRACT

REVISIONS




STONY BROOK ENERGY CENTER
MMWEC PHASE 1 PROJECT

R.W. BECK and ASSOCIATES
ANALYTICAL AND CONSULTING ENGINEERS

Seattle, Washington Denver, Colorado

345KV. TRANSMISSION LINE NO.334
PLAN AND PROFILE (SHEET 1)
(INCLUDES STRUCTURES 1-4)

Des. RLD	Eng. RLD	Appr. RLD	Appr. Des.	Appr. Des.	Appr. Des.
Des. C/S	Chg. RLD	Appr.			
File. MC-8244-ET2-4A	Date. 1-31-80				2296-C-201.4



NOTE:
POTENTIAL NHESP SENSITIVE HABITAT
BETWEEN 30+29 AND STRUCTURE #5

WESTOVER METROPOLITAN DEVELOPMENT CORPORATION

MERIDIAN OF
THE WESTOVER COORDINATE
SYSTEM WHICH EQUALS THAT
OF THE MASSACHUSETTS COORDINATE
SYSTEM, MAINLAND ZONE


NOTES

1. PROVIDE SIGN READING "DANGER-KEEP AWAY".

CONTINENTAL FIELD SERVICE CORPORATION
ELMSFORD, NEW YORK

No.	Date	By	Chd	Revision
7	1/17/12	JPH	MD	ADDED NHESP NOTE
6	3/1/11	YL	JP	CONVERTED TO CAD
5	7/10/11	SLK		CONFIRMING TO CONSTRUCTION RECORDS,
4	4/17/11	SLK		ADDED "DANGER KEEP AWAY" SIGNS
3	5-4-11	CJS	RLD	ADDED STRUCTURE NUMBERS
2	3/1/11	WB	RLD	ADDED ANCHORS AT PI No.3
1	1/15/11	CJS	RLD	T6-TRANSMISSION LINE CONSTRUCTION CONTRACT
0	1/21/11	CJS	RLD	T5-CLEARING CONTRACT

REVISIONS



MMWEC
A PUBLIC CORPORATION
STONY BROOK ENERGY CENTER
MMWEC PHASE 1 PROJECT


R.W. BECK and ASSOCIATES
ANALYTICAL AND CONSULTING ENGINEERS
Seattle, Washington Denver, Colorado

345KV. TRANSMISSION LINE NO.334
PLAN AND PROFILE (SHEET 2)
(INCLUDES STRUCTURES 5-9)

Des. C/S	Eng. RLD	App. C/S	App. RLD	App. Date	App. Desc.
				1-31-80	2296-C-202.5

NOTES

1. PROVIDE SIGN READING "DANGER-KEEP AWAY".



CONTINENTAL FIELD SERVICE CORPORATION
ELMSFORD, NEW YORK

No.	Date	By	Clad	Revision
7	5/11	JPH	MD	ADDED WEST ST. AND WETLAND
6	3/1/00	YL	JP	CONVERTED TO CAD
5	7/10/00	SLK	SLK	CONFORMING TO CONSTRUCTION RECORDS.
4	6/17/00	SLK	RLD	ADDED "DANGER KEEP AWAY" SIGNS.
3	5-8-00	CJS	RLD	ADDED STRUCTURE NUMBERS AND DAMPERS
2	3/1/00	W.B	RLD	ADDED ANCHORS AT P.I. NO 4 AND P.I. NO 5
1	1/13/00	CJS	RLD	T5-TRANSMISSION LINE CONSTRUCTION CONTRACT
0	1/13/00	CJS	RLD	T5-CLEARING CONTRACT

REVISIONS

MMWEC
Massachusetts Municipal Wholesale Electric Company
A PUBLIC CORPORATION

STONY BROOK ENERGY CENTER
MMWEC PHASE 1 PROJECT

R.W. BECK and ASSOCIATES
ANALYTICAL AND CONSULTING ENGINEERS
Seattle, Washington Denver, Colorado

345KV. TRANSMISSION LINE NO.334
PLAN AND PROFILE (SHEET 3)
(INCLUDES STRUCTURES 10-14)

Des. RLD	Eng. RLD	Appr. RLD	Appr. RLD	Appr. RLD
Des. C/S	Eng. C/S	Appr. C/S	Appr. C/S	Appr. C/S
File: MC-6244-ET2-4A	Date: 1-31-00	2296-C-203.5		

NOTES
 1. FIELD VERIFY ALL ANGLES TO LANDMARKS, DISTANCES FROM C CONSTRUCTION TO RIGHT OF WAY LIMITS, AND ALL PROFILE ELEVATIONS BETWEEN P.I.6 AND P.I.8

CONTINENTAL FIELD SERVICE CORPORATION
 ELMSFORD, NEW YORK

No.	Date	By	Revised	Revision
6	6/11	JPH	MD	ADDED WETLAND
5	6/24/04	YL	JP	CONVERTED TO CAD
4	7/14/04	SLK	SLK	CONFORMING TO CONSTRUCTION RECORDS.
3	7-1-03	CJS	RLD	ADDED NOTE NO.1 REMOVED R-O-W.DIM.
2	9-4-03	CJS	RLD	ADDED STRUCTURE NUMBERS AND DAMPERS
1	2/14/03	CJS	RLD	T6-TRANSMISSION LINE CONSTRUCTION CONTRACT
0	1/21/03	CJS	RLD	T5-CLEARING CONTRACT

REVISION


MMWEC
 Massachusetts Municipal Wholesale Electric Company
 A PUBLIC CORPORATION

STONY BROOK ENERGY CENTER
 MMWEC PHASE 1 PROJECT

R.W.BECK and ASSOCIATES
 ANALYTICAL AND CONSULTING ENGINEERS
 Seattle, Washington Denver, Colorado


345KV. TRANSMISSION LINE NO.334
 PLAN AND PROFILE (SHEET 4)
 (INCLUDES STRUCTURES 15-18)

Des. RLD	Eng. RLD	Appr. Eng. Des.	Appr. Const. Des.	Appr. Const. Des.
Des. C/S	Des. RLD	Appr.		
File: MC-6244-E2-4A	Date: 1-31-00	2296-C-204.4		



NOTES

1. FIELD VERIFY ALL ANGLES TO LANDMARKS, DISTANCES FROM @ CONSTRUCTION TO RIGHT OF WAY LIMITS, AND ALL PROFILE ELEVATIONS BETWEEN P.I.B AND P.L.B
2. FOR ST. 1 STRUCTURES LOCATE TWO ARMS NO. OF THE @ CONSTRUCTION AND ONE ARMS NO. OF THE @ CONSTRUCTION.
3. PROVIDED "DANGER-KEEP AWAY" SIGNS.



CONTINENTAL FIELD SERVICE CORPORATION
ELMSFORD, NEW YORK

No.	Date	By	Chd.	Revision
7	4/11	JPH	MD	ADDED WETLAND, POTENTIAL WELL AND AGR.
6	2/28/94	YL	JP	CONVERTED TO CAD
5	7/10/94	SLK	SLK	CONFORMING TO CONSTRUCTION RECORDS
4	6/29/94	SLK		ADDED DANGER KEEP AWAY SIGNS
3	7-1-92	CJS	RLD	ADDED NOTE NO.1 & NO.2 REMOVED R-0-W, DIM.
2	8-8-89	CJS	RLD	ADDED STRUCTURE NUMBERS AND DAMPERS
1	2/18/94	CJS	RLD	T6-TRANSMISSION LINE CONSTRUCTION CONTRACT
0	1/21/94	CJS	RLD	T5-CLEARING CONTRACT

REVISIONS

MMWEC
Massachusetts Municipal Wholesale Electric Company
A PUBLIC CORPORATION

STONY BROOK ENERGY CENTER
MMWEC PHASE 1 PROJECT


R.W. BECK and ASSOCIATES
ANALYTICAL AND CONSULTING ENGINEERS
Seattle, Washington Denver, Colorado

345KV. TRANSMISSION LINE NO.334
PLAN AND PROFILE (SHEET 5)
(INCLUDES STRUCTURES 19-22)

Des. RLD	Eng. RLD	Appr. Eng. Dist.	Appr. Mach. Dist.	Appr. Survey. Dist.
Des. C/S	Chd. RLD	Appr.		
File. MC-6244-E72-MA	Date. 1-31-80	2296-C-205.5		

NOTES

1. ALL STATIONING IS TO THE ϵ OF SURVEY. WHERE ϵ OF SURVEY AND ϵ OF CONSTRUCTION DO NOT COINCIDE AN OFFSET DIMENSION FROM ϵ OF SURVEY WILL BE PROVIDED.
2. FIELD VERIFY ALL OFFSET DIMENSIONS FOR TANGENT AND ANGLE POLES.



CONTINENTAL FIELD SERVICE CORPORATION
ELMSFORD, NEW YORK

No.	Date	By	Clad	Revision
6	8/11	JPH	MD	ADDED WETLAND
5	2/28/08	YL	JP	CONVERTED TO CAD
4	7/10/07	SLK	SLK	CONFORMING TO CONSTRUCTION RECORDS.
3	8-8-03	CJS	RLD	ADDED STRUCTURE NUMBERS
2	2/4/03	W.B	RLD	ADDED ANCHORS AT P.I.No 9A
1	2/18/03	CJS	RLD	T5-TRANSMISSION LINE CONSTRUCTION CONTRACT
0	1/21/03	CJS	RLD	T5-CLEARING CONTRACT

REVISIONS

MMWEC
Massachusetts Municipal Wholesale Electric Company
A PUBLIC CORPORATION

STONY BROOK ENERGY CENTER
MMWEC PHASE 1 PROJECT


R.W.BECK and ASSOCIATES
ANALYTICAL AND CONSULTING ENGINEERS
Seattle, Washington Denver, Colorado

345KV. TRANSMISSION LINE NO.334
PLAN AND PROFILE (SHEET 8)
(INCLUDES STRUCTURES 32-36)

Des. RLD	Eng. RLD	Appr. Eng. Dist.	Appr. Eng. Dist.	Appr. Eng. Dist.
Des. C/S	Clad. RLD	Appr.	Appr.	Appr.
File: MC-6244-ET2-MA			Date: 1-31-00	2296-C-208.4

MERIDIAN OF THE WESTOVER COORDINATE SYSTEM WHICH EQUELS THAT OF THE MASSACHUSETTS COORDINATE SYSTEM, MAINLAND ZONE

MUNSING



NOTES

1. ALL STATIONING IS TO THE C OF SURVEY. WHERE C OF SURVEY AND C OF CONSTRUCTION DO NOT COINCIDE AN OFFSET DIMENSION FROM C OF SURVEY WILL BE PROVIDED.
2. FIELD VERIFY ALL OFFSET DIMENSIONS FOR ALL TANGENT AND ANGLE POLES.

CONTINENTAL FIELD SERVICE CORPORATION
ELMSFORD, NEW YORK.

6	1/11	JPH	MD	ADDED WETLAND TEXT
5	1/11/00	YL	JP	CONVERTED TO CAD
4	7/14/01	SLK	SLK	CONFORMING TO CONSTRUCTION RECORDS.
3	1/14/02	CJS	RLD	REV PROFILE STA. 258+00 TO 267+00. REV STR. 39, 40 & 41. REV SAG CURVES
2	3-2-02	CJS	RLD	ADDED STRUCTURE NUMBERS AND DAMPERS
1	1/14/02	CJS	RLD	T6-TRANSMISSION LINE CONSTRUCTION CONTRACT
0	1/14/02	CJS	RLD	T5-CLEARING CONTRACT
No.	Date	By	Clad	Revision

REVISIONS



STONY BROOK ENERGY CENTER
MMWEC PHASE 1 PROJECT




R.W. BECK and ASSOCIATES
ANALYTICAL AND COMPUTING ENGINEERS

Seattle, Washington Denver, Colorado

345KV. TRANSMISSION LINE NO.334
PLAN AND PROFILE (SHEET 9)
(INCLUDES STRUCTURES 37-41)

Des. RLD	Eng. RLD	Appr. Dist.	Appr. Dist.	Appr. Dist.
Des. C/S	Chk. RLD	Appr.	Appr.	Appr.
File. MC-8244-E12-MA	Date. 1-31-00	2296-C-209.4		

THE MERIDIAN OF THE WESTOVER COORDINATE SYSTEM WHICH EQUALS THAT OF THE MASSACHUSETTS COORDINATE SYSTEM, MAINLAND ZONE



CONTINENTAL FIELD SERVICE CORPORATION
ELMSFORD, NEW YORK

No.	Date	By	Clad	Revision
5	3/1/88	YL	JP	CONVERTED TO CAD
4	7/16/88	SLK	SLK	CONFORMING TO CONSTRUCTION RECORDS.
3	8/2/88	CJS	RLD	REV. POLES @ STR. 42
2	8-8-88	CJS	RLD	ADDED STRUCTURE NUMBERS
1	2/18/88	CJS	RLD	TS-TRANSMISSION LINE CONSTRUCTION CONTRACT
0	1/31/88	CJS	RLD	TS-CLEARING CONTRACT

REVISIONS



STONY BROOK ENERGY CENTER
MMWEC PHASE 1 PROJECT

R.W. BECK and ASSOCIATES
ANALYTICAL AND CONSULTING ENGINEERS
Seattle, Washington Denver, Colorado




345KV. TRANSMISSION LINE NO. 334
PLAN AND PROFILE (SHEET 10)
(INCLUDES STRUCTURE 42)

Des.	Eng.	Appr.	Appr.	Appr.
C/S	Chd.	Plan.	Dist.	Dist.
RLD	RLD	RLD	RLD	RLD
MO-8244-ET2-MA	1-31-88	2296-C-210.4		

YOP
MAPS
HANCOCK
AND
LANESBOROUGH

Legend

- Estimated Tower Locations
- Estimated Access Road
- MA Towns



LOCUS MAP

Berkshire Wind Power Brodie Mountain Road Hancock and Lanesborough, MA

USGS TOPOGRAPHIC QUADRANGLE MAP
HANCOCK, MASSACHUSETTS, 1988

Project No:	15.0166117.00
Drawn by:	ERH
Checked by:	DMN
Date:	6/3/2011
Figure No:	1

**CHEMICAL
MATERIAL
SAFETY
DATA
SHEETS**

MATERIAL SAFETY DATA SHEET



Emergency Phone: 800-992-5994
Dow AgroSciences LLC
Indianapolis, IN 46268

RODEO* HERBICIDE

Effective Date: 3/23/04
Product Code: 84825
MSDS: 006694

1. PRODUCT AND COMPANY IDENTIFICATION:

PRODUCT: Rodeo* Herbicide

COMPANY IDENTIFICATION:

Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268-1189

2. COMPOSITION/INFORMATION ON INGREDIENTS:

Glyphosate IPA: N-(phosphono-methyl) glycine, Isopropylamine Salt	CAS # 038641-94-0	53.8%
Balance, Total		46.2%

3. HAZARDOUS IDENTIFICATIONS:

EMERGENCY OVERVIEW

Clear, pale yellow liquid. May cause eye irritation. Slightly toxic to aquatic organisms.

EMERGENCY PHONE NUMBER: 800-992-5994

4. FIRST AID:

EYE: Flush eyes thoroughly with water for several minutes. Remove contact lenses after initial 1-2 minutes and continue flushing for several additional minutes. If effects occur, consult a physician, preferably an ophthalmologist.

SKIN: Wash skin with plenty of water.

INGESTION: No emergency medical treatment necessary.

INHALATION: Remove person to fresh air; if effects occur, consult a physician.

NOTE TO PHYSICIAN: No specific antidote. Treatment of exposure should be directed at the control of symptoms and the clinical condition of the patient.

5. FIRE FIGHTING MEASURES:

FLASH POINT: >214°F (>101°C)

METHOD USED: Setaflash

FLAMMABLE LIMITS:

LFL: Not applicable

UFL: Not applicable

EXTINGUISHING MEDIA: Foam, CO₂, Dry Chemical

FIRE AND EXPLOSION HAZARDS: Foam fire extinguishing system is preferred because uncontrolled water can spread possible contamination. Toxic irritating gases may be formed under fire conditions.

FIRE-FIGHTING EQUIPMENT: Use positive-pressure, self-contained breathing apparatus and full protective equipment.

6. ACCIDENTAL RELEASE MEASURES:

ACTION TO TAKE FOR SPILLS: Absorb small spills with an inert absorbent material such as Hazorb, Zorbball, sand, or dirt. Report large spills to Dow AgroSciences on 800-992-5994.

7. HANDLING AND STORAGE:

PRECAUTIONS TO BE TAKEN IN HANDLING AND STORAGE: Keep out of reach of children. Do not swallow. Avoid contact with eyes, skin, and clothing. Avoid breathing vapors and spray mist. Handle concentrate in ventilated area. Wash thoroughly with soap and water after handling and before eating, chewing gum, using tobacco, using the toilet or smoking. Keep away from food, feedstuffs, and water supplies. Store in original container with the lid tightly closed. Store above 10°F (-12°C) to keep from crystallizing.

8. EXPOSURE CONTROLS/PERSONAL PROTECTION:

These precautions are suggested for conditions where the potential for exposure exists. Emergency conditions may require additional precautions.

EXPOSURE GUIDELINES: None established

ENGINEERING CONTROLS: Good general ventilation should be sufficient for most conditions. Local exhaust ventilation may be necessary for some operations.

RECOMMENDATIONS FOR MANUFACTURING, COMMERCIAL BLENDING, AND PACKAGING WORKERS:

EYE/FACE PROTECTION: Use safety glasses.

SKIN PROTECTION: No precautions other than clean body-covering clothing should be needed.

MATERIAL SAFETY DATA SHEET



Emergency Phone: 800-992-5994
Dow AgroSciences LLC
Indianapolis, IN 46268

Effective Date: 3/23/04
Product Code: 84825
MSDS: 006694

RODEO* HERBICIDE

RESPIRATORY PROTECTION: For most conditions, no respiratory protection should be needed; however, if discomfort is experienced, use a NIOSH approved air-purifying respirator.

APPLICATIONS AND ALL OTHER HANDLERS: Please refer to the product label for personal protective clothing and equipment.

9. PHYSICAL AND CHEMICAL PROPERTIES:

APPEARANCE: Clear, pale yellow liquid

DENSITY: 10.0 - 10.5 lbs/gal

pH: 4.8 - 5.0

ODOR: None

SOLUBILITY IN WATER: Miscible

SPECIFIC GRAVITY: 1.21 gm/L

FREEZING POINT: -7°F - -10°F (-21°C - -25°C)

10. STABILITY AND REACTIVITY:

STABILITY: (CONDITIONS TO AVOID) Stable under normal storage conditions.

INCOMPATIBILITY: (SPECIFIC MATERIALS TO AVOID) Galvanized or unlined steel (except stainless steel) containers or spray tanks may produce hydrogen gas which may form a highly combustible gas mixture.

HAZARDOUS DECOMPOSITION PRODUCTS: None known.

HAZARDOUS POLYMERIZATION: Not known to occur.

11. TOXICOLOGICAL INFORMATION:

EYE: May cause slight temporary eye irritation. Corneal injury is unlikely.

SKIN: Essentially non-irritating to skin. Prolonged skin contact is unlikely to result in absorption of harmful amounts. The LD₅₀ for skin absorption in rabbits is >5000 mg/kg. Did not cause allergic skin reactions when tested in guinea pigs.

INGESTION: Very low toxicity if swallowed. Harmful effects not anticipated from swallowing small amounts. The oral LD₅₀ for rats is >5000 mg/kg.

INHALATION: Brief exposure (minutes) is not likely to cause adverse effects. The aerosol LC₅₀ for rats is >6.37 mg/L for 4 hours.

SYSTEMIC (OTHER TARGET ORGAN) EFFECTS: For a similar material, glyphosate, in animals, effects have been reported on the following organ: liver.

CANCER INFORMATION: A similar material, glyphosate, did not cause cancer in laboratory animals.

TERATOLOGY (BIRTH DEFECTS): For glyphosate IPA, available data are inadequate for evaluation of potential to cause birth defects.

REPRODUCTIVE EFFECTS: For glyphosate IPA, available data are inadequate to determine effects on reproduction.

MUTAGENICITY: For a similar material, glyphosate, in-vitro and animal genetic toxicity studies were negative.

12. ECOLOGICAL INFORMATION:

ENVIRONMENTAL DATA:

ECOTOXICOLOGY:

Material is practically non-toxic to aquatic organisms on an acute basis (LC₅₀ or EC₅₀ is >100 mg/L in most sensitive species tested).

Acute LC₅₀ for rainbow trout (*Oncorhynchus mykiss*) is >2500 mg/L.

Acute immobilization EC₅₀ in water flea (*Daphnia magna*) is 918 mg/L.

Material is practically non-toxic to birds on an acute basis (LD₅₀ is >2000 mg/kg).

Acute oral LD₅₀ in bobwhite (*Colinus virginianus*) is >2000 mg/kg.

The LC₅₀ in earthworm *Eisenia foetida* is >1000 mg/kg.

Acute contact LD₅₀ in honey bee (*Apis mellifera*) is >100 µg/bee.

Acute oral LD₅₀ in honey bee (*Apis mellifera*) is >100 µg/bee.

Growth inhibition EC₅₀ in green alga (*Selenastrum capricornutum*) is 127 mg/L.

Growth inhibition EC₅₀ in duckweed (*Lemna sp.*) is 24.4 mg/L.

13. DISPOSAL CONSIDERATIONS:

DISPOSAL METHOD: If wastes and/or containers cannot be disposed of according to the product label directions, disposal of this material must be in accordance with your local or area regulatory authorities.

MATERIAL SAFETY DATA SHEET

Emergency Phone: 800-992-5994
Dow AgroSciences LLC
Indianapolis, IN 46268

Effective Date: 3/23/04
Product Code: 84825
MSDS: 006694



RODEO* HERBICIDE

This information presented below only applies to the material as supplied. The identification based on characteristic(s) or listing may not apply if the material has been used or otherwise contaminated. It is the responsibility of the waste generator to determine the toxicity and physical properties of the material generated to determine the proper waste identification and disposal methods in compliance with applicable regulations.

If the material as supplied becomes a waste, follow all applicable regional, national and local laws and regulations.

14. TRANSPORT INFORMATION:

U.S. DEPARTMENT OF TRANSPORTATION (DOT) INFORMATION:

For all package sizes and modes of transportation:
This material is not regulated for transport.

15. REGULATORY INFORMATION:

NOTICE: The information herein is presented in good faith and believed to be accurate as of the effective date shown above. However, no warranty, express or implied, is given. Regulatory requirements are subject to change and may differ from one location to another; it is the buyer's responsibility to ensure that its activities comply with federal, state or provincial, and local laws. The following specific information is made for the purpose of complying with numerous federal, state or provincial, and local laws and regulations.

U.S. REGULATIONS

SARA 313 INFORMATION: To the best of our knowledge, this product contains no chemical subject to SARA Title III Section 313 supplier notification requirements.

SARA HAZARD CATEGORY: This product has been reviewed according to the EPA "Hazard Categories" promulgated under Sections 311 and 312 of the Superfund Amendment and Reauthorization Act of 1986 (SARA Title III) and is considered, under applicable definitions, to meet the following categories:

Not to have met any hazard category

TOXIC SUBSTANCES CONTROL ACT (TSCA): All ingredients are on the TSCA inventory or are not required to be listed on the TSCA inventory.

STATE RIGHT-TO-KNOW: This product is not known to contain any substances subject to the disclosure requirements of

New Jersey
Pennsylvania

OSHA HAZARD COMMUNICATION STANDARD: This product is a "Hazardous Chemical" as defined by the OSHA Hazard Communication Standard, 29 CFR 1910.1200.

COMPREHENSIVE ENVIRONMENTAL RESPONSE COMPENSATION AND LIABILITY ACT (CERCLA, or SUPERFUND): To the best of our knowledge, this product contains no chemical subject to reporting under CERCLA.

NATIONAL FIRE PROTECTION ASSOCIATION (NFPA) RATINGS:

<u>CATEGORY</u>	<u>RATING</u>
Health	1
Flammability	1
Reactivity	0

16. OTHER INFORMATION:

MSDS STATUS: Revised Sections: 3,4,11,12,13,14 & 15
Reference: DR-0361-8028
Replaces MSDS Dated: 1/12/00
Document Code: D03-148-002
Replaces Document Code: D03-148-001

The Information Herein Is Given In Good Faith, But No Warranty, Express Or Implied, Is Made. Consult Dow AgroSciences For Further Information.


DuPont™ Escort® XP Herbicide

Version 2.0

Revision Date 01/22/2010

Ref. 130000036195

This SDS adheres to the standards and regulatory requirements of the United States and may not meet the regulatory requirements in other countries.

SECTION 1. PRODUCT AND COMPANY IDENTIFICATION

Product name : DuPont™ Escort® XP Herbicide
 Tradename/Synonym : DPX-T6376 60 XP
 Metsulfuron Methyl 60 XP
 Escort 60 DF
 B11495142
 METSULFURON METHYL (Methyl 2-[[[(4-methoxy-6-methyl-1,3,4-triazin-2-yl)amino]carbonyl]amino]sulfonyl]benzoate)

MSDS Number : 130000036195
 Manufacturer : DuPont
 1007 Market Street
 Wilmington, DE 19898

Product Information : 1-800-441-7515 (outside the U.S. 1-302-774-1000)
 Medical Emergency : 1-800-441-3637 (outside the U.S. 1-302-774-1139)
 Transport Emergency : CHEMTREC: 1-800-424-9300 (outside the U.S. 1-703-527-3887)

SECTION 2. HAZARDS IDENTIFICATION
Emergency Overview
CAUTION!

Causes eye irritation. Avoid contact with skin, eyes and clothing. Avoid breathing dust. Avoid breathing vapours or mist.

Potential Health Effects

This section includes potential acute adverse effects which could occur if this material is not used according to the label.

Skin : May cause: Irritation with discomfort or pain, redness or rash, itching or swelling.

Eyes : May cause: Irritation with discomfort, pain, redness, or visual impairment.

Carcinogenicity

None of the components present in this material at concentrations equal to or greater than 0.1% are listed by IARC, NTP, or OSHA, as a carcinogen.

SECTION 3. COMPOSITION/INFORMATION ON INGREDIENTS

Component	CAS-No.	Concentration
Metsulfuron methyl	74223-64-6	60 %
Other Ingredients		40 %

**DuPont™ Escort® XP Herbicide**

Version 2.0

Revision Date 01/22/2010

Ref. 130000036195

SECTION 4. FIRST AID MEASURES

- Skin contact : Take off all contaminated clothing immediately. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
- Eye contact : Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
- Inhalation : No specific intervention is indicated as the compound is not likely to be hazardous. Consult a physician if necessary.
- Ingestion : No specific intervention is indicated as the compound is not likely to be hazardous. Consult a physician if necessary.
- General advice : Have the product container or label with you when calling a poison control center or doctor, or going for treatment.
For medical emergencies involving this product, call toll free 1-800-441-3637. See Label for Additional Precautions and Directions for Use.

SECTION 5. FIRE-FIGHTING MEASURES

- Flammable Properties
Flash point : not applicable
- Suitable extinguishing media : Water spray, Dry chemical, Foam, Carbon dioxide (CO2)
- Unsuitable extinguishing media : High volume water jet, (contamination risk)
- Firefighting Instructions : In the event of fire, wear self-contained breathing apparatus. Wear full protective equipment. (on small fires) If area is heavily exposed to fire and if conditions permit, let fire burn itself out since water may increase the area contaminated. Cool containers / tanks with water spray.

SECTION 6. ACCIDENTAL RELEASE MEASURES

NOTE: Review FIRE FIGHTING MEASURES and HANDLING (PERSONNEL) sections before proceeding with clean-up. Use appropriate PERSONAL PROTECTIVE EQUIPMENT during clean-up.

- Safeguards (Personnel) : Wear personal protective equipment. Refer to protective measures listed in sections 7 and 8.
- Spill Cleanup : Clean-up methods - small spillage Sweep up or vacuum up spillage and collect in suitable container for disposal. Clean-up methods - large spillage Prevent further leakage or spillage. Use approved industrial vacuum cleaner for removal. Shovel into suitable container for disposal. If spill area is on



DuPont™ Escort® XP Herbicide

Version 2.0

Revision Date 01/22/2010

Ref. 130000036195

ground near valuable plants or trees, remove 5 cm of top soil after initial clean-up.

Accidental Release Measures : Prevent material from entering sewers, waterways, or low areas. Never return spills in original containers for re-use. Dispose of in accordance with local regulations.

SECTION 7. HANDLING AND STORAGE

- Handling (Personnel) : Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Handling (Physical Aspects) : Dust may form explosive mixture in air. Keep away from heat and sources of ignition.
- Storage : Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Store product in original container only, away from other pesticides, fertilizer, food or feed. Keep containers tightly closed in a cool, well-ventilated place. Keep out of the reach of children.

SECTION 8. EXPOSURE CONTROLS/PERSONAL PROTECTION

- Personal protective equipment
 - Skin and body protection : Applicators and other handlers must wear:
Long sleeved shirt and long pants
Shoes plus socks
PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:
Coveralls
Shoes plus socks
- Protective measures : Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Exposure Guidelines
Exposure Limit Values
Metsulfuron methyl
AEL * (DUPONT) 10 mg/m3 8 & 12 hr. TWA

* AEL is DuPont's Acceptable Exposure Limit. Where governmentally imposed occupational exposure limits which are lower than the AEL are in effect, such limits shall take precedence.

SECTION 9. PHYSICAL AND CHEMICAL PROPERTIES


DuPont™ Escort® XP Herbicide

Version 2.0

Revision Date 01/22/2010

Ref. 130000036195

Reproductive toxicity : Animal testing did not show any effects on fertility.

Teratogenicity : Animal testing showed no developmental toxicity.

SECTION 12. ECOLOGICAL INFORMATION

Aquatic Toxicity

Metsulfuron methyl

96 h LC50 : Oncorhynchus mykiss (rainbow trout) > 150 mg/l

96 h LC50 : Lepomis macrochirus (Bluegill sunfish) > 150 mg/l

Toxicity to other organisms

Metsulfuron methyl

LD50 : Anas platyrhynchos (Mallard duck) > 2,510 mg/kg

LC50 : Colinus virginianus (Bobwhite quail) > 5,620 mg/kg

Additional ecological information : Environmental Hazards: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

SECTION 13. DISPOSAL CONSIDERATIONS

Waste Disposal : Do not contaminate water, food or feed by disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Treatment, storage, transportation, and disposal must be in accordance with applicable federal, state/provincial, and local regulations.

Container Disposal : Refer to the product label for instructions.

SECTION 14. TRANSPORT INFORMATION

 IMDG UN-Number : 3077
 Proper shipping name : Environmentally hazardous substance, solid, n.o.s.
 (Metsulfuron methyl)
 Class : 9
 Packaging group : III
 Labelling No. : 9

Marine pollutant

Not regulated as a hazardous material by DOT.

Not regulated as a hazardous material by IATA.

Optional classification as per IATA Special Provision A97.



DuPont™ Escort® XP Herbicide

Version 2.0

Revision Date 01/22/2010

Ref. 13000036195

SECTION 15. REGULATORY INFORMATION

SARA 313 Regulated Chemical(s) : SARA 313: This material does not contain any chemical components with known CAS numbers that exceed the threshold (De Minimis) reporting levels established by SARA Title III, Section 313.

EPA Reg. No. : 352-439
 In the United States this product is regulated by the US Environmental Protection Agency (EPA) under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA). It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read and follow all label directions. This product is excluded from listing requirements under EPA/TSCA.

SECTION 16. OTHER INFORMATION

	NFPA	HMIS
Health :	1	1
Flammability :	1	1
Reactivity/Physical hazard :	0	0

MSDS preparation date : 01/22/2010

DuPont™, Escort® are trademarks of E. I. du Pont de Nemours and Company

Contact person : DuPont Crop Protection, Wilmington, DE, 19898, Phone: 1-888-638-7668

The information provided in this Safety Data Sheet is correct to the best of our knowledge, information and belief at the date of its publication. The information given is designed only as a guidance for safe handling, use, processing, storage, transportation, disposal and release and is not to be considered a warranty or quality specification. The information relates only to the specific material designated and may not be valid for such material used in combination with any other materials or in any process, unless specified in the text.

Significant change from previous version is denoted with a double bar.

CHEMICAL
TECHNICAL
DATA
SHEETS

Specimen Label



Rodeo®

Herbicide

For control of annual and perennial weeds and woody plants in forests, non-crop sites, and in and around aquatic sites; also for use in wildlife habitat areas, for perennial grass release, and grass growth suppression and grazed areas on these sites.

Avoid contact of herbicide with foliage, green stems, exposed non-woody roots or fruit of crops, desirable plants and trees, because severe injury or destruction may result.

Active Ingredient(s):	
glyphosate [†] N-(phosphonomethyl)glycine,	
isopropylamine salt	53.8%
Other Ingredients	46.2%
Total Ingredients	100.0%

[†] Contains 5.4 pounds per gallon glyphosate, isopropylamine salt (4 pounds per gallon glyphosate acid).

EPA Reg. No. 62719-324

Keep Out of Reach of Children

CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful If Inhaled

Avoid breathing spray mist. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE (Personal Protective Equipment). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

Do not contaminate water when cleaning equipment or disposing of equipment washwaters. Treatment of aquatic weeds can result in oxygen depletion or loss due to decomposition of dead plants. This oxygen loss can cause fish suffocation.

In case of leak or spill, soak up and remove to a landfill.

Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

Do not mix, store or apply this product or spray solutions of this product in galvanized steel or unlined steel (except stainless steel) containers or spray tanks. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas, which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Terms and Conditions of Use, Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

This is an end-use product. Dow AgroSciences does not intend and has not registered it for reformulation. See individual container label for repackaging limitations.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has dried.

Storage and Disposal

Do not contaminate water, food, feed or seed by storage or disposal.

Pesticide Storage: Store above 10°F (-12°C) to keep product from crystallizing. Crystals will settle to the bottom. If allowed to crystallize, place in a warm room 68°F (20°C) for several days to redissolve and roll or shake container or recirculate in mini-bulk containers to mix well before using.

Pesticide Disposal: Wastes resulting from use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Nonrefillable containers 5 gallons or less:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers larger than 5 gallons:

Container Reuse: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Nonrefillable containers larger than 5 gallons:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

General Information (How this product works)

This product is a water-soluble liquid, which mixes readily with water and nonionic surfactant to be applied as a foliar spray for the control or destruction of many herbaceous and woody plants. This product is intended for control of annual and perennial weeds and woody plants in forests, pine straw plantations, non-crop sites such as utility rights-of-way, and in and around aquatic sites; also for use in wildlife habitat areas, for perennial grass release, and grass growth suppression and grazed areas on these sites.

The active ingredient in this product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, 7 days or more on most perennial weeds, and 30 days or more on most woody plants. Extremely cool or cloudy weather following treatment may slow the activity of this product and delay visual effects of control. Visible effects include gradual wilting and yellowing of the plant which advances to complete browning of above-ground growth and deterioration of underground plant parts.

Unless otherwise directed on this label, delay application until vegetation has emerged and reached the stages described for control of such vegetation under the "Weeds Controlled" section of this label.

Unemerged plants arising from unattached underground rhizomes or root stocks of perennials or brush will not be affected by the spray and will continue to grow. For this reason best control of most perennial weeds or brush is obtained when treatment is made at late growth stages approaching maturity.

Always use the higher rate of this product and surfactant within the recommended range when vegetation is heavy or dense, when treating dense multi-canopied sites or woody vegetation or difficult-to-control herbaceous or woody plants.

Do not treat weeds, brush or trees under poor growing conditions such as drought stress, disease or insect damage, as reduced control may result. Reduced control of target vegetation may also occur if foliage is heavily covered with dust at the time of treatment.

Reduced control may result when applications are made to woody plants or weeds following site disturbance or plant top growth removal from grazing, mowing, logging or mechanical brush control. For best results, delay treatment of such areas until resprouting and foliar growth has restored the target vegetation to the recommended stage of growth for optimum herbicide exposure and control.

Rainfall or irrigation occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours after application may wash the product off the foliage and a repeat treatment may be required.

This product does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

Note: The maximum rates stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate or sulfosate containing products does not exceed the maximum use rates.

Grazing Restrictions: This product may be used to treat undesirable vegetation in utility rights-of-way that pass through pastures, rangeland, and forestry sites that are being grazed. For tank mix applications, comply with all restrictions appearing on the tank mix product label.

Except for lactating dairy animals there are no grazing restrictions following the labeled applications of this product.

- For lactating dairy animals there are no grazing restrictions for the following labeled applications of this product:
 - ▶ Where the spray can be directed onto undesirable woody brush and trees, such as in handgun spray-to-wet or low volume directed spray treatments.
 - ▶ For tree injection of frill applications and for cut stump treatments
- For broadcast applications, observe the following restrictions for lactating dairy animals:
 - ▶ For application rates of greater than 4.5 but not to exceed 7.5 quarts per acre, no more than 15 percent of the available grazing area may be treated.
 - ▶ For application rates that do not exceed 4.5 quarts per acre, no more than 25 percent of the available grazing area may be treated.
- These restrictions do not apply to pastures, rangeland or forestry sites outside of utility rights-of-way.

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. When not in use, keep container closed to prevent spills and contamination.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product or other materials that are not expressly recommended in this label. Mixing this product with herbicides or other materials not recommended in this label may result in reduced performance.

ATTENTION: Avoid drift. Extreme care must be used when applying this product to prevent injury to desirable plants and crops.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of plant or crop injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray drift to occur. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. **Avoid applying at excessive speed or pressure.**

Spray Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the following **Aerial Drift Reduction Advisory Information**:

Importance of Droplet Size: The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversion section of this label).

Controlling Droplet Size: Volume-Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows product larger droplets.

Pressure-Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of nozzles-Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation-Orienting nozzles so that the spray is released backwards, parallel to the airstream will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

Nozzle Type-Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.

Boom Length-For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application-Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment: When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

Wind: Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Temperature and Humidity: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions: Applications should not occur during a temperature inversion, because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a connected cloud (under low wind conditions) indicates an inversion, while smoke that moves upwards and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas: The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Mixing And Application Instructions

Apply these spray solutions in properly maintained and calibrated equipment capable of delivering desired volumes. Hand-gun applications should be properly directed to avoid spraying desirable plants. Note: reduced results may occur if water containing soil is used, such as water from ponds and unlined ditches.

Mixing

This product mixes readily with water. Mix spray solutions of this product as follows:

1. Fill the mixing or spray tank with the required amount of water while adding the required amount of this product (see "Directions for Use" and "Weeds Controlled" sections of this label).
2. Near the end of the filling process, add the required surfactant and mix well. Remove hose from tank immediately after filling to avoid siphoning back into the water source.

Note: If tank mixing with Garlon® 3A herbicide, ensure that Garlon 3A is well mixed with at least 75 percent of the total spray volume before adding this product to the spray tank to avoid incompatibility.

During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, place the filling hose below the surface of the spray solution (only during filling), terminate by-pass and return lines at the bottom of the tank, and, if needed, use an approved anti-foam or defoaming agent.

Keep by-pass line on or near bottom of tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh. Carefully select correct nozzle to avoid spraying a fine mist. For best results with conventional ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

IMPORTANT: When using this product, unless otherwise specified, mix with a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient. For conifer release (pine release) use only surfactants that are approved for conifer release, and specified on the surfactant label as safe for use in conifer release (pine release). Always read and follow the manufacturer's surfactant label recommendations for best results.

Colorants or marking dyes approved for use with herbicides may be added to spray mixtures of this product. Colorants or dyes used in spray solutions of this product may reduce performance, especially at lower rates or dilutions. Use colorants or dyes according to the manufacturer's label recommendations.

Clean sprayer and parts immediately after using this product by thoroughly flushing with water and dispose of rinsate according to labeled use or disposal instructions.

Carefully observe all cautionary statements and other information appearing in the surfactant label.

Application Equipment And Techniques

ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE EXERCISED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift, or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to crops, plants, or other areas on which the treatment was not intended. The likelihood of plant or crop injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray drift to occur. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. **AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.**

Note: Use of this product in a manner not consistent with this label may result in injury to persons, animals, or crops, or other unintended consequences. When not in use, keep container closed to prevent spills and contamination.

Aerial Equipment

For aerial application of this product in California, refer to Federal supplemental label for this product entitled "For Aerial Application in California Only". In California, aerial application may be made in aquatic sites and noncrop areas, including aquatic sites present in noncrop areas that are part of the intended treatment.

For control of weed or brush species listed in this label using aerial application equipment: For aerial broadcast application, unless otherwise specified, apply the rates of this product and surfactant recommended for broadcast application in a spray volume of 3 to 20 gallons of water per acre. See the "Weeds Controlled" section of this label for labeled annual and herbaceous weeds and woody plants and broadcast rate recommendations. Aerial applications of this product may only be made as specifically recommended in this label.

AVOID DRIFT. Do not apply during inversion conditions, when winds are gusty or under any other condition which will allow drift. Drift may cause damage to any vegetation contacted to which treatment is not intended. To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure.

Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing in the additive label. The use of a drift control agent for conifer and herbaceous release applications may result in conifer injury and is not recommended.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. **Prolonged exposure of this product to uncoated steel surfaces may result in corrosion and possible failure of the part. Landing gear are most susceptible.** The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion.

Ground Broadcast Equipment

For control of weed or brush species listed in this label using conventional boom equipment: For ground broadcast application, unless otherwise specified, apply the rates of this product and surfactant recommended for broadcast application in a spray volume of 3 to 30 gallons of water per acre. See the "Weeds Controlled" section of this label for labeled annual and herbaceous weeds and woody plants and broadcast rate recommendations. As density of vegetation increases, spray volume should be increased within the recommended range to ensure complete coverage. Carefully select correct nozzle to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

Forestry and Utility Rights-of-Way Sites: This product is recommended for broadcast applications using suitable ground equipment in forestry sites, utility sites, and utility rights-of way. Apply the recommended rates of this product and surfactant in a spray volume of 10 to 60 gallons per acre. Check for even distribution of spray droplets.

**Hand-Held and High-Volume Equipment
(Use Coarse Sprays Only)**

For control of weeds listed in this label using knapsack sprayers or high-volume spraying equipment utilizing handguns or other suitable nozzle arrangements:

High volume sprays: Prepare a 3/4 to 2 percent solution of this product in water, add a nonionic surfactant and apply to foliage of vegetation to be controlled. For specific rates of application and instructions for control of various annual and perennial weeds, see the "Weeds Controlled" section in this label.

Prepare the desired volume of spray solution by mixing the amount of this product in water, shown in the following table:

Spray Solution

Desired Volume	Amount of this product							
	3/4%	1%	1 1/4%	1 1/2%	2%	5%	8%	10%
1 gal	1 fl oz	1 1/3 fl oz	1 2/3 fl oz	2 fl oz	2 2/3 fl oz	6 1/2 fl oz	10 1/4 fl oz	12 3/4 fl oz
25 gal	1 1/2 pt	1 qt	1 1/4 qt	1 1/2 qt	2 qt	5 qt	2 gal	2.5 gal
100 gal	3 qt	1 gal	1 1/4 gal	1 1/2 gal	2 gal	5 gal	8 gal	10 gal

2 tablespoons = 1 fluid ounce

For use in knapsack sprayers, it is suggested that the recommended amount of this product be mixed with water in a larger container. Fill the knapsack sprayer with the mixed solution and add the correct amount of surfactant.

Selective Equipment

This product may be applied through shielded sprayers or wiper application equipment. This equipment may be used to selectively control undesirable vegetation without harming desirable vegetation.

Shielded sprayers direct the herbicide solution onto weeds while shielding desirable vegetation from the spray solution. Any recommended rate or tank mixture of this product may be used employing this equipment.

Wiper applicators physically wipe product directly onto undesirable vegetation. Care should be taken to avoid wiping desirable vegetation. Use a 33 to 100 percent solution of this product, diluted in water for wiper applications. Use a 33 percent solution for wick or gravity feed systems. Higher concentrations may be used in pressurized systems that are capable of handling thicker solutions. Addition of a nonionic surfactant at a rate of 10 percent by volume of total herbicide solution is recommended.

Applications should be made on a spray-to-wet basis. Spray coverage should be uniform and complete. Do not spray to point of runoff.

Low volume directed sprays: This product may be used as a 5 to 10 percent solution in low-volume directed sprays for spot treatment of trees and brush. This treatment method is most effective in areas where there is a low density of undesirable trees or brush. If a straight stream nozzle is used, start the application at the top of the targeted vegetation and spray from top to bottom in a lateral zig-zag motion. Ensure that at least 50 percent of the leaves are contacted by the spray solution. For flat fan and cone nozzles and with hand-directed mist blowers, mist the application over the foliage of the targeted vegetation. Small, open-branched trees need only be treated from one side. If the foliage is thick or there are multiple root sprouts, applications must be made from several sides to ensure adequate spray coverage.

Weeds Controlled

Annual Weeds

Apply to actively growing annual grasses and broadleaf weeds.

Allow at least 3 days after application before disturbing treated vegetation. After this period the weeds may be mowed, tilled or burned. See "Directions for Use," "General Information" and "Mixing and Application Instructions" for labeled uses and specific application instructions.

Broadcast Application Rates: For weeds less than 6 inches tall, use 1 1/2 pints of this product per acre plus a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient. If weeds are greater than 6 inches tall, use 2 1/2 pints of this product per acre plus a non-ionic surfactant containing 80% or greater active ingredient..

Hand-Held, High-Volume Application Rates: Use a 3/4 percent solution of this product in water plus a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient. Apply to foliage of vegetation to be controlled.

When applied as directed, this product plus a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient will control the following annual weeds:

Common Name	Scientific Name
Balsamapple [†]	<i>Momordica charantia</i>
Barley	<i>Hordeum vulgare</i>
Barnyardgrass	<i>Echinochloa crus-galli</i>
Bassia, fivehook	<i>Bassia hyssopifolia</i>
Bluegrass, annual	<i>Poa annua</i>
Bluegrass, bulbous	<i>Poa bulbosa</i>
Brome	<i>Bromus spp.</i>
Buttercup	<i>Ranunculus spp.</i>
Cheat	<i>Bromus secalinus</i>
Chickweed, mouseear	<i>Cerastium vulgatum</i>
Cocklebur	<i>Xanthium strumarium</i>
Corn, volunteer	<i>Zea mays</i>
Crabgrass	<i>Digitaria spp.</i>
Dwarf dandelion	<i>Krigia cespitosa</i>
Falseflax, smallseed	<i>Camelina microcarpa</i>
Fiddleneck	<i>Amsinckia spp.</i>
Flaxleaf fleabane	<i>Conyza bonariensis</i>
Fleabane	<i>Erigeron spp.</i>
Foxtail	<i>Setaria spp.</i>
Foxtail, Carolina	<i>Alopecurus carolinianus</i>
Groundsel, common	<i>Senecio vulgaris</i>
Horseweed/Marestail	<i>Conyza canadensis</i>
Kochia	<i>Kochia scoparia</i>
Lambsquarters, common	<i>Chenopodium album</i>
Lettuce, prickly	<i>Lactuca serriola</i>
Morningglory	<i>Ipomoea spp.</i>
Mustard, blue	<i>Chorispora tenella</i>
Mustard, tansy	<i>Descurainia pinnata</i>
Mustard, tumble	<i>Sisymbrium altissimum</i>
Mustard, wild	<i>Sinapis arvensis</i>
Oats, wild	<i>Avena fatua</i>
Panicum	<i>Panicum spp.</i>
Pennycress, field	<i>Thlaspi arvense</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pigweed, smooth	<i>Amaranthus hybridus</i>
Ragweed, common	<i>Ambrosia artemisiifolia</i>
Ragweed, giant	<i>Ambrosia trifida</i>
Rocket, London	<i>Sisymbrium irio</i>
Rye	<i>Secale cereale</i>
Ryegrass, Italian ^{**}	<i>Lolium multiflorum</i>
Sandbur, field	<i>Cenchrus spp.</i>
Shattercane	<i>Sorghum bicolor</i>
Shepherd's-purse	<i>Capsella bursa-pastoris</i>
Signalgrass, broadleaf	<i>Brachiaria platyphylla</i>
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>
Sowthistle, annual	<i>Sonchus oleraceus</i>
Spanishneedles ^{**}	<i>Bidens bipinnata</i>
Stinkgrass	<i>Eragrostis ciliaris</i>
Sunflower	<i>Helianthus annuus</i>
Thistle, Russian	<i>Salsola kali</i>
Spurry, umbrella	<i>Holosteum umbellatum</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Wheat	<i>Triticum aestivum</i>
Witchgrass	<i>Panicum capillare</i>

[†] Apply with hand-held equipment only.

^{**} Apply 3 pints of this product per acre.

Annual weeds will generally continue to germinate from seed throughout the growing season. Repeat treatments will be necessary to control later germinating weeds.

Perennial Weeds

Apply this product to control most vigorously growing perennial weeds. Unless otherwise directed, apply when target plants are actively growing and most have reached early head or early bud stage of growth. Unless otherwise directed, allow at least 7 days after application before disturbing vegetation.

NOTE: If weeds have been mowed or tilled, do not treat until regrowth has reached the recommended stages. Fall treatments must be applied before a killing frost.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed.

Specific Weed Control Recommendations: For perennial weeds, apply the recommended rate plus a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient. Use of this product without surfactant will result in reduced herbicide performance. Refer to the "Mixing and Application Instructions" section of this label and the surfactant manufacturer label for more information.

When applied as directed, this product plus a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient will control the following perennial weeds: (Numbers in parentheses "(-)" following common name of a listed weed species refer to "Specific Perennial Weed Control Recommendations" for that weed which follow the species listing.)

Common Name	Scientific Name
Alfalfa (31)	<i>Medicago sativa</i>
Alligatorweed [†] (1)	<i>Alternanthera philoxeroides</i>
Anise/Fennel (31)	<i>Foeniculum vulgare</i>
Artichoke, Jerusalem (31)	<i>Helianthus tuberosus</i>
Bahiagrass (31)	<i>Paspalum notatum</i>
Bermudagrass (2)	<i>Cynodon dactylon</i>
Bindweed, field (3)	<i>Convolvulus arvensis</i>
Bluegrass, Kentucky (12)	<i>Poa pratensis</i>
Blueweed, Texas (3)	<i>Helianthus ciliaris</i>
Brackenfern (4)	<i>Pteridium spp.</i>
Bromegrass, smooth (12)	<i>Bromus inermis</i>
Canarygrass, reed (12)	<i>Phalaris arundinacea</i>
Cattail (5)	<i>Typha spp.</i>
Clover, red (31)	<i>Trifolium pratense</i>
Clover, white (31)	<i>Trifolium repens</i>
Cogongrass (6)	<i>Imperata cylindrica</i>
Cordgrass (7)	<i>Spartina spp.</i>
Cutgrass, giant [†] (8)	<i>Zizaniopsis miliacea</i>
Dallisgrass (31)	<i>Paspalum dilatatum</i>
Dandelion (31)	<i>Taraxacum officinale</i>
Dock, curly (31)	<i>Rumex crispus</i>
Dogbane, hemp (9)	<i>Apocynum cannabinum</i>
Fescue (31)	<i>Festuca spp.</i>
Fescue, tall (10)	<i>Mesembryanthemum crystallinum</i>
Guineagrass (11)	<i>Panicum maximum</i>
Hemlock, poison (31)	<i>Conium maculatum</i>
Horsenettle (31)	<i>Solanum carolinense</i>
Horseradish (9)	<i>Armoracia rusticana</i>
Ice Plant (22)	<i>Mesembryanthemum crystallinum</i>
Johnsongrass (12)	<i>Sorghum halepense</i>
Kikuyugrass (21)	<i>Pennisetum clandestinum</i>

Common Name	Scientific Name
Knapweed (9)	<i>Centaurea repens</i>
Lantana (13)	<i>Lantana camara</i>
Lespedeza, common (31)	<i>Lespedeza striata</i>
Lespedeza, sericea (31)	<i>Lespedeza cuneata</i>
Loosestrife, purple (14)	<i>Lythrum salicaria</i>
Lotus, American (15)	<i>Nelumbo lutea</i>
Maidencane (16)	<i>Panicum hematomon</i>
Milkweed (17)	<i>Asclepias spp.</i>
Muhly, wirestem (21)	<i>Muhlenbergia frondosa</i>
Mullein, common (31)	<i>Verbascum thapsus</i>
Napiergrass (31)	<i>Pennisetum purpureum</i>
Nightshade, silverleaf (3)	<i>Solanum elaeagnifolium</i>
Nutsedge, purple (18)	<i>Cyperus rotundus</i>
Nutsedge, yellow (18)	<i>Cyperus esculentus</i>
Orchardgrass (12)	<i>Dactylis glomerata</i>
Pampasgrass (19)	<i>Cortaderia jubata</i>
Paragrass (16)	<i>Brachiaria mutica</i>
Phragmites ^{††} (20)	<i>Phragmites spp.</i>
Quackgrass (21)	<i>Agropyron repens</i>
Reed, giant (22)	<i>Arundo donax</i>
Ryegrass, perennial (12)	<i>Lolium perenne</i>
Smartweed, swamp (31)	<i>Polygonum coccineum</i>
Spatdock (23)	<i>Nuphar luteum</i>
Starthistle, yellow (31)	<i>Centaurea solstitialis</i>
Sweet potato, wild [†] (24)	<i>Ipomoea pandurata</i>
Thistle, artichoke (25)	<i>Cynara cardunculus</i>
Thistle, Canada (25)	<i>Cirsium arvense</i>
Timothy (12)	<i>Phleum pratense</i>
Torpedograss [†] (26)	<i>Panicum repens</i>
Tules, common (27)	<i>Scirpus acutus</i>
Vaseygrass (31)	<i>Paspalum urvillei</i>
Velvetgrass (31)	<i>Holcus spp.</i>
Waterhyacinth (28)	<i>Eichornia crassipes</i>
Waterlettuce (29)	<i>Pistia stratiotes</i>
Waterprimrose (30)	<i>Ludwigia spp.</i>
Wheatgrass, western (12)	<i>Agropyron smithii</i>

[†] Partial control.

^{††} Partial control in southeastern states. See "Specific Weed Control Recommendations" below.

Specific Perennial Weed Control Recommendations:

- Alligatorweed:** Apply 6 pints of this product per acre as a broadcast spray or as a 1 1/4 percent solution with hand-held equipment to provide partial control of alligatorweed. Apply when most of the target plants are in bloom. Repeat applications will be required to maintain such control.
- Bermudagrass:** Apply 7 1/2 pints of this product per acre as a broadcast spray or as a 1 1/2 percent solution with hand-held equipment. Apply when target plants are actively growing and when seedheads appear.
- Bindweed, field / Silverleaf Nightshade / Texas Blueweed:** Apply 6 to 7 1/2 pints of this product per acre as a broadcast spray west of the Mississippi River and 4 1/2 to 6 pints of this product per acre east of the Mississippi River. With hand-held equipment, use a 1 1/2 percent solution. Apply when target plants are actively growing and are at or beyond full bloom. For silverleaf nightshade, best results can be obtained when application is made after berries are formed. Do not treat when weeds are under drought stress. New leaf development indicates active growth. For best results apply in late summer or fall.
- Brackenfern:** Apply 4 1/2 to 6 pints of this product per acre as a broadcast spray or as a 3/4 to 1 percent solution with hand-held equipment. Apply to fully expanded fronds which are at least 18 inches long.
- Cattail:** Apply 4 1/2 to 6 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Apply when target plants are actively growing and are at or beyond the early-to-full bloom stage of growth. Best results are achieved when application is made during the summer or fall months.
- Cogongrass:** Apply 4 1/2 to 7 1/2 pints of this product per acre as a broadcast spray. Apply when cogongrass is at least 18 inches tall and actively growing in late summer or fall. Allow 7 or more days after application before tillage or mowing. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to maintain control.
- Cordgrass:** Apply 4 1/2 to 7 1/2 pints of this product per acre as a broadcast spray or as a 1 to 2 percent solution with hand-held equipment. Schedule applications in order to allow 6 hours before treated plants are covered by tidewater. The presence of debris and silt on the cordgrass plants will reduce performance. It may be necessary to wash targeted plants prior to application to improve uptake of this product into the plant.
- Cutgrass, giant:** Apply 6 pints of this product per acre as a broadcast spray or as a 1 percent solution with hand-held equipment to provide partial control of giant cutgrass. Repeat applications will be required to maintain such control, especially where vegetation is partially submerged in water. Allow for substantial regrowth to the 7 to 10-leaf stage prior to retreatment.
- Dogbane, hemp / Knapweed / Horseradish:** Apply 6 pints of this product per acre as a broadcast spray or as a 1 1/2 percent solution with hand-held equipment. Apply when target plants are actively growing and most have reached the late bud-to-flower stage of growth. For best results, apply in late summer or fall.
- Fescue, tall:** Apply 4 1/2 pints of this product per acre as a broadcast spray or as a 1 percent solution with hand-held equipment. Apply when target plants are actively growing and most have reached the boot-to-head stage of growth. When applied prior to the boot stage, less desirable control may be obtained.
- Guineagrass:** Apply 4 1/2 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Apply when target plants are actively growing and when most have reached at least the 7-leaf stage of growth.
- Johnsongrass / Bluegrass, Kentucky / Bromegrass, smooth / Canarygrass, reed / Orchardgrass / Ryegrass, perennial / Timothy / Wheatgrass, western:** Apply 3 to 4 1/2 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Apply when target plants are actively growing and most have reached the boot-to-head stage of growth. When applied prior to the boot stage, less desirable control may be obtained. In the fall, apply before plants have turned brown.
- Lantana:** Apply this product as a 3/4 to 1 percent solution with hand-held equipment. Apply to actively growing lantana at or beyond the bloom stage of growth. Use the higher application rate for plants that have reached the woody stage of growth.
- Loosestrife, purple:** Apply 4 pints of this product per acre as a broadcast spray or as a 1 to 1 1/2 percent solution using hand-held equipment. Treat when plants are actively growing at or beyond the bloom stage of growth. Best results are achieved when application is made during summer or fall months. Fall treatments must be applied before a killing frost.

15. **Lotus, American:** Apply 4 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Treat when plants are actively growing at or beyond the bloom stage of growth. Best results are achieved when application is made during summer or fall months. Fall treatments must be applied before a killing frost. Repeat treatment may be necessary to control regrowth from underground parts and seeds.
16. **Maidencane / Paragrass:** Apply 6 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Repeat treatments will be required, especially to vegetation partially submerged in water. Under these conditions, allow for regrowth to the 7 to 10-leaf stage prior to retreatment.
17. **Milkweed, common:** Apply 4 1/2 pints of this product per acre as a broadcast spray or as a 1 1/2 percent solution with hand-held equipment. Apply when target plants are actively growing and most have reached the late bud-to-flower stage of growth.
18. **Nutsedge: purple, yellow:** Apply 4 1/2 pints of this product per acre as a broadcast spray, or as a 3/4 percent solution with hand-held equipment to control existing nutsedge plants and immature nutlets attached to treated plants. Apply when target plants are in flower or when new nutlets can be found at rhizome tips. Nutlets which have not germinated will not be controlled and may germinate following treatment. Repeat treatments will be required for long-term control.
19. **Pampasgrass:** Apply a 1 1/2 percent solution of this product with hand-held equipment when plants are actively growing.
20. **Phragmites:** For partial control of phragmites in Florida and the counties of other states bordering the Gulf of Mexico, apply 7 1/2 pints per acre as a broadcast spray or apply a 1 1/2 percent solution with hand-held equipment. In other areas of the U.S., apply 4 to 6 pints per acre as a broadcast spray or apply a 3/4 percent solution with hand-held equipment for partial control. For best results, treat during late summer or fall months when plants are actively growing and in full bloom. Due to the dense nature of the vegetation, which may prevent good spray coverage and uneven stages of growth, repeat treatments may be necessary to maintain control. Visual control symptoms will be slow to develop.
21. **Quackgrass / Kikuyugrass / Muhly, wirestem:** Apply 3 to 4 1/2 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment when most quackgrass or wirestem muhly is at least 8 inches in height (3 to 4-leaf stage of growth) and actively growing. Allow 3 or more days after application before tillage.
22. **Reed, giant / ice plant:** For control of giant reed and ice plant, apply a 1 1/2 percent solution of this product with hand-held equipment when plants are actively growing. For giant reed, best results are obtained when applications are made in late summer to fall.
23. **Spatterdock:** Apply 6 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Apply when most plants are in full bloom. For best results, apply during the summer or fall months.
24. **Sweet potato, wild:** Apply this product as a 1 1/2 percent solution using hand-held equipment. Apply to actively growing weeds that are at or beyond the bloom stage of growth. Repeat applications will be required. Allow the plant to reach the recommended stage of growth before retreatment.
25. **Thistle, Canada / artichoke:** Apply 3 to 4 1/2 pints of this product per acre as a broadcast spray or as a 1 1/2 percent solution with hand-held equipment for Canada thistle. To control artichoke thistle, apply a 2 percent solution as a spray-to-wet application. Apply when target plants are actively growing and are at or beyond the bud stage of growth.
26. **Torpedograss:** Apply 6 to 7 1/2 pints of this product per acre as a broadcast spray or as a 3/4 to 1 1/2 percent solution with hand-held equipment to provide partial control of torpedograss. Use the lower rates under terrestrial conditions, and the higher rates under partially submerged or a floating mat condition. Repeat treatments will be required to maintain such control.
27. **Tules, common:** Apply this product as a 1 1/2 percent solution with hand-held equipment. Apply to actively growing plants at or beyond the seedhead stage of growth. After application, visual symptoms will be slow to appear and may not occur for 3 or more weeks.
28. **Waterhyacinth:** Apply 5 to 6 pints of this product per acre as a broadcast spray or apply a 3/4 to 1 percent solution with hand-held equipment. Apply when target plants are actively growing and at or beyond the early bloom stage of growth. After application, visual symptoms may require 3 or more weeks to appear with complete necrosis and decomposition usually occurring within 60 to 90 days. Use the higher rates when more rapid visual effects are desired.
29. **Waterlettuce:** For control, apply a 3/4 to 1 percent solution of this product with hand-held equipment to actively growing plants. Use higher rates where infestations are heavy. Best results are obtained from mid-summer through winter applications. Spring applications may require retreatment.
30. **Waterprimrose:** Apply this product as a 3/4 percent solution using hand-held equipment. Apply to plants that are actively growing at or beyond the bloom stage of growth, but before fall color changes occur. Thorough coverage is necessary for best control.
31. **Other perennial weeds listed above:** Apply 4 1/2 to 7 1/2 pints of this product per acre as a broadcast spray or apply as a 3/4 to 1 1/2 percent solution with hand-held equipment.

Woody Brush and Trees

NOTE: If brush has been mowed or tilled or trees have been cut, do not treat until regrowth has reached the recommended stage of growth.

Application Rates and Timing

When applied as a 5 to 8 percent solution as a directed application as described in the "Hand-Held and High-Volume Equipment" section, this product will control or partially control all wood brush and tree species listed in this section of this label. Use the higher rate of application for dense stands and larger woody brush and trees.

Specific Brush or Tree Control Recommendations: Numbers in parentheses "(-)" following the common name of a listed brush or tree species refer to "Specific Brush or Tree Control Recommendations" which follow the species listing. See this section for specific application rates and timing for listed species.

For woody brush and trees, apply the recommended rate plus a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient. Use of this product without surfactant will result in reduced herbicide performance. Refer to the "Mixing and Application Instructions" section of this label and the surfactant manufacturer label for more information. Make applications when plants are actively growing and, unless otherwise directed, after full-leaf expansion. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In arid areas, best results are obtained when application is made in the spring or early summer when brush species are at high moisture content and are flowering. Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with fall treatments.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

See the "Directions for Use" and "Mixing and Application Instructions" sections in this label for labeled use and specific application instructions. **When applied as directed, this product plus a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient will control the following woody brush plants and trees:** (Numbers in parentheses "(-)" following common name of a listed brush or tree species refer to "Specific Brush or Tree Control Recommendations" for that species which follow the species listing.)

Common Name	Scientific Name
Alder (1)	<i>Alnus spp.</i>
Ash [†] (20)	<i>Fraxinus spp.</i>
Aspen, quaking (2)	<i>Populus tremuloides</i>
Bearclover, Bearmat (20)	<i>Chamaebatia foliolosa</i>
Birch (3)	<i>Betula spp.</i>
Blackberry (1)	<i>Rubus spp.</i>
Broom, French (4)	<i>Cytisus monspessulanus</i>
Broom, Scotch (4)	<i>Cytisus scoparius</i>
Buckwheat, California [†] (5)	<i>Eriogonum fasciculatum</i>
Cascara [†] (20)	<i>Rhamnus purshiana</i>
Catsclaw [†] (6)	<i>Acacia greggi</i>
Ceanothus (20)	<i>Ceanothus spp.</i>
Chamise (17)	<i>Adenostoma fasciculatum</i>
Cherry, bitter (7)	<i>Prunus emarginata</i>
Cherry, black (7)	<i>Prunus serotina</i>
Cherry, pin (7)	<i>Prunus pensylvanica</i>
Coyote brush (8)	<i>Baccharis consanguinea</i>
Creeper, Virginia [†] (20)	<i>Parthenocissus quinquefolia</i>
Dewberry (1)	<i>Rubus trivialis</i>
Dogwood (9)	<i>Cornus spp.</i>
Elderberry (3)	<i>Sambucus spp.</i>
Elm [†] (20)	<i>Ulmus spp.</i>
Eucalyptus, bluegum (10)	<i>Eucalyptus globulus</i>
Hasardia [†] (5)	<i>Haplopappus squamosus</i>
Hawthorn (2)	<i>Crataegus spp.</i>
Hazel (3)	<i>Corylus spp.</i>
Hickory (9)	<i>Carya spp.</i>
Holly, Florida (11) (Brazilian peppertree)	<i>Schinus terebinthifolius</i>
Honeysuckle (1)	<i>Lonicera spp.</i>
Hornbeam, American (20)	<i>Carpinus caroliniana</i>
Kudzu (12)	<i>Pueraria lobata</i>
Locust, black [†] (20)	<i>Robinia pseudoacacia</i>
Manzanita (20)	<i>Arctostaphylos spp.</i>
Maple, red [†] (13)	<i>Acer rubrum</i>
Maple, sugar (14)	<i>Acer saccharum</i>
Maple, vine [†] (20)	<i>Acer circinatum</i>
Monkey flower [†] (5)	<i>Mimulus guttatus</i>
Oak, black [†] (20)	<i>Quercus velutina</i>
Oak, northern pin (14)	<i>Quercus palustris</i>
Oak, post (1)	<i>Quercus stellata</i>

Common Name	Scientific Name
Oak, red (14)	<i>Quercus rubra</i>
Oak, southern red (7)	<i>Quercus falcata</i>
Oak, white [†] (20)	<i>Quercus alba</i>
Persimmon [†] (20)	<i>Diospyros spp.</i>
Poison-ivy (15)	<i>Rhus radicans</i>
Poison-oak (15)	<i>Rhus toxicodendron</i>
Poplar, yellow [†] (20)	<i>Liriodendron tulipifera</i>
Prunus (7)	<i>Prunus spp.</i>
Raspberry (1)	<i>Rubus spp.</i>
Redbud, eastern (20)	<i>Cercis canadensis</i>
Rose, multiflora (16)	<i>Rosa multiflora</i>
Russian-olive (20)	<i>Elaeagnus angustifolia</i>
Sage: black (17), white	<i>Salvia spp.</i>
Sagebrush, California (17)	<i>Artemisia californica</i>
Salmonberry (3)	<i>Rubus spectabilis</i>
Salt cedar [†] (9)	<i>Tamarix spp.</i>
Saltbush, sea myrtle (18)	<i>Baccharis halimifolia</i>
Sassafras (20)	<i>Sassafras aibidum</i>
Sourwood [†] (20)	<i>Oxydendrum arboreum</i>
Sumac, poison [†] (20)	<i>Rhus vernix</i>
Sumac, smooth [†] (20)	<i>Rhus glabra</i>
Sumac, winged [†] (20)	<i>Rhus copallina</i>
Sweetgum (7)	<i>Liquidambar styraciflua</i>
Swordfern [†] (20)	<i>Polystichum munitum</i>
Tallowtree, Chinese (17)	<i>Sapium sebiferum</i>
Thimbleberry (3)	<i>Rubus parviflorus</i>
Tobacco, tree [†] (5)	<i>Nicotiana glauca</i>
Trumpet creeper (2)	<i>Campsis radicans</i>
Waxmyrtle, southern [†] (11)	<i>Myrica cerifera</i>
Willow (19)	<i>Salix spp.</i>

[†] Partial control (See below for control or partial control instructions.)

Specific Brush or Tree Control Recommendations:

- Alder / Blackberry / Dewberry / Honeysuckle / Oak, Post / Raspberry:** For control, apply 4 1/2 to 6 pints per acre as a broadcast spray or as a 3/4 to 1 1/4 percent solution with hand-held equipment.
- Aspen, Quaking / Hawthorn / Trumpet creeper:** For control, apply 3 to 4 1/4 pints of this product per acre as a broadcast spray or as a 3/4 to 1 1/4 percent solution with hand-held equipment.
- Birch / Elderberry / Hazel / Salmonberry / Thimbleberry:** For control, apply 3 pints per acre of this product as a broadcast spray or as a 3/4 percent solution with hand-held equipment.
- Broom, French / Broom, Scotch:** For control, apply a 1 1/4 to 1 1/2 percent solution with hand-held equipment.
- Buckwheat, California / Hasardia / Monkey flower / Tobacco, tree:** For partial control of these species, apply a 3/4 to 1 1/2 percent solution of this product as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.
- Catsclaw:** For partial control, apply a 1 1/4 to 1 1/2 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.
- Cherry, bitter / Cherry, black / Cherry, pin / Oak, southern red / Sweetgum / Prunus:** For control, apply 3 to 7 1/2 pints of this product per acre as a broadcast spray or as a 1 to 1 1/2 percent solution with hand-held equipment.
- Coyote brush:** For control, apply a 1 1/4 to 1 1/2 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.

9. **Dogwood / Hickory / Salt cedar:** For partial control, apply a 1 to 2 percent solution of this product with hand-held equipment or 6 to 7 1/2 pints per acre as a broadcast spray.
10. **Eucalyptus, bluegum:** For control of eucalyptus resprouts, apply a 1 1/2 percent solution of this product with hand-held equipment when resprouts are 6 to 12-feet tall. Ensure complete coverage. Apply when plants are actively growing. Avoid application to drought-stressed plants.
11. **Holly, Florida / Waxmyrtle, southern:** For partial control, apply this product as a 1 1/2 percent solution with hand-held equipment.
12. **Kudzu:** For control, apply 6 pints of this product per acre as a broadcast spray or as a 1 1/2 percent solution with hand-held equipment. Repeat applications will be required to maintain control.
13. **Maple, red:** For control, apply as a 3/4 to 1 1/4 percent solution with hand-held equipment when leaves are fully developed. For partial control, apply 2 to 7 1/2 pints of this product per acre as a broadcast spray.
14. **Maple, sugar / Oak: northern pin / Oak, red:** For control, apply as a 3/4 to 1 1/4 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.
15. **Poison-ivy / Poison-oak:** For control, apply 6 to 7 1/2 pints of this product per acre as a broadcast spray or as a 1 1/2 percent solution with hand-held equipment. Repeat applications may be required to maintain control. Fall treatments must be applied before leaves lose green color.
16. **Rose, multiflora:** For control, apply 3 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Treatments should be made prior to leaf deterioration by leaf-feeding insects.
17. **Sage, black / Sagebrush, California / Chamise / Tallowtree, Chinese:** For control of these species, apply a 3/4 percent solution of this product as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.
18. **Saltbush, sea myrtle:** For control, apply this product as a 1 percent solution with hand-held equipment.
19. **Willow:** For control, apply 4 1/2 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment.
20. **Other woody brush and trees listed above:** For partial control, apply 3 to 7 1/2 pints of this product per acre as a broadcast spray or as a 3/4 to 1 1/2 percent solution with hand-held equipment.

Aquatic and other Noncrop Sites

Apply this product as directed and under conditions described to control or partially control weeds and woody plants listed in the "Weeds Controlled" section in industrial, recreational and public areas or other similar aquatic or terrestrial sites on this label.

Noncrop Sites

This product may be used to control the listed weeds in and around aquatic sites and on noncrop sites such as :

- Airports
- Golf Courses
- Habitat Restoration & Management Areas
- Highways & Roadsides
- Industrial Plant Sites
- Lumberyards
- Parking Areas
- Parks
- Petroleum Tank Farms

- Pipeline, Power, Telephone & Utility Rights-of-Way
- Pumping Installations
- Railroads
- Schools
- Storage Areas
- Similar Sites

Aquatic Sites

This product may be applied to emerged weeds in all bodies of fresh and brackish water which may be flowing, nonflowing or transient. This includes lakes, rivers, streams, ponds, estuaries, rice levees, seeps, irrigation and drainage ditches, canals, reservoirs, wastewater treatment facilities, wildlife habitat restoration and management areas and similar sites.

If aquatic sites are present in the noncrop area and are part of the intended treatment, read and observe the following directions:

- **This product does not control plants which are completely submerged or have a majority of their foliage under water.**
- There is no restriction on the use of treated water for irrigation, recreation or domestic purposes.
- Consult local state fish and game agency and water control authorities before applying this product to public water. Permits may be required to treat such water.
- **NOTE:** Do not apply this product directly to water within 1/2 mile upstream of an active potable water intake in flowing water (i.e., river, stream, etc.) or within 1/2 mile of an active potable water intake in a standing body of water such as lake, pond or reservoir. To make aquatic applications around and within 1/2 mile of active potable water intakes, the water intake must be turned off for a minimum period of 48 hours after the application. The water intake may be turned on prior to 48 hours if the glyphosate level in the intake water is below 0.7 parts per million as determined by laboratory analysis. These aquatic applications may be made **only** in those cases where there are alternative water sources or holding ponds which would permit the turning off of an active potable water intake for a minimum period of 48 hours after the applications. This restriction does not apply to intermittent inadvertent overspray of water in terrestrial use sites.
- For treatments after drawdown of water or in dry ditches, allow 7 or more days after treatment before reintroduction of water to achieve maximum weed control. Apply this product within 1 day after drawdown to ensure application to actively growing weeds.
- Floating mats of vegetation may require retreatment. Avoid wash-off of sprayed foliage by spray boat or recreational boat backwash or by rainfall within 6 hours of application. Do not re-treat within 24 hours following the initial treatment.
- Applications made to moving bodies of water must be made while traveling upstream to prevent concentration of this herbicide in water. When making any bankside applications, do not overlap more than 1 foot into open water. Do not spray in bodies of water where weeds do not exist. The maximum application rate of 7 1/2 pints per acre must not be exceeded in any single broadcast application that is being made over water.

- When emerged infestations require treatment of the total surface area of impounded water, treating the area in strips may avoid oxygen depletion due to decaying vegetation. Oxygen depletion may result in fish kill.

Forestry Sites and Utility Rights-of-Way

In forest and utility sites, this product is recommended for the control or partial control of woody brush, trees, and annual and perennial herbaceous weeds. This product is also recommended for use in preparing or establishing wildlife openings within these sites, in pine straw plantations for maintaining logging roads, and for side trimming along utility rights-of-way.

In forestry sites, this product is recommended for use in site preparation prior to planting any tree species, including Christmas trees and silvicultural nursery sites.

In utility sites, this product is recommended for use along electrical power, pipeline, and telephone rights-of-way, and in other utility sites associated with these rights-of-way, such as substations.

Application Rates [†]:

Method of Application	Application Rate	Spray Volume (gal/acre)
Broadcast		
Aerial	1.5 to 7.5 qt/acre	5 to 30
Ground	1.5 to 7.5 qt/acre	10 to 60
Spray-to-Wet		
Handgun, Backpack Mistblower	0.75 to 2% by volume	spray-to-wet
Low Volume Directed Spray ^{††}		
Handgun, Backpack Mistblower	5% to 10% by volume	partial coverage

[†]Where repeat applications are necessary, do not exceed 8.0 quarts per acre per year.

^{††}For low volume directed spray applications, coverage should be uniform with at least 50 percent of the foliage contacted. For best results, coverage of the top one-half of the plant is important.

In forestry site preparation and utility rights-of-way applications, this product requires use with a surfactant such as a non-ionic surfactant containing greater than 80 percent active ingredient. Use of this product without surfactant will result in reduced herbicide performance. Refer to the "Mixing and Application Instructions" section of this label and the surfactant manufacturer label for more information.

Use higher rates of this product within the recommended rate ranges for control or partial control of woody brush, trees and hard-to-control perennial herbaceous weeds. For best results, apply to actively growing woody brush and trees after full leaf expansion and before fall color and leaf drop. Use increased rates within the recommended rate range to control of perennial herbaceous weeds from emergence up to the appearance of seedheads, flowers or berries appear. Use lower rates within the recommended rate range to control annual herbaceous weeds and actively growing perennial herbaceous weeds after seedheads, flowers or berries appear. Apply to foliage of actively growing annual herbaceous weeds anytime after emergence.

Tank Mixtures

This product may be used in tank mix combination with other herbicide products to broaden the spectrum of vegetation controlled. When tank mixing, read and observe applicable use directions, precautions and limitations on the respective product labels. Use according to the most restrictive precautionary statements for each product on the mixture. Any recommended rate of this product may be used in a tank mix.

Note: For forestry site preparation, make sure the tank mix product is approved for use prior to planting the desired species. Observe planting interval restrictions. For side trimming treatments in utility rights-of-way, tank mixtures with Arsenal 2WSL herbicide are not recommended. For side trimming treatments, it is recommended that this product be used alone as recommended, or as a tank mix with Garlon.

Product	Broadcast Rate	Use Sites
Arsenal Applicators Concentrate	2 to 16 fl oz/acre	Forestry site preparation
Oust	1 to 4 oz/acre	Forestry site preparation, utility sites
Garlon 3A [†]	1 to 4 qt/acre	Forestry site preparation, utility sites
Garlon 4	1 to 4 qt/acre	Forestry site preparation, utility sites
Arsenal 2WSL	2 to 32 fl oz/acre	Utility sites
	Spray-to-Wet Rates	
Arsenal Applicators Concentrate	1/32% to 1/2% by volume	Forestry site preparation
Arsenal 2WSL	1/32% to 1/2% by volume	Utility sites
	Low Volume Directed Spray Rates	
Arsenal Applicators Concentrate	1/8% to 1/2% by volume	Forestry site preparation
Arsenal 2WSL	1/8% to 1/2% by volume	Utility sites

[†]Ensure that Garlon 3A is thoroughly mixed with water before adding this product. Agitation is required while mixing this product with Garlon 3A to avoid compatibility problems.

For control of herbaceous weeds, use the lower recommended tank mixture rates. For control of dense stands or difficult-to-control woody brush and trees, use the higher recommended rates.

Forestry Conifer and Hardwood Release

Directed Sprays and Selective Equipment

This product may be applied as a directed spray or by using selective equipment in forestry conifer and hardwood sites, including Christmas tree plantations and silvicultural nurseries. This product requires use with a surfactant. Use only surfactants that are approved for conifer release and specified on the surfactant label as safe for use in conifer release (pine release). Use of this product without surfactant will result in reduced herbicide performance. Refer to the "Mixing and Application Instructions" section of this label and the surfactant manufacturer label for more information.

Tank Mixing: In hardwood plantations, tank mixtures with Oust may be used. In pine plantations, tank mixtures with Garlon 4 or Arsenal AC may be used. Comply with all site restrictions, forestry species limitations, and precautions on the tank mix product labels.

Avoid contact of spray drift, mist or drips with foliage, green bark or non-woody surface roots of desirable plant species. See "Application Equipment and Techniques" section of this label for specific recommendations and precautions.

Spray-to-Wet Applications: Use a 2 percent spray solution to control undesirable woody brush and trees. To control herbaceous weeds, use a 1 to 2 percent spray solution.

Low Volume Directed Spray Applications: Use a 5 to 10 percent spray solution. Coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one-half of the unwanted vegetation is important.

Broadcast Applications: For equipment calibrated for broadcast applications, use 1 1/2 to 7 1/2 quarts of this product per acre. Apply in 10 to 60 gallons of clean water per acre. Shielded application equipment may be used to avoid contact of the spray solution with desirable plants. Shields should be adjusted to prevent spray contact with the foliage of green bark of desirable vegetation.

Wiper Application Equipment: See the "Selective Equipment" section of this label for equipment and application rate recommendations.

Broadcast Application

Note: Except where specifically recommended below, make broadcast applications of this product only where conifers have been established for more than one year.

Broadcast application must be made after formation of final conifer resting buds in the fall or prior to initial bud swelling in the spring.

Injury may occur to conifers treated for release, especially where spray patterns overlap or the higher rates are applied. Damage can be accentuated if applications are made when conifers are actively growing, or are under stress from drought, flood water, improper planting, insects, animal damage or diseases.

This product requires use with a surfactant. Use a surfactant that is labeled/recommended for use in over-the-top release applications. Use of this product without a surfactant will result in reduced herbicide performance. Refer to the "Mixing and Application Instructions" section of this label and the surfactant manufacturer label for more information.

For release of the following conifer species outside the Southeastern United States:

Douglas fir (*Pseudotsuga menziesii*)

Fir (*Abies* species)

Hemlock^{††} (*Tsuga* species)

Pines[†] (*Pinus* species)

Redwood, California^{††} (*Sequoia* species)

[†] Includes all species except loblolly pine, longleaf pine, shortleaf pine or slash pine.

^{††} Use of a surfactant is not recommended for release of hemlock species or California redwood. In mixed conifer stands, injury to these species may result if a surfactant is used.

Application Rate for Conifer Release: Apply 3/4 to 1 1/2 quarts per acre as a broadcast spray. In Maine and New Hampshire, up to 2 1/4 quarts per acre of this product may be used for the control and suppression of difficult-to-control hardwood species.

To release Douglas fir, and pine and spruce species at the end of the first growing season (except in California), apply 3/4 to 1 1/8 quarts per acre of this product. Make sure that all conifers are well hardened off.

Note: For release of Douglas fir with this product or recommended tank mixtures, a nonionic surfactant recommended for over-the-top foliar spray may be used. To avoid possible conifer injury, nonionic surfactants may be used at 2 fluid ounces per acre at elevations above 1500 feet, or 1 fluid ounce per acre in the coastal range or at elevations below 1500 feet. Use of surfactant rates exceeding those listed above may result in unacceptable conifer injury and are not recommended. Make sure that the nonionic surfactant has been adequately tested for safety to Douglas fir before use.

Tank Mixtures with Oust: To release jack pine, white pine and white spruce, apply 3/4 to 1 1/2 quarts of this product with 1 to 3 ounces (1 to 1 1/2 ounces for white pine) of Oust per acre. Make applications to actively growing weeds as a broadcast spray over the top of established conifers. Applications at these rates should be made after formation of conifer resting buds in the late summer or fall.

Tank Mixtures with Arsenal Applicators Concentrate: This product may be tank mixed with Arsenal Applicators Concentrate for release of Douglas fir. Tank mix 3/4 to 1 1/8 quarts of this product with 2 to 6 fluid ounces of Arsenal Applicators Concentrate per acre. For release of balsam fir and red spruce, apply a mixture of 1 1/2 quarts of this product with 1 to 2 1/2 fluid ounces of Arsenal Applicators Concentrate per acre.

In Maine and New Hampshire for the release of red pine, balsam fir, red spruce, white spruce, Norway spruce, and black spruce with dense tough-to-control brush and where maples make up a large component of the undesirable trees, up to 2 1/4 quarts per acre of this product may be tank mixed with 1 to 2 1/2 fluid ounces per acre of Arsenal Applicators Concentrate herbicide and applied as a broadcast spray.

Tank mixtures with Arsenal Applicators Concentrate and Oust or Oust XP Herbicides: In Maine and New Hampshire for release of red pine, balsam fir, red spruce, white spruce, Norway spruce and black spruce with heavy grass and herbaceous weed densities, tough-to-control brush and where maples make up a large component of the undesirable trees up to 2 1/4 quarts per acre of this product may be tank mixed with 1 to 2.5 fluid ounces per acre of Arsenal Applicators Concentrate and 1 to 3 oz of Oust or Oust XP herbicides and applied as a broadcast spray.

For release of the following conifer species in the Southeastern United States:

Loblolly pine (*Pinus taeda*)

Eastern white pine (*Pinus strobus*)

Shortleaf pine (*Pinus echinata*)

Slash pine (*Pinus elliotii*)

Virginia pine (*Pinus virginiana*)

Longleaf pine (*Pinus palustris*)

Apply 1 1/8 to 1 7/8 quarts of this product per acre as a broadcast spray during late summer or early fall after the conifers have hardened off. For applications at the end of the first growing season, use 3/4 quart of this product alone or in a recommended tank mixture.

Tank Mixtures with Arsenal Applicators Concentrate: For conifer release, apply 3/4 to 1 1/2 quarts of this product with 2 to 16 fluid ounces of Arsenal Applicators Concentrate per acre as a broadcast spray. Use only on conifer species that are labeled for over-the-top spray for both products. Use the higher recommended rates for dense tough-to-control wood brush and trees.

Read and observe label claims, cautionary statements and all information on the labels of each product used in these tank mixtures. Use according to the most restrictive precautionary statements for each product in the mixture.

Herbaceous Release

When applied as directed, this product plus listed residual herbicides provides postemergence control of the annual weeds and control or suppression of the perennial weeds listed in this label, and residual control of the weeds listed in the residual herbicide label. Make applications to actively growing weeds as a broadcast spray over the top of labeled conifers.

Tank Mixtures with Oust: To release loblolly pines, tank mix 12 to 18 fluid ounces of this product with 2 to 4 ounces of Oust per acre.

To release slash pines, tank mix 9 to 12 fluid ounces of this product with 2 to 4 ounces of Oust per acre.

In Maine and New Hampshire for release of red pine, balsam fir, red spruce, white spruce, Norway spruce, and black spruce with heavy grass and herbaceous weeds infesting the site, up to 2 1/4 quarts per acre of this product may be tank mixed with 1 to 3 oz of Oust herbicide or Oust XP herbicide to control grass, herbaceous weeds and woody brush, and applied as a broadcast spray.

For tank mixtures with Oust use a surfactant that is labeled/ recommended for use in over-the-top herbaceous release applications. Use of this product without a surfactant will result in reduced herbicide performance. Refer to the "Mixing and Application Instructions" section of this label and the surfactant manufacturer label for more information.

Weed control may be reduced if water volumes exceed 25 gallons per acre for these treatments.

Tank Mixture with Atrazine: To release Douglas fir, apply 3/4 quart of this product with 4 pounds a.i. of atrazine per acre. Apply only over Douglas fir that has been established for at least one full growing season. Apply in the early spring, usually mid-March through early April. Injury will occur if applications are made after bud swell in the spring. For this use, do not add surfactant to the tank mixture.

Always read and follow the manufacturer's label for all herbicides and surfactants used.

Wetland Sites

This product may be used in and around water (aquatic areas) and wetlands found in forestry and in power, telephone and pipeline rights-of-way sites, including where these sites are adjacent to or surrounding domestic water supply reservoirs, supply streams, lakes and ponds. Read and observe the following before making applications in and around water.

Consult local public water control authorities before applying this product in and around public water. Permits may be required to treat in such areas.

There is no restriction on the use of treated water for irrigation, recreation or domestic purposes.

Note: Do not apply this product directly to water within 1/2 mile up-stream of an active potable water intake in flowing water (i.e., river, stream, etc.) or within 1/2 mile of an active potable water intake in a standing body of water such as a lake, pond or reservoir. To make aquatic applications around and within 1/2 mile of active potable water intakes, the water intake must be turned off for a minimum period of 48 hours after application. These aquatic applications may be made ONLY in those cases where there are alternative water sources or holding ponds which would permit the turning off of an active potable water intake for a minimum period of 48 hours after the application. This restriction does not apply to intermittent inadvertent overspray of water in terrestrial use sites.

Do not spray open bodies of water where woody brush, trees and herbaceous weeds do not exist. The maximum application rate of 3 3/4 quarts per acre must not be exceeded in a single over-water broadcast application except as follows, where any recommended rate may be applied:

- Stream crossings in utility right-of-way.
- Where applications will result in less than 20 percent of the total water area being treated.

Wildlife Habitat Restoration and Management Areas

This product is recommended for the restoration and/or maintenance of native habitat and in wildlife management areas.

Habitat Restoration and Maintenance: When applied as directed, exotic and other undesirable vegetation may be controlled in habitat management areas. Applications may be made to allow recovery of native plant species, to open up water to attract waterfowl, and for similar broad-spectrum vegetation control requirements in habitat management areas. Spot treatments may be made to selectively remove unwanted plants for habitat enhancement. For spot treatments, care should be exercised to keep spray off of desirable plants.

Wildlife Food Plots: This product may be used as a site preparation treatment prior to planting wildlife food plots. Apply as directed to control vegetation in the plot area. Any wildlife food species may be planted after applying this product, or native species may be allowed to reinfest the area. If tillage is needed to prepare a seedbed, wait 7 days after applying this product before tilling to allow for maximum effectiveness.

Wiper Applications

For wick or wiper applications, mix 1 gallon of this product with 2 gallons of clean water to make a 33 percent solution. Addition of a nonionic surfactant at a rate of 10 percent by volume of total herbicide solution is recommended.

Wiper applications can be used to control or suppress annual and perennial weeds listed on this label. In heavy weed stands, a double application in opposite directions may improve results. See the "Weed Controlled" section in this label for recommended timing, growth stage and other instructions for achieving optimum results

Cut Stump Application

Woody vegetation may be controlled by treating freshly cut stumps of trees and resprouts with this product. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut vegetation close to the soil surface. **Apply a 50 to 100 percent solution of this product to freshly cut surface immediately after cutting.** Delay in applying this product may result in reduced performance. For best results, trees should be cut during periods of active growth and full leaf expansion.

When used according to directions for cut stump application, this product will **control, partially control or suppress** most woody brush and tree species, some of which are listed below:

Common Name	Scientific Name
Alder	<i>Alnus spp.</i>
Coyote brush †	<i>Baccharis consanguinea</i>
Dogwood †	<i>Cornus spp.</i>
Eucalyptus	<i>Eucalyptus spp.</i>
Hickory †	<i>Carya spp.</i>
Madrone	<i>Arbutus menziesii</i>
Maple †	<i>Acer spp.</i>
Oak	<i>Quercus spp.</i>
Poplar †	<i>Populus spp.</i>
Reed, giant	<i>Arundo donax</i>
Salt cedar	<i>Tamarix spp.</i>
Sweet gum †	<i>Liquidambar styraciflua</i>
Sycamore †	<i>Platanus occidentalis</i>
Tan oak	<i>Lithocarpus densiflorus</i>
Willow	<i>Salix spp.</i>

† This product is not approved for this use on these species in the state of California.

Injection and Frill Applications

Woody vegetation may be controlled by injection or frill application of this product. Apply this product using suitable equipment which must penetrate into living tissue. Apply the equivalent of 1 ml of this product per 2 to 3 inches of trunk diameter. This is best achieved by applying 25 to 100 percent concentration of this product either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying dilute material to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow runoff to occur from frill or cut areas in species that exude sap freely after frills or cutting. In species such as these, make frill or cut at an oblique angle so as to produce a cupping effect and use undiluted material. For best results, applications should be made during periods of active growth and full leaf expansion.

This treatment will control the following woody species:

Common Name	Scientific Name
Oak	<i>Quercus spp.</i>
Poplar	<i>Populus spp.</i>
Sweet gum	<i>Liquidambar styraciflua</i>
Sycamore	<i>Platanus occidentalis</i>

This treatment will suppress the following woody species:

Common Name	Scientific Name
Black gum †	<i>Nyssa sylvatica</i>
Dogwood	<i>Cornus spp.</i>
Hickory	<i>Carya spp.</i>
Maple, red	<i>Acer rubrum</i>

† This product is not approved for this use on this species in the state of California.

Release of Bermudagrass or Bahiagrass on Noncrop Sites

Release Of Dormant Bermudagrass And Bahiagrass

When applied as directed, this product will provide control or suppression of many winter annual weeds and tall fescue for effective release of dormant bermudagrass or bahiagrass. Make applications to dormant bermudagrass or bahiagrass.

For best results on winter annuals, treat when weeds are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall fescue, treat when fescue is in or beyond the 4 to 6-leaf stage.

Weeds Controlled

Rate recommendations for control or suppression of winter annuals and tall fescue are listed below.

Apply the recommended rates of this product in 10 to 25 gallons of water per acre plus a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient.. Use of this product without surfactant will result in reduced herbicide performance. Refer to the "Mixing and Application Instructions" section of this label and the surfactant manufacturer label for more information.

Weeds Controlled or Suppressed †

Note: C = Controlled; S = Suppressed

Weed Species	Rate (Fluid Ounces Per Acre)					
	6	9	12	18	24	48
Barley, little <i>Hordeum pusillum</i>	S	C	C	C	C	C
Bedstraw, catchweed <i>Galium aparine</i>	S	C	C	C	C	C
Bluegrass, annual <i>Poa annua</i>	S	C	C	C	C	C
Chervil <i>Chaerophyllum tainturieri</i>	S	C	C	C	C	C
Chickweed, common <i>Stellaria media</i>	S	C	C	C	C	
Clover, crimson <i>Trifolium incarnatum</i>	•	S	S	C	C	C
Clover, large hop <i>Trifolium campestre</i>	•	S	S	C	C	C
Speedwell, corn <i>Veronica arvensis</i>	S	C	C	C	C	C
Fescue, tall <i>Festuca arundinacea</i>	•	•	•	•	S	S
Geranium, Carolina <i>Geranium carolinianum</i>	•	•	S	S	C	C
Henbit <i>Lamium amplexicaule</i>	•	S	C	C	C	C
Ryegrass, Italian <i>Lolium multiflorum</i>	•	•	S	C	C	C
Vetch, common <i>Vicia sativa</i>	•	•	S	C	C	C

†These rates apply only to sites where an established competitive turf is present.

Release Of Actively Growing Bermudagrass

NOTE: Use only on sites where bahiagrass or bermudagrass are desired for ground cover and some temporary injury or yellowing of the grasses can be tolerated.

When applied as directed, this product will aid in the release of bermudagrass by providing control of annual species listed in the "Weeds Controlled" section in this label, and suppression or partial control of certain perennial weeds.

For control or suppression of those annual species listed in this label, use 3/4 to 2 1/4 pints of this product as a broadcast spray in 10 to 25 gallons of spray solution per acre, plus a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient.. Use of this product without surfactant will result in reduced herbicide performance. Refer to the "Mixing and Application Instructions" section of this label and the surfactant manufacturer label for more information. Use the lower rate when treating annual weeds below 6 inches in height (or length of runner in annual vines). Use the higher rate as size of plants increases or as they approach flower or seedhead formation.

Use the higher rate for partial control or longer-term suppression of the following perennial species. Use lower rates for shorter-term suppression of growth.

Bahiagrass	Johnsongrass [†]
Dallisgrass	Trumpetcreeper ^{††}
Fescue (tall)	Vaseygrass

[†] Johnsongrass is controlled at the higher rate.

^{††} Suppression at the higher rate only.

Use only on well-established bermudagrass. Bermudagrass injury may result from the treatment but regrowth will occur under moist conditions. Repeat applications in the same season are not recommended, since severe injury may result.

Bahiagrass Seedhead and Vegetative Suppression

When applied as directed in the "Noncrop Sites" section in this label, this product will provide significant inhibition of seedhead emergence and will suppress vegetative growth for a period of approximately 45 days with single applications and approximately 120 days with sequential applications.

Apply this product 1 to 2 weeks after full green-up of bahiagrass or after the bahiagrass has been mowed to a uniform height of 3 to 4 inches. Applications must be made prior to seedhead emergence. Apply 5 fluid ounces per acre of this product in 10 to 25 gallons of water per acre, plus a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient. Use of this product without surfactant will result in reduced herbicide performance. Refer to the "Mixing and Application Instructions" section of this label and the surfactant manufacturer label for more information.

Sequential applications of this product plus nonionic surfactant may be made at approximately 45-day intervals to extend the period of seedhead and vegetative growth suppression. For continued vegetative growth suppression, sequential applications must be made prior to seedhead emergence.

Apply no more than 2 sequential applications per year. As a first sequential application, apply 3 fluid ounces of this product per acre plus nonionic surfactant. A second sequential application of 2 to 3 fluid ounces per acre plus nonionic surfactant may be made approximately 45 days after the last application.

Annual Grass Growth Suppression

For growth suppression of some annual grasses, such as annual ryegrass, wild barley and wild oats growing in coarse turf on roadsides or other industrial areas, apply 3 to 4 ounces of this product in 10 to 40 gallons of water per acre plus a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient. Use of this product without surfactant will result in reduced herbicide performance. Refer to the "Mixing and Application Instructions" section of this label and the surfactant manufacturer label for more information. Applications should be made when annual grasses are actively growing and before the seedheads are in the boot stage of development. Treatments made after seedhead emergence may cause injury to the desired grasses.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. To the fullest extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. To the fullest extent permitted by law, in no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®Trademark of Dow AgroSciences LLC

**Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268**

Label Code: D02-148-005
Replaces Label: D02-148-004
LOES Number: 010-01471

EPA-accepted 07/13/06


Revisions:

1. Updated Storage and Disposal



DuPont™ Escort® XP

herbicide



“..... A Growing Partnership With Nature”



DuPont™ Escort® XP

herbicide

Dry Flowable

<i>Active Ingredient</i>	<i>By Weight</i>
Metsulfuron methyl Methyl 2-[[[(4-methoxy-6-methyl- 1,3,5-triazin-2-yl)amino]- carbonyl]amino]sulfonyl]benzoate	60%
<i>Inert Ingredients</i>	40%
TOTAL	100%

EPA Reg. No. 352-439

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-441-3637 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION! Causes eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust or spray mist.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

Long-sleeved shirt and long pants.

Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

This herbicide is injurious to plants at extremely low concentrations. Nontarget plants may be adversely effected from drift and run-off.

IMPORTANT

DO NOT USE ON FOOD OR FEED CROPS EXCEPT AS RECOMMENDED BY THIS LABEL OR SUPPLEMENTAL LABELING. Injury to or loss of desirable trees or other plants may result if the precautions listed below are not followed.

- Do not apply DuPont™ ESCORT® XP herbicide (except as recommended), or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend or in locations where the product may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, tennis courts, or similar areas.
- Prevent drift of spray to desirable plants.
- Do not contaminate any body of water, including irrigation water.
- Keep from contact with fertilizers, insecticides, fungicides and seeds.

Low rates of ESCORT® XP can kill or severely injure most crops. Following an ESCORT® XP application, the use of spray equipment to apply other pesticides to crops on which ESCORT® XP is not registered may result in their damage. The most effective way to reduce this crop damage potential is to use dedicated mixing and application equipment.

GENERAL INFORMATION

ESCORT® XP herbicide is a dispersible granule that is mixed in water and applied as a spray by ground or aerial application.

ESCORT® XP is recommended for the control of annual and perennial weeds and unwanted woody plants on private, public and military lands, on rights-of-way, industrial sites, non-crop areas, ditchbanks of dry drainage ditches, certain types of unimproved turf grass, and conifer and hardwood plantations, including grazed areas on these sites. Do not use on irrigation ditches.

ESCORT® XP controls weeds and woody plants primarily by postemergent activity. Although ESCORT® XP has preemergence activity, best results are generally obtained when ESCORT® XP is applied to foliage after emergence or dormancy break. Generally, for the control of annual weeds, ESCORT® XP provides the best results when applied to young, actively growing weeds. For the control of perennial weeds, applications made at the bud/bloom stage or while the target weeds are in the fall rosette stage may provide the best results. The use rate depends upon the weed species and size at the time of application.

The degree and duration of control may depend on the following:

- weed spectrum and infestation intensity
- weed size at application
- environmental conditions at and following treatment
- soil pH, soil moisture, and soil organic matter.

ESCORT® XP may be applied on conifer and hardwood plantations, and non-crop sites that contain areas of temporary surface water caused by the collection of water between planting beds, in equipment ruts, or in other

depressions created by management activities. It is permissible to treat intermittently flooded low lying sites, seasonally dry flood plains and transitional areas between upland and lowland sites when no water is present. It is also permissible to treat marshes, swamps and bogs after water has receded as well as seasonally dry flood deltas. DO NOT make applications to natural or man-made bodies of water such as lakes, reservoirs, ponds, streams and canals.

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

ESCORT® XP is absorbed primarily through the foliage of plants, and by the roots to a lesser degree. Plant cell division is generally inhibited in sensitive plants within a few hours following uptake. Two to 4 weeks after application, leaf growth slows followed by discoloration and tissue death. The final effects on annual weeds are evident about 4 to 6 weeks after application. The ultimate effect on perennial weeds and woody plants occurs in the growing season following application.

Warm, moist conditions following treatment promote the activity of ESCORT® XP, while cold, dry conditions may reduce or delay activity. Weeds and brush hardened off by cold weather or drought stress may not be controlled.

The use of a surfactant is recommended to enhance the control of susceptible plants, except where noted. Apply at a minimum rate (concentration) of 1/4% volume/volume (1 quart per 100 gallons of spray solution), or at the manufacturer's recommended rate. Use only EPA approved surfactants containing at least 80% active ingredient. Certain types of surfactants, such as those incorporating acetic acid (i.e. LI- 700), may not be compatible with ESCORT® XP and may result in decreased performance. Certain surfactants may not be suitable for use on desirable plants, such as turf and conifers, listed on this label. Consult the surfactant manufacturer's label for appropriate uses. Weed and brush control may be reduced if rainfall occurs soon after application.

RESISTANCE

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistant weed biotypes, it may be necessary to change cultural practices within and between crop seasons such as using a combination of tillage, retreatment, tank-mix partners and/or sequential herbicide applications that have a different site of action. Weed escapes that are allowed to go to seed will promote the spread of resistant biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.

INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DuPont™ ESCORT® XP should be used only in accordance with recommendations on this label or in separately published DuPont recommendations.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically recommended by DuPont. User assumes all risks associated with such non-recommended use.

Do not apply more than 4 ounces of ESCORT® XP per acre per year.

Do not use on food or feed crops except as recommended by this label or supplemental labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency in your State responsible for pesticide regulation.

TANK MIXES

ESCORT® XP may be tank mixed with other herbicides registered for the use sites described in this label. Use only those tank mix partners which are labeled for the appropriate use site. When tank mixing, use the most restrictive label limitations for each of the products being used in the tank mix.

AGRICULTURAL USES

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Shoes plus socks

CONIFER PLANTATIONS

Application Information

ESCORT® XP is recommended for the control of many species of weeds and deciduous trees on sites where conifers are growing or are to be planted. Apply by ground equipment or by air (helicopter only). Refer to the "Weeds Controlled" and "Brush Species Controlled" for a listing of susceptible species.

Application Timing

Apply ESCORT® XP after weeds have emerged or after undesirable hardwoods have broken winter dormancy and have reached the point of full leaf expansion.

Conifer Site Preparation

--Application Before Transplanting

After consulting the "Weeds Controlled" and "Brush Species Controlled" tables, apply the rates of ESCORT® XP recommended for the most difficult to control species on the site.

Southeast—Apply up to 4 ounces per acre for loblolly and slash pines. Transplant the following planting season.

Northeast and Lake States—Apply up to 2 ounces per acre for red pine. Transplant the following planting season. Apply up to 2 ounces per acre for black, white and Norway spruce. Transplant the following spring.

West—Apply up to 2 ounces per acre prior to planting Douglas Fir, Sitka Spruce, Western Red Cedar, Western Hemlock, Ponderosa Pine, and Grand Fir in the Coast Rangeland and western slope of the Cascades in Oregon and Washington. These conifer species listed can be planted anytime after application. Other conifer species can be planted providing the user has prior experience indicating acceptable tolerance to ESCORT® XP soil residues.

Without prior experience, it is recommended that other species be planted on a small scale to determine selectivity before large-scale plantings are made as unacceptable injury may occur. DuPont will not assume responsibility for injury to any conifer species not listed on this label.

Tank Mix Combinations—

For broader spectrum control, the following products are recommended in combination with DuPont™ ESCORT® XP.

Glyphosate (4 pound active per gallon)

Tank mix 1 to 2 ounces of ESCORT® XP with 2 to 10 quarts of glyphosate per acre. Refer to the product container for a list of species controlled.

Imazapyr (4 pound active per gallon)

Tank mix 1 to 2 ounces of ESCORT® XP with 10 to 24 fluid ounces of imazapyr per acre. Loblolly and slash pines may be transplanted the planting season following application. This combination controls ash, black gum, cherry, hawthorn, honeysuckle, hophornbeam, persimmon, oaks (red, white and water), sassafras, sweetgum, Vaccinium species, and suppresses blackberry, dogwood, elms, myrtle dahoon, hickories, and red maple.

Glyphosate (4 pound active per gallon) + Imazapyr (4 pound active per gallon)

Tank mix 1/2 to 1 ounce of ESCORT® XP with 16 to 64 fluid ounces of glyphosate and 10 to 12 fluid ounces of imazapyr per acre. Slash and loblolly pines may be transplanted the planting season following application. This combination controls cherry, dogwood, elms, oaks (red and water), persimmon, sassafras, sweetgum and suppresses hickory.

DuPont™ VELPAR® L or VELPAR® DF

Tank mix 1 to 2 ounces of ESCORT® XP per acre with VELPAR® L or VELPAR® DF at the rates recommended on the container for various soil textures. Loblolly and slash pines may be transplanted the planting season following application. Refer to the product container for a list of species controlled.

DuPont™ OUST® EXTRA

Tank mix 1/2 to 1 1/2 ounces of ESCORT® XP with 2 to 3 ounces of OUST® EXTRA per acre for herbaceous weed control. Refer to the product container and the "Weeds Controlled" section of this label for a listing of the weeds controlled. Loblolly and slash pines may be transplanted the planting season following application. Tank mix 2 ounces of ESCORT® XP with 3 ounces of OUST® EXTRA per acre for herbaceous weed control and early spring suppression of bull thistle and Canada thistle in the Coast Rangeland and western slope of the Cascade Mountains. Douglas fir may be transplanted at least 90 days following application.

Release--Hardwood Control and Suppression

ESCORT® XP is recommended for application over the top of established slash and loblolly pine to control the species listed in "Weeds Controlled" and "Brush Species Controlled" section of this label. Apply 1 to 4 ounces per acre to control the species indicated, including kudzu.

Tank Mix Combinations—

For broader spectrum control the following products are recommended in combination with ESCORT® XP.

Imazapyr (4 pound active per gallon)

Tank mix 1 to 2 ounces of ESCORT® XP with 8 to 16 fluid ounces of imazapyr per acre for application to loblolly pine. Refer to the imazapyr label regarding the use of surfactants and the appropriate application timing with respect to the age and development stage of the pines. This combination controls ash, black gum, cherry, hawthorn, honeysuckle, hophornbeam, oaks (red, white and water), sassafras, sweetgum, Vaccinium species, and suppresses blackberry, dogwood, elms, myrtle dahoon, hickories, persimmon, and red maple.

VELPAR® L or VELPAR® DF

Tank mix 1 to 2 ounces of ESCORT® XP with VELPAR® L or VELPAR® DF at the rates recommended on the container for various soil textures. This combination may be applied to loblolly and slash pines.

Release--Herbaceous Weed Control

ESCORT® XP may be applied to transplanted loblolly and slash pine for the control of herbaceous competition. Consult the "Weeds Controlled" for a listing of the susceptible species and recommended application rates. Best results are obtained when ESCORT® XP is applied just before weed emergence until shortly after weed emergence.

Tank Mix Combinations—

For broader spectrum control the following products are recommended in combination with ESCORT® XP.

Imazapyr (4 pound active per gallon)

Tank mix 1/2 to 1 ounce of ESCORT® XP with 4 fluid ounces of imazapyr per acre. The tank mix may be used on loblolly pine.

VELPAR® L or VELPAR® DF

Tank mix 1/2 to 1 ounce of ESCORT® XP with VELPAR® L or VELPAR® DF at the rates recommended on the container for various soil textures. This combination may be applied to loblolly and slash pines.

Release - Directed Spray in Conifers

Western US

To release conifers from competing brush species, such as, blackberry, salmonberry, snowberry, thimbleberry and wild roses, mix 2 to 4 ounces of ESCORT® XP per 100 gallons of spray solution. Direct spray onto the foliage of competing brush species using a knapsack or backpack sprayer. For best results, apply any time after the brush species have reached full leaf stage but before autumn coloration. At application, the majority of the brush species should be less than six feet in height to help ensure adequate spray coverage. Thorough coverage of the target foliage is necessary to optimize results. Care should be taken to direct the ESCORT® XP spray solution away from the conifer foliage.

NOTE:

DuPont™ ESCORT® XP may cause temporary yellowing and or growth suppression when the spray solution contacts conifer foliage. The use of a surfactant with ESCORT® XP may improve brush control results. When using a surfactant with ESCORT® XP, extra precaution must be taken to avoid contact with conifer foliage. Excessive drift onto conifers may result in severe injury.

IMPORTANT PRECAUTIONS

—CONIFER PLANTATIONS ONLY

- Applications of ESCORT® XP made to conifers that are suffering from loss of vigor caused by insects, diseases, drought, winter damage, animal damage, excessive soil moisture, planting shock, or other stresses may injure or kill the trees.
- Applications of ESCORT® XP made for herbaceous release should only be made after adequate rainfall has closed the planting slit and settled the soil around the roots following transplanting.
- Do not apply ESCORT® XP to conifers grown as ornamentals.
- ESCORT® XP applications may result in damage and mortality to other species of conifers when they are present on sites with those listed in the preceding recommendations for conifer plantations.

HARDWOOD PLANTATIONS

Application Information

ESCORT® XP is recommended at rates of up to 2 ounces per acre for the control of many weed species on sites where yellow poplar is growing or is to be planted, and on sites where red alder is to be planted. Apply by ground equipment or by air (helicopter only). Refer to the "Weeds Controlled" sections of this label for a listing of susceptible species.

Application Timing

ESCORT® XP may be applied as a site preparation treatment prior to planting red alder or yellow poplar. As a prior to planting site preparation treatment for red alder, ESCORT® XP may be tank mixed with other herbicides labeled for this use.

ESCORT® XP may also be applied over-the-top of planted yellow poplar seedlings after the soil has settled around the root system, but before the seedlings have broken dormancy (prior to bud break).

Release--Herbaceous Weed Control

ESCORT® XP may be applied to yellow poplar for the control of herbaceous competition. Consult the "Weeds Controlled" for a listing of the susceptible species and recommended application rates. Best results are obtained when ESCORT® XP is applied just before weed emergence until shortly after weed emergence.

Tank Mix Combinations—

Tank mix 1/2 ounce of ESCORT® XP with 4 to 6 pints of DuPont™ VELPAR® L as recommended on the package label for "RELEASE--HERBACEOUS WEED CONTROL" in pine plantations in the eastern U.S. Follow the VELPAR® L label recommendations regarding altering the application rate by soil texture.

IMPORTANT PRECAUTIONS

—HARDWOOD PLANTATIONS ONLY

- Application of VELPAR® L and ESCORT® XP made to yellow poplar that are suffering from loss of vigor caused by insects, disease, drought, winter damage, animal damage, excessive soil moisture, planting shock or other stresses may injure or kill the seedlings.
- Applications of ESCORT® XP made for release should only be made after adequate rainfall has closed the planting slit and settled the soil around the roots following transplanting.
- The use of surfactant is not recommended for applications made over the tops of trees.
- Careful consideration must be given by an experienced and knowledgeable forester to match the requirements of yellow poplar and/or red alder to the conditions of the site. Treatment of yellow poplar and/or red alder planted on a site inadequate to meet its requirements may injure or kill the seedlings.

NON-AGRICULTURAL USES

NON-AGRICULTURAL USE REQUIREMENTS
 The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.
 Do not enter or allow others to enter the treated area until sprays have dried.
Non-crop industrial weed control and selective weed control in turf (industrial, unimproved only) are not within the scope of the Worker Protection Standard.

NON-CROP SITES

Application Information

ESCORT® XP is recommended for general weed control on private, public and military lands as follows: Uncultivated nonagricultural areas (such as airports, highway, railroad and utility rights-of-way, sewage disposal areas, etc.); uncultivated agricultural areas - non-crop producing (such as farmyards, fuel storage areas, fence rows, soil bank land, barrier strips, etc.); industrial sites - outdoor (such as lumberyards, pipeline and tank farms, etc.) including grazed areas on these sites. It is also recommended for the control of certain noxious and troublesome weeds.

Consult the "Weeds Controlled" and "Brush Species Controlled" tables to determine the appropriate application rate.

ESCORT® XP may be applied in tank mixture with other herbicides labeled for use on non-crop sites. Fully read the labels and follow all directions and restrictions on each label.

Applications may be made by ground or air. Use a sufficient volume of water to ensure thorough coverage of the target vegetation with the application equipment being used.

NATIVE GRASSES

DuPont™ ESCORT® XP is recommended for weed control and suppression in the establishment and maintenance of native grasses. It may be used where blue grama, bluestems (big, little, plains, sand, ww spar) bromegrasses (meadow), buffalograss, green sprangletop, indiangrass, kleingrass, lovegrasses (atherstone, sand, weeping, wilman), orchardgrass, sideoats grama, switchgrass (blackwell), wheatgrass (bluebunch, intermediate, pubescent Siberian, slender, streamband, tall, thickspike, western), and Russian wildrye are established. It may also be applied over these species in the seedling stage, except for orchardgrass and Russian wildrye.

Application Information

Apply ESCORT® XP at the rate of 1/10 ounce per acre for the control and suppression* of bur buttercup (testiculate), common purslane, common sunflower*, cutleaf eveningprimrose*, flixweed*, lambsquarters* (common and slimleaf), marestail*, pigweed (redroot and tumble), snow speedwell, tansymustard* and tumble mustard (Jim Hill mustard).

* Suppression is a visual reduction in weed competition (reduced population or vigor) as compared to untreated areas. Degree of suppression will vary with the size of weed and environmental conditions following treatment.

Application Timing

For established grasses, apply when weeds are in the seedling stage.

For grasses in the seedling stage, apply preplant or preemergence where the soil (seed bed) has been cultivated.

IMPORTANT PRECAUTIONS

—NATIVE GRASSES

- Grass species or varieties may differ in their response to various herbicides. If no information is available, limit the initial use of ESCORT® XP to a small area. Components in a grass seed mixture will vary in tolerance to ESCORT® XP, so the final stand may not reflect the seed ratio.
- Under certain conditions such as heavy rainfall, high pH, prolonged cold weather, or wide fluctuations in day/night temperatures prior to or soon after ESCORT® XP application, temporary discoloration and/or grass injury may occur. ESCORT® XP should not be applied to grass that is stressed by severe weather conditions, drought, low fertility, water-saturated soils, disease, or insect damage as grass injury may result. Severe winter stress, drought, disease, or insect damage before or following application also may result in grass injury.

GRASS REPLANT INTERVALS

Following an application of ESCORT® XP to non-crop areas, the treated sites may be replanted with various species of grasses at the intervals recommended below.

For soils with a pH of 7.5 or less, observe the following replant intervals:

Species	Rate (ounces per acre)	Replant Interval (months)
Brome, Meadow	1/2—1	2
	1—2	3
Brome, Smooth	1/2—1	2
	1—2	4
Fescue, Alta	1/2—1	2
	1—2	4
Fescue, Red	1/2—1	2
	1—2	4
Fescue, Sheep	1/2—1	1
	1—2	4
Foxtail, Meadow	1/2—1	2
	1—2	4
Green Needlegrass	1/2—2	1
Orchardgrass	1/2—1	2
	1—2	4
Russian wildrye	1/2—1	1
	1	2
	2	3
Switchgrass	1/2—1	1
	1—2	3
Timothy	1/2—1	2
	1—2	4
Wheatgrass, Western	1/2—1	2
	1—2	3

For soils with a pH of 7.5 or greater observe the following replant intervals:

Species	Rate (ounces per acre)	Replant Interval (months)
Alkali Sacaton	1/2—1	1
	1—2	3
Bluestem, Big	1/2—2	3
Brome, Mountain	1/2—1	1
	1—2	2
Grama, Blue	1/2—2	1
Grama, Sideoats	1/2	2
	>1/2	>3
Switchgrass	1/2	2
	>1/2	>3
Wheatgrass, Thickspike	1/2—2	1
Wheatgrass, Western	1—2	2
	1/2—1	3

The recommended intervals are for applications made in the Spring to early Summer. Because ESCORT® XP degradation is slowed by cold or frozen soils, applications made in the late Summer or Fall should consider the intervals as beginning in the Spring following treatment. Testing has indicated that there is considerable variation in response among the species of grasses when seeded into areas treated with ESCORT® XP. If species other than those listed above are to be planted into areas treated with ESCORT® XP, a field bioassay should be performed, or previous experience may be used, to determine the feasibility of replanting treated sites.

**ADDITIONAL GRASS INFORMATION
APPLICATION INFORMATION FOR GRASS
ESTABLISHMENT**

DuPont™ ESCORT® XP is recommended for the control or suppression of broadleaf weeds to aid in the establishment of the following perennial native or improved grasses:

Blue Grama	Sideoats grama
Bluestems --	Switchgrass –
Big	Blackwell
Little	Wheatgrasses –
Plains	bluebunch
Sand	crested
W W spar	intermediate
Buffalograss	pubescent
Green sprangletop	Siberian
Kleingrass	slender
Lovegrasses --	steambank
Atherstone	tall
Sand	thickspike
Weeping	western
Wilman	Wildrye grass –
Orchardgrass	Russian

Maximize potential for grass establishment by consulting with the Natural Resource and Conservation Service of other government agencies or local experts concerning planting techniques and other cultural practices. Performance from ESCORT® XP may not always be satisfactory due to the inability of newly planted grass stands to sufficiently compete with weeds, and the severity of weed pressure in new grass stands.

An additional herbicide application or mowing may be needed.

Use Rates and Application Timing for Grass Establishment Preplant (prior to planting) or Preemergence (after planting but before grass emergence)

Do not use more than 1/10 ounce per acre of ESCORT® XP for grass establishment.

Apply ESCORT® XP at 1/10 ounce per acre on all labeled grasses except orchardgrass and Russian wildrye grass. Do not apply ESCORT® XP preplant or preemergence to orchardgrass and Russian wildrye grass as severe crop injury may result.

Early postemergence to new plantings

Apply ESCORT® XP at 1/10 ounce per acre, plus a non-ionic surfactant at the rate of 2 to 4 pints per 100 gallons of spray solution on all labeled grasses anytime after grass emergence.

Do not use a spray adjuvant other than non-ionic surfactant. Because grass species differ in time of emergence, apply only after the majority of grasses are in the 3 to 4 leaf stage.

Postemergence to stands with 1 – 5 leaf grasses planted the previous season

Apply ESCORT® XP at 1/10 ounce per acre plus a non-ionic surfactant at the rate of 2 to 4 pints per 100 gallons of spray solution, on all labeled grasses when the majority of the grasses have one or more leaves.

Do not use a spray adjuvant other than non-ionic surfactant.

APPLICATION INFORMATION FOR ESTABLISHED GRASSES

Use Rates for Established Grasses

Apply up to 1 ounce ESCORT® XP per acre as a broadcast application to established grasses. For spot applications, use 1 ounce per 100 gallons of water. Do not apply more than 1 2/3 ounces of ESCORT® XP per acre per year.

Refer to the Weeds Controlled section of this label for a listing of the weeds controlled by ESCORT® XP and the appropriate use rate to obtain control.

Application Timing – Established Grasses

ESCORT® XP may be applied to established native grasses such as bluestems and grama, and on other established grasses such as bermudagrass, bluegrass, orchardgrass, bromegrass, fescue and timothy that were planted the previous growing season (or earlier) and are fully tillered, unless otherwise directed on this label. Specific application timing information on several of these grass species follows:

Grass	Minimum time from Grass establishment ESCORT® XP application
Bermudagrass	2 months
Bluegrass, bromegrass, Orchardgrass	6 months
Timothy	12 months
Fescue	24 months

Fescue Precautions:

Note that ESCORT® XP may temporarily stunt tall fescue, cause it to turn yellow, or cause seedhead suppression. To minimize these symptoms, take the following precautions:

- Do not use more than 4/10 ounce per acre of ESCORT® XP
 - Tank mix ESCORT® XP with 2,4-D
 - Use the lowest recommended rate for target weeds
 - Use a non-ionic surfactant at 1/2 to 1 pint per 100 gallons of spray solution
 - Make application later in the spring after the new growth is 5 to 6 inches tall, or in the fall
 - Do not use surfactant when liquid nitrogen is used as a carrier
 - Do not use a spray adjuvant other than non-ionic surfactant
- The first cutting yields may be reduced due to seedhead suppression resulting from treatment with ESCORT® XP.

Timothy Precautions:

Timothy should be at least 6 inches tall at application and be actively growing. Applications of ESCORT® XP to timothy under any other conditions may cause crop yellowing and/or stunting. To minimize these symptoms, take the following precautions:

- Do not use more than 4/10 ounce per acre ESCORT® XP
 - Tank mix ESCORT® XP with 2, 4-D
 - Use the lowest recommended rate for target weeds
 - Use a non-ionic surfactant at 1/2 pint per 100 gallons of spray solution (1/16%)
 - Make applications in the late summer or fall
 - Do not use surfactant when liquid nitrogen is used as a carrier
 - Do not use spray adjuvant other than non-ionic surfactant
- Application of ESCORT® XP to Pensacola bahiagrass, ryegrass (Italian or perennial) and Garrison’s creeping foxtail may cause severe injury to and/or loss of forage.

Other Grasses:

Varieties and species of forage grasses differ in their tolerance to herbicides. When using DuPont™ ESCORT® XP on a particular grass for the first time, limit use to a small area. In no injury occurs throughout the season, larger acreage may be treated the following season.

Broadleaf forage species, such as alfalfa and clover, are highly sensitive to ESCORT® XP and will be severely stunted or injured by ESCORT® XP.

CROP ROTATION

Before using ESCORT® XP, carefully consider your crop rotation plans and options.

Minimum Rotational Intervals

Minimum rotation intervals* are determined by the rate of breakdown of ESCORT® XP applied. ESCORT® XP breakdown in the soil is affected by soil pH, presence of soil microorganisms, soil temperature, and soil moisture. Low soil pH, high soil temperature, and high soil moisture increase ESCORT® XP breakdown in soil, while high soil pH, low soil temperature, and low soil moisture slow ESCORT® XP breakdown.

Of these 3 factors, only soil pH remains relatively constant. Soil temperature, and to a greater extent, soil moisture, can vary significantly from year to year and from area to area. For this reason, soil temperatures and soil moisture should be monitored regularly when considering crop rotations.

* The minimum rotation interval represents the period of time from the last application to the anticipated date of the next planting.

Soil pH Limitations

ESCORT® XP should not be used on soils having a pH above 7.9, as extended soil residual activity could extend crop rotation intervals beyond normal. Under certain conditions, ESCORT® XP could remain in the soil for 34 months or more, injuring wheat and barley. In addition, other crops planted in high-pH soils can be extremely sensitive to low concentrations of ESCORT® XP.

Checking Soil pH

Before using ESCORT® XP, determine the soil pH of the areas of intended use. To obtain a representative pH value for the test area, take several 0" to 4" samples from different areas of the field and analyze them separately. Consult local extension publications for additional information on recommended soil sampling procedures.

BIOASSAY

A field bioassay must be completed before rotating to any crop or grass species/variety not listed in the Rotation Intervals Table, or if the soil pH is not in the specified range, or if the use rate applied is not specified in the table. To conduct a field bioassay, grow test strips of the crop(s) or grass(es) you plan to grow the following year in fields previously treated with ESCORT® XP. Crop or grass response to the bioassay will indicate whether or not to rotate to the crop(s) or grass(es) grown in the test strips. If a field bioassay is planned, check with your local Agricultural dealer or DuPont representative for information detailing the field bioassay procedure.

Rotation Intervals for Overseeding and Renovation

Location	Crop or Grass Species	Maximum ESCORT® XP Rate (oz per A)	Minimum Rotation Interval (months)
AL, AR, FL, GA, KY, LA, MS, NC, OK, SC, TN, TX, VA, WV	Alfalfa, red clover, white clover, sweet clover, bermudagrass, bluegrass, ryegrass, tall fescue	1/10 to 3/10	4
	Wheat (except durum)	1/10 to 3/10	1
	Durum, barley, oat	1/10 to 3/10	10
ALL STATES NOT INCLUDED ABOVE	Red clover, white clover, and sweet clover	1/10 to 2/10	12
	Bermudagrass, bluegrass, ryegrass	1/10 to 2/10	6
	Tall Fescue	1/10 to 2/10	18
	Wheat (except durum)	1/10 to 2/10	1
	Durum, barley, oat	1/10 to 2/10	10
ALL AREAS WITH SOIL PH OF 7.5 OR LESS	Russian wildrye	1/10 to 1/2	1
	Green needlegrass, switchgrass, sheep fescue	1/10 to 1	1
	Meadow brome, smooth brome, alta fescue, red fescue, meadow foxtail, orchardgrass, Russian wildrye, timothy	1/10 to 1	2
	Alkali sacaton, mountain brome, blue grama thickspike wheatgrass	1/10 to 1	1
ALL AREAS WITH SOIL PH OF 7.9 OR LESS	Sideoats grama, switchgrass	1/10 to 1/2	2
	Western wheatgrass	1/10 to 1	2
	Sideoats grama, switchgrass, big bluestem	1/10 to 1	3

When used as directed, there is no grazing or haying restriction for use rates of 1 2/3 ounce per acre or less. At use rates greater than 1 2/3 ounce per acre and up to 3 1/3 ounce per acre, forage grasses may be cut for hay, fodder or green forage and fed to livestock, including lactating animals, 3 days after treatment.

IMPORTANT PRECAUTIONS

- Grass species or varieties may differ in their response to various herbicides. If no information is available, limit the initial use of DuPont™ ESCORT® XP to a small area.
- Components in a grass seed mixture will vary in tolerance to ESCORT® XP so the final stand may not reflect the seed ratio.
- Under certain conditions such as heavy rainfall, high pH, prolonged cold weather, or wide fluctuations in day/night temperatures prior to or soon after ESCORT® XP application, temporary discoloration and/or grass injury may occur. ESCORT® XP should not be applied to grass that is stressed by severe weather conditions, drought, low fertility, water-saturated soils, disease, or insect damage as grass injury may result. Severe winter stress, drought, disease, or insect damage before or following application also may result in grass injury.
- Applications of ESCORT® XP to lands undersown with legumes may cause injury to the legumes. Legumes in a seeding mixture may be severely injured or killed following an application of ESCORT® XP.
- The control of weeds in wheel track areas may be reduced if ground applications are made when dry, dusty field conditions exist. The addition of 2,4-D or MCPA should improve weed control under these conditions.

WEEDS CONTROLLED

1/3 to 1/2 ounce per acre

Annual sowthistle	Goldenrod
Aster	Lambsquarters
Bahiagrass	Marestail/horseweed****
Beebalm	Maximillion sunflower
Bittercress	Miners lettuce
Bitter sneezeweed	Pennsylvania smartweed
Blackeyed-susan	Plains coreopsis
Blue mustard	Plantain
Bur buttercup	Redroot pigweed
Chicory	Redstem filaree
Clover	Rough fleabane
Cocklebur	Shepherd's purse
Common chickweed	Silky crazyweed (locoweed)
Common groundsel	Smallseed falseflax
Common purslane	Smooth pigweed
Common yarrow	Sweet clover
Conical catchfly	Tansymustard
Corn cockle	Treacle mustard
Cow cockle	Tumble mustard
Crown vetch	Wild carrot
Dandelion	Wild garlic
Dogfennel	Wild lettuce
False chamomile	Wild mustard
Fiddleneck tarweed	Woolly croton
Field pennycress	Wood sorrel
Flixweed	Yankeweed

1/2 to 1 ounce per acre

Blackberry	Honeysuckle
Black henbane	Multiflora rose and other wild roses
Broom snakeweed*	Musk thistle***
Buckhorn plantain	Oxeye daisy
Bull thistle	Plumeless thistle
Common crupina	Prostrate knotweed
Common sunflower	Rosering gaillardia
Curly dock	Seaside arrowgrass
Dewberry	Sericea lespezeza
Dyer's woad	Tansy ragwort
Garlic mustard	Teasel
Gorse	Wild caraway
Halogeton	
Henbit	

1 to 2 ounces per acre

Common mullein	Purple loosestrife
Common tansy	Purple scabious
Field bindweed**	Scotch thistle
Greasewood	Scouringrush
Gumweed	Salsify
Houndstongue	Snowberry
Lupine	St. Johnswort
Old world climbing fern (Lygodium)	Sulphur cinquefoil
Perennial pepperweed	Western salsify
Poison hemlock	Whitetop (hoary cress)
	Wild Iris

1 1/2 to 2 ounces per acre

Canada thistle**	Tall larkspur
Dalmation toadflax**	Wild parsnip
Duncecap larkspur	Yellow toadflax**
Russian knapweed**	

3 to 4 ounces per acre

Kudzu

- * Apply fall through spring.
- ** Suppression, which is a visual reduction in weed competition (reduced population or vigor) as compared to untreated areas. Apply as a full coverage spray for best performance.
- *** Certain biotypes of musk thistle are more sensitive to ESCORT® XP and may be controlled with rates of 1/4 to 1/2 ounce per acre. Treatments of ESCORT® XP may be applied from rosette through bloom stages of development.
- **** Certain biotypes of marestail/horsetail are less sensitive to ESCORT® XP and may be controlled by tank mixes with herbicides with a different mode of action.

Problem Weed Control

For broader spectrum control and for use on certain biotypes of broadleaf weeds which may be resistant to ESCORT® XP and herbicides with the same mode of action, the following tank mixes are recommended.

Dicamba + 2,4-D

Weed	Rate of ESCORT® XP (fluid ounces/acre)	Rate of dicamba (fluid ounces/acre)	Rate of 2,4-D (fluid ounces/acre)
Kochia control	1/2	8	16
Spotted knapweed control	1/2	8	16
Rush skeletonweed suppression	1	8	16

TURF, INDUSTRIAL (UNIMPROVED ONLY)

Application Information

ESCORT® XP is recommended for selective weed control in unimproved industrial turf where certain grasses are well established and desired as ground cover. ESCORT® XP is also recommended for the control of certain noxious and troublesome weeds in turf.

In addition to conventional spray equipment, ESCORT® XP may also be applied with invert emulsion equipment. When using an invert emulsion, mix the prescribed rate of ESCORT® XP in the water phase.

Consult the "Weeds Controlled" table to determine which weeds will be controlled by the following recommendations:

Turf Type	Rate of DuPont™ ESCORT® XP (ounces/acre)
Fescue and Bluegrass	1/4 to 1/2
Crested Wheatgrass and Smooth Brome	1/4 to 1
Bermudagrass	1/4 to 2

Application Timing

Applications may be made at anytime of the year, except when the soil is frozen.

When a spring application is made on fescue or bluegrass, a second application may be made during the summer after full seedhead maturation.

Growth Suppression and Seedhead Inhibition (Chemical Mowing)

Application Information

ESCORT® XP is recommended for growth suppression and seedhead inhibition in well established fescue and bluegrass turf at the use rate of 1/4 to 1/2 ounce per acre.

Tank Mix Combination

ESCORT® XP may be tank mixed with "Embark" for improved performance in the regulation of growth and seedhead suppression. Tank mix 1/4 to 1/2 ounce of ESCORT® XP with 1/8 to 1/4 pint of "Embark".

Application Timing

Application may be made after at least 2 to 3 inches of new growth has emerged until the appearance of the seed stalk.

Fescue Precautions:

ESCORT® XP may temporarily stunt tall fescue, cause it to turn yellow, or cause seedhead suppression. To minimize these symptoms, take the following precautions:

- Do not use more than 4/10 ounce per acre of ESCORT® XP.
- Tank mix ESCORT® XP with 2,4-D.
- Use the lowest recommended rate for target weeds.
- Use a non-ionic surfactant at 1/2 to 1 pint per 100 gallons of spray solution.
- Make application later in the spring after the new growth is 5 to 6 inches tall, or in the fall.
- Do not use a surfactant when liquid nitrogen is used as a carrier.
- Do not use a spray adjuvant other than non-ionic surfactant.
- The yields from the first cutting may be reduced due to seedhead suppression resulting from treatment with ESCORT® XP.

IMPORTANT PRECAUTIONS

—INDUSTRIAL TURF ONLY

- An application of ESCORT® XP may cause temporary discoloration (chlorosis) of the grasses. Use the lower recommended rates for minimum discoloration.
- With fescue and bluegrass, sequential applications made during the same or consecutive growth periods (i.e. spring and fall) may result in excessive injury to turf.

- Excessive injury may result when ESCORT® XP is applied to turf that is under stress from drought, insects, disease, cold temperatures (winter injury) or poor fertility.
- ESCORT® XP is not recommended for use on bahiagrass.

BRUSH CONTROL

Application Information

ESCORT® XP is recommended for the control of undesirable brush growing in non-crop areas including grazed areas on these sites. Applications may be made by air, high volume ground application, low volume ground application and ultra-low volume ground application. Except as noted for multiflora rose, ESCORT® XP should be applied as a spray to the foliage.

The application volume required will vary with the height and density of the brush and the application equipment used. Generally, aerial applications will require 15 to 25 gallons of water per acre; high volume ground application will require 100 to 400 gallons of water per acre; low volume ground application will require 20 to 50 gallons of water per acre; and ultra-low volume ground application will require 10 to 20 gallons of water per acre.

Regardless of the application volume and equipment used, thorough coverage of the foliage, particularly the terminal growing points, is necessary to optimize results.

BRUSH SPECIES CONTROLLED

Species	High Volume Rate (ounces/100 gallon)	Broadcast Rate (ounces/acre)
Ash	1—2	1—3
Aspen	1—2	1—3
Black locust	1—2	1—3
Blackberry	1—2	1—3
Camelthorn	1—2	1—3
Cherry	1—2	1—3
Cottonwood	1—2	2—3
Eastern red cedar	1—2	2—3
Elder	1—2	2—3
Elm	1—2	1—3
Firs	3	1—2
Hawthorn	1—2	1—3
Honeysuckle	1—2	1/2—1
Mulberry	1—2	2—3
Multiflora rose	1—2	1—3
Muscadine (wild grape)	1—2	2—3
Oaks	1—2	1—3
Ocean spray (<i>Holodiscus</i>)	1—2	2—3
Osage orange	1—2	2—3
Red maple	1—2	2—3
Salmonberry	1/2—1	1—3
Snowberry	1/2—1	1—3
Spruce (black and white)	3	2—3
Thimbleberry	1/2—1	1—3
Tree of heaven (<i>Ailanthus</i>)	1—2	1—2
Wild roses	1/2—1	1—3
Willow	1/2—1	1—3
Yellow poplar	1/2—1	1—3

For low volume and ultra-low volume ground applications, mix 4 to 8 ounces of ESCORT® XP per 100 gallons of spray solution.

Application Timing

Make a foliar application of the recommended rate of DuPont™ ESCORT® XP during the period from full leaf expansion in the spring until the development of full fall coloration on deciduous species to be controlled.

Coniferous species may be treated at anytime during the growing season.

Spot Treatment

ESCORT® XP is recommended for the control of many species of weeds including noxious/invasive weeds in certain established grasses growing on non-crop areas. Refer to the "Weeds Controlled" section for a listing of susceptible weed species and the application rate per acre per the target weed.

Or, mix one gram of ESCORT® XP per one gallon of water along with a surfactant. Spray to the point of wetting the entire surface of the target weeds, approximately 40 gallons of solution per acre.

Tank Mix Combinations—

ESCORT® XP may be tank mixed with any product labeled for non-crop brush control at the application rates specified on the companion product's label for the pests specified on the product's companion label. Read and follow the label instructions of both products when tank mixing. Follow the most restrictive limitations of any of the product labels being tank mixed.

Low Rate Applications

Imazapyr (2 pound active per gallon)

Combine 1 to 2 ounces of ESCORT® XP with 1 to 4 pints of imazapyr herbicide per acre and apply as a broadcast spray. Aerial applications should use a minimum of 15 gallons per acre spray volume. In addition to species listed above controlled by ESCORT® XP, this combination controls black gum, hophornbeam, sassafras, sweetgum, Vaccinium species, dogwood, myrtle dahoon, hickories, and persimmon.

Picloram (2 pound active per gallon) + Imazapyr (2 pound active per gallon)

Combine 1 to 1 1/2 ounce of ESCORT® XP with 2 to 8 fluid ounces of imazapyr and 1 to 2 pints of picloram per 100 gallons of water. Apply as a high volume spray. This tank mix controls cherry, elms, box elder, maples, hackberry, redbud, ash, oaks (including shingle oak), black locust and sassafras.

*Picloram is a restricted use pesticide.

Spotgun Basal Soil Treatment

For control of multiflora rose, prepare a spray suspension of ESCORT® XP by mixing 1 ounce per gallon of water. Mix vigorously until the ESCORT® XP is dispersed and agitate periodically while applying the spray suspension.

Apply the spray preparation with an exact delivery handgun applicator. Apply at the rate of 4 milliliters for each 2 feet of rose canopy diameter. Direct the treatment to the soil within 2 feet of the stem union. When treating large plants and more than one delivery is required, make applications on opposite sides of the plant.

Applications should be made from early spring to summer.

IMPORTANT PRECAUTIONS

—NON-CROP BRUSH ONLY

- When using tank mixtures of ESCORT® XP with companion herbicides, read and follow all use instructions, application rates, warnings and precautions appearing on the labels. Follow the most restrictive label instructions for each of the herbicides used.

SPRAY EQUIPMENT

Low rates of ESCORT® XP can kill or severely injure most crops. Following an ESCORT® XP application, the use of spray equipment to apply other pesticides to crops on which ESCORT® XP is not registered may result in their damage. The most effective way to reduce this crop damage potential is to use dedicated mixing and application equipment.

The selected sprayer should be equipped with an agitation system to keep ESCORT® XP suspended in the spray tank. Use a sufficient volume of water to thoroughly cover the foliage of undesirable weeds, generally 10 to 40 gallons per acre. Select a spray volume and delivery system that will deliver a uniform spray pattern. Be sure the sprayer is calibrated before use. Avoid overlapping and shut off spray booms while starting, turning, slowing or stopping to avoid injury to desired plants.

Refer to the brush control section of this label for information unique to that particular use.

MIXING INSTRUCTIONS

1. Fill the tank 1/4 to 1/3 full of water.
2. While agitating, add the required amount of ESCORT® XP.
3. Continue agitation until the ESCORT® XP is fully dispersed, at least 5 minutes.
4. Once the ESCORT® XP is fully dispersed, maintain agitation and continue filling tank with water. ESCORT® XP should be thoroughly mixed with water before adding any other material.
5. As the tank is filling, add tank mix partners (if desired) then add the necessary volume of nonionic surfactant. Always add surfactant last.
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. ESCORT® XP spray preparations are stable if they are pH neutral or alkaline and stored at or below 100° F.
8. If ESCORT® XP and a tank mix partner are to be applied in multiple loads, pre-slurry the ESCORT® XP in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the ESCORT® XP.

USE PRECAUTIONS

- Do not drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the product may be washed or moved into contact with their roots, as injury or loss of desirable trees or other plants may result

- Treatment of powdery, dry soil or light, sandy soil when there is little likelihood of rainfall soon after treatment may result in off target movement and possible damage to susceptible crops when soil particles are moved by wind or water. Injury to crops may result if treated soil is washed, blown, or moved onto land used to produce crops. Exposure to DuPont™ ESCORT® XP may injure or kill most crops. Injury may be more severe when the crops are irrigated. Do not apply ESCORT® XP when these conditions are identified and powdery, dry soil or light or sandy soils are known to be prevalent in the area being treated.
- Applications made where runoff water flows onto agricultural land may injure crops. Applications made during periods of intense rainfall, to soils saturated with water, to surfaces paved with materials such as asphalt or concrete, or to soils through which rainfall will not readily penetrate may result in runoff and movement of ESCORT® XP. Do not treat frozen soil. Treated soil should be left undisturbed to reduce the potential for ESCORT® XP movement by soil erosion due to wind or water.
- Do not use on lawns, walks, driveways, tennis courts or similar areas.
- Do not apply through any type of irrigation system.
- When used as directed, there are no grazing or haying restrictions for use rates of 1 2/3 ounce per acre and less. At use rates of 1 2/3 to 3 1/3 ounce per acre, forage grasses may be cut for hay, fodder or green forage and fed to livestock, including lactating animals, 3 days after treatment.
- Do not use this product in the following counties of Colorado: Saguache, Rio Grande, Alamosa, Costilla and Conejos.
- Do not use this product in California.

SPRAYER CLEANUP

Spray equipment must be cleaned before ESCORT® XP is sprayed. Follow the cleanup procedures specified on the labels of previously applied products. If no directions are provided, follow the six steps outlined below.

At the End of the Day

When multiple loads of ESCORT® XP herbicide are applied, it is recommended that at the end of each day of spraying, the interior of the tank be rinsed with fresh water and then partially filled, and the boom and hoses flushed. This will prevent the buildup of dried pesticide deposits that can accumulate in the application equipment.

1. Drain tank; thoroughly rinse spray tanks, boom, and hoses with clean water. Loosen and physically remove any visible deposits.
2. Fill the tank with clean water and 1 gal of household ammonia* (contains 3% active) for every 100 gal of water. Flush the hoses, boom, and nozzles with the cleaning solution. Then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 min. Flush the hoses, boom, and nozzles again with the cleaning solution, and then drain

the tank.

3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
 4. Repeat step 2.
 5. Rinse the tank, boom, and hoses with clean water.
 6. If only ammonia is used as a cleaner, the rinsate solution may be applied back to the crop(s) recommended on this label. Do not exceed the maximum labeled use rate. If other cleaners are used, consult the cleaner label for rinsate disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.
- * Equivalent amounts of an alternate-strength ammonia solution or a DuPont-approved cleaner can be used in the cleanout procedure. Carefully read and follow the individual cleaner instructions. Consult your agricultural dealer, applicator, or DuPont representative for a listing of approved cleaners.

Notes:

1. **Attention:** Do not use chlorine bleach with ammonia, as dangerous gases will form. Do not clean equipment in an enclosed area.
2. Steam-cleaning aerial spray tanks is recommended prior to performing the above cleanout procedure to facilitate the removal of any caked deposits.
3. When ESCORT® XP is tank mixed with other pesticides, all required cleanout procedures should be examined and the most rigorous procedure should be followed.
4. In addition to this cleanout procedure, all precleanout guidelines on subsequently applied products should be followed as per the individual labels.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (>150 - 200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS! See **Wind, Temperature and Humidity**, and **Temperature Inversions** sections of this label.

Controlling Droplet Size - General Techniques

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

Controlling Droplet Size - Aircraft

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- **Boom Length** - The boom length should not exceed 3/4 of the wing or rotor length - longer booms increase drift potential.
- **Application Height** - Application more than 10 ft above the canopy increases the potential for spray drift.

BOOM HEIGHT

Setting the boom at the lowest labeled height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. AVOID GUSTY OR WINDLESS CONDITIONS. Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft

smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

STORAGE AND DISPOSAL

Pesticide Storage: Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Store in a cool, dry place.

Pesticide Disposal: Do not contaminate water, food, or feed by disposal. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: For Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. **For Fiber Sacks:** Completely empty fiber sack by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into manufacturing or application equipment. Then dispose of sack in a sanitary landfill or by incineration if allowed by State and local authorities. **For Fiber Drums With Liners:** Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner. **For Bags Containing Water Soluble Packets:** Do not reuse the outer box or the resealable plastic bag. When all water-soluble packets are used, the outer packaging should be clean and may be disposed of in a sanitary landfill or by incineration, or if allowed by State and local authorities, by open burning. If burned, stay out of smoke. If the resealable plastic bag contacts the formulated product in any way, the bag must be triple rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer wrap as described above. **For Metal Containers (non aerosol):** Triple rinse (or equivalent) the container. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. **For Paper and Plastic Bags:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

The DuPont Oval Logo, DuPont™, ESCORT® and VELPAR® are trademarks or registered trademarks of E.I. duPont de Nemours & Company.

“Embark” is a registered trademark of PBI Gordon Corporation.

SL - 1252A 111507 11-08-07

LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product; crop injury, or; injury to non-target crops or plants.

DuPont does not agree to be an insurer of these risks. **TO THE FULLEST EXTENT PERMITTED BY LAW, WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE FULLEST EXTENT PERMITTED BY LAW, IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

For product information call: 1-888-6-DUPONT

Internet address: <http://cropprotection.dupont.com/>

© 2001-2007 E. I. du Pont de Nemours and Company, Wilmington, Delaware 19898.

All rights reserved.