



# Natural Heritage & Endangered Species Program

Massachusetts Division of Fisheries & Wildlife

Route 135, Westborough, MA 01581

tel: (508) 389-6360; fax: (508) 389-7891

[www.nhesp.org](http://www.nhesp.org)

## Oak Hairstreak

*Satyrrium favonius*

State Status: **Special Concern**

Federal Status: **None**

**Description:** The Oak Hairstreak has dark brown wings above, often with a round, rust-colored patch in the middle of the forewing and small spot of the same color at the anal angle of the hind wing. On the underside, the wings are tan with a white and black postmedial line crossing both wings and forming a “W” near the anal angle of the hind wing; the hind wing has orange submarginal spots and an iridescent blue patch flanked by two black spots at the anal angle. Wingspan is 26-32 mm. The larva is of the typical slug-like lycaenid form, covered with short, dense setae, green with a faint lateral line and oblique dorsolateral dashes.

**Habitat:** In Massachusetts, the Oak Hairstreak inhabits xeric and open oak woodland and barrens on rocky uplands and sandplains. Adult butterflies are often found nectaring in weedy or scrub areas within or near such habitat, including old fields, clearings, powerline or pipeline cuts, abandoned gravel pits, etc. New Jersey tea (*Ceanothus americanus*), dogbane (*Apocynum* spp.), and milkweed (*Asclepias* spp.) are favored nectar sources, although others are used.

**Life History:** Adult butterflies fly in late June and July. Eggs are laid on oak (*Quercus*); the particular oak species used in Massachusetts have not been documented. Eggs overwinter and hatch in the spring; larvae feed on catkins and new foliage, pupating by early June.

**Geographic Range:** The Oak Hairstreak is spottily distributed from southern New England south to Florida and west to Illinois, Colorado, and Arizona. In Massachusetts, this species occurs from Worcester to near Boston and southeastward; as well as in western Massachusetts in the Connecticut River Valley and the southern Berkshires.




*Satyrrium favonius* • MA: Norfolk Co., Canton • 28 Jun 2004 • Photo by M.W. Nelson

### Adult Flight Period in Massachusetts

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec

### Threats

- Habitat loss
- Fire suppression
- Introduced generalist parasitoids
- Insecticide spraying
- Clear-cut timber harvest



Updated December 2012  
M.W. Nelson

Please allow the Natural Heritage & Endangered Species Program to continue to conserve the biodiversity of Massachusetts with a contribution for 'endangered wildlife conservation' on your state income tax form, as these donations comprise a significant portion of our operating budget.