

EIPP COMPREHENSIVE HEALTH ASSESSMENT: Postpartum

Instructions: Complete at intake by the end of second unit rate reimbursed HV (IV-2), if intaked into EIPP following birth, and at key developmental stages by the RN and/or SW.

ID Number: _____

Last Name: _____ First Name: _____ DOB: _____ Age: _____

Street Address: _____ City: _____ State: MA Zip Code: _____

Mailing Address: _____ City: _____ State: MA Zip Code: _____

Phone Number: _____ Alternate Contact: _____

OB-GYN/Midwife Name: _____ OB-GYN/Midwife Phone Number: _____

Prenatal Care Began (circle one): 1st 2nd 3rd Number of Prenatal Visits: _____

Check here if there are no changes to the street and mailing address.

OVERALL LEVEL OF STRENGTH:

USE CLINICAL JUDGEMENT BASED ON ASSESSMENT OF INDICATORS AND COPING STRATEGIES USED					
Initial	Date Completed: _____	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
2 Month	Date Completed: _____	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
4 Month	Date Completed: _____	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
6 Month	Date Completed: _____	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
8 Month	Date Completed: _____	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
12 Month	Date Completed: _____	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)

PROBLEM LIST:

Problem	Date Identified	Actions Taken	Date Resolved

INFANT & POSTPARTUM INFORMATION:

Baby's First Name: _____ Baby's Last Name: _____ Baby's DOB: _____ Baby's Birth weight: _____ Baby's Gender (circle one): M/F

Type of Delivery: _____ Birth Complications: _____ Pediatrician's Name: _____ Pediatrician's Phone Number: _____

Immunizations: HepB: 1 2 3 Dtap: 1 2 3 Hib: 1 2 3 IPV: 1 2 3 PCV7: 1 2 3

HOUSEHOLD COMPOSITION:

Name	Gender (M/F)	# weeks pregnant	Due Date	Plan to Breastfeed (Y/N)	Age	DOB	Relation	Living in Home (Y/N)	English (Y/N)	Concerns
	F						Self	Yes		

KEY ASSESSMENT AREAS:

KAA#1: Access and Utilization of Care				
Indicators	1 (low)	2 (moderate)	3 (high)	
1a: Housing	<ul style="list-style-type: none"> ▪ Repeated Hx /current homelessness ▪ At risk of eviction ▪ No working phone 	<ul style="list-style-type: none"> ▪ Hx/at risk of homelessness ▪ Owes back rent, fuel & utilities ▪ Phone recently disconnected 	<ul style="list-style-type: none"> ▪ Hx of maintaining stable housing ▪ Fuel/Utilities on, paid, & functioning long term ▪ Working telephone available 	
1b: Health	<ul style="list-style-type: none"> ▪ No health insurance ▪ No access to medical or dental care 	<ul style="list-style-type: none"> ▪ Healthy Start or Health Safety Net Fund ▪ At risk of losing health insurance ▪ Some barriers to accessing medical and/or dental care 	<ul style="list-style-type: none"> ▪ MassHealth or private insurance ▪ Receives medical home with regular care ▪ Receives regular dental care 	
1c: Education	<ul style="list-style-type: none"> ▪ Less than 10th grade education ▪ Unable to set educational or career goals 	<ul style="list-style-type: none"> ▪ High school diploma or GED ▪ Sets and pursues short term educational and career goals 	<ul style="list-style-type: none"> ▪ College or advanced degree ▪ Sets and pursues long term educational and career goals 	
1d: Economics	<ul style="list-style-type: none"> ▪ No or minimal income ▪ FOB not contributing financially ▪ Loss of job/school due to pregnancy ▪ No child care available ▪ No transportation ▪ Unable to prioritize or budget 	<ul style="list-style-type: none"> ▪ Unable to meet all basic financial needs ▪ FOB inconsistently contributes financially ▪ Job/school threatened by pregnancy ▪ Sporadic child care available ▪ Sporadic transportation available ▪ Needs assistance budget/prioritizing 	<ul style="list-style-type: none"> ▪ Adequate income for living expenses ▪ FOB contributes financially ▪ Job/school accommodating pregnancy ▪ Adequate child care available ▪ Reliable transportation available ▪ Able to budget and prioritize 	
1e: Community Support	<ul style="list-style-type: none"> ▪ No culturally & linguistically appropriate information about community resources and services available ▪ Overwhelmed and immobilized ▪ Unable to access services ▪ No social supports and isolated 	<ul style="list-style-type: none"> ▪ Limited culturally & linguistically appropriate information about community resources and services available ▪ Hx of difficulty in accessing services ▪ Limited friends and family in area are supportive/helpful when needed 	<ul style="list-style-type: none"> ▪ Has culturally & linguistically appropriate information about community resources and services available ▪ Hx of success in accessing services ▪ Friends and family in area are supportive/ helpful when needed 	
USE CLINICAL JUDGEMENT BASED ON ASSESSMENT OF INDICATORS AND COPING STRATEGIES USED				
Initial Postpartum CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
2 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
4 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
6 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
8 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
12 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
Additional Comments:				

KAA#2: Women's Health				
Indicators	1 (low)	2 (moderate)	3 (high)	
2a: Family Planning	<ul style="list-style-type: none"> No knowledge of BCM Never used/no plan for BCM Unplanned pregnancy Recent STI Unable to negotiate BCM w/ partner 	<ul style="list-style-type: none"> Verbalizes BCM options Used BCM in past Unplanned pregnancy Hx of STI Conflict with partner on BCM use 	<ul style="list-style-type: none"> Success with BCM Planned pregnancy No Hx of STIs Partner supportive of BCM 	
2b: STI's/HIV	<ul style="list-style-type: none"> Never practices safer sex High risk for STI 	<ul style="list-style-type: none"> Usually practices safer sex Moderate risk for STI 	<ul style="list-style-type: none"> Practices safer sex Low risk for STIs 	
2c: Reproductive Health	<ul style="list-style-type: none"> Inadequate PNC No PP appointment No knowledge of normal maternal physical changes in first year of parenthood 	<ul style="list-style-type: none"> Adequate PNC PP appt scheduled Some knowledge of normal maternal physical changes in first year of parenthood 	<ul style="list-style-type: none"> Adequate PNC Keeps PP appointments Knowledgeable of normal maternal physical changes in first year of parenthood 	
2d: Other Women's Health Concerns	<ul style="list-style-type: none"> No knowledge of health concerns PCP not identified Inadequate Hx of physical exam with screenings 	<ul style="list-style-type: none"> Some knowledge of health concerns Identifies PCP Irregular Hx of physical exams with screenings 	<ul style="list-style-type: none"> Understands health concerns Identifies PCP Annual physical exam with screenings-knows results 	
USE CLINICAL JUDGEMENT BASED ON ASSESSMENT OF INDICATORS AND COPING STRATEGIES USED				
Initial Postpartum CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
2 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
4 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
6 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
8 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
12 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
Additional Comments:				

KAA#3: Oral Health			
Indicators	1 (low)	2 (moderate)	3 (high)
3a: Mom's Oral Health	<ul style="list-style-type: none"> Never received dental care No cleaning during pregnancy Irregular brushing Does not use floss No Fluoride 	<ul style="list-style-type: none"> Irregular dental care No cleaning during pregnancy Brushes daily Irregular floss Occasional fluoride in diet 	<ul style="list-style-type: none"> Regular dental care Cleaning during pregnancy Brushes at least 2x day Flosses 1x a day Fluoride in diet
3b: Baby (Newborn to 6 months) oral health	<ul style="list-style-type: none"> Bottle propping observed Gums are not cleaned No fluoride in diet Verbalizes lack of knowledge of role of fluoride and strong teeth 	<ul style="list-style-type: none"> May bottle prop occasionally Gums cleaned irregularly Occasional fluoride in diet Verbalizes some knowledge of role of fluoride in making strong teeth 	<ul style="list-style-type: none"> No bottle propping Gums are cleaned daily Fluoride in diet Verbalizes role of fluoride in making strong teeth
3c: Baby (6-12 months) oral health	<ul style="list-style-type: none"> Gums and teeth are not cleaned Sippee cup not introduced at 6 months Baby walking around with bottle No knowledge of bottle weaning at 12-14 months 	<ul style="list-style-type: none"> Gums and teeth cleaned irregularly Sippee cup introduced after 6 months Baby walk with bottle occasionally Some knowledge of benefits of bottle weaning at 12-14 months 	<ul style="list-style-type: none"> Cleans baby's gums or teeth daily Sippee cup introduced at six months Baby does not walk with bottle Verbalizes benefits of bottle weaning at 12-14 months

USE CLINICAL JUDGEMENT BASED ON ASSESSMENT OF INDICATORS AND COPING STRATEGIES USED				
Initial Postpartum CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
2 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
4 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
6 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
8 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
12 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
Additional Comments:				

KAA#4: Nutrition				
Indicators	1 (low)	2 (moderate)	3 (high)	
4a: Maternal Nutrition and Weight Gain	<ul style="list-style-type: none"> ▪ BMI ▪ Lacks food representing most of the food groups for meeting family's needs ▪ Major barriers to securing, preparing, and/or feeding infant healthy foods 	<ul style="list-style-type: none"> ▪ BMI ▪ May lack food representing some of the food groups for meeting family's needs ▪ Some barriers to securing, preparing, and/or feeding infant healthy foods 	<ul style="list-style-type: none"> ▪ BMI ▪ Adequate food representing all food groups for meeting family's needs ▪ No barriers to securing, preparing, and/or feeding infant healthy foods 	
4b: Infant Nutrition and fluid intake	<ul style="list-style-type: none"> ▪ Major concerns for infant weight loss/gain ▪ Inadequate response to feeding cues ▪ Infant brought to ER or PEDI for dehydration ▪ Does not show proper preparation or storage of breastmilk or formula ▪ No knowledge of impact of infant growth spurts on feeding and sleeping patterns 	<ul style="list-style-type: none"> ▪ Some concerns for infant weight loss/gain ▪ Inconsistent response to feeding cues ▪ Willing to learn S+S of dehydration and # of wet diapers ▪ Some knowledge of how to store breastmilk or formula ▪ Some knowledge of impact of infant growth spurts on feeding/ sleeping 	<ul style="list-style-type: none"> ▪ Adequate infant weight gain (4-6 oz/wk) ▪ Adequately responds to feeding cues ▪ Knows proper hydration (4-8 wet diapers/day) ▪ Knows proper preparation and storage of breastmilk or formula ▪ Knowledge of impact of infant growth spurts on feeding and sleeping patterns 	
USE CLINICAL JUDGEMENT BASED ON ASSESSMENT OF INDICATORS AND COPING STRATEGIES USED				
Initial Postpartum CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
2 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
4 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
6 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
8 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
12 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
Additional Comments:				

KAA#5: Breastfeeding				
Indicators	1 (low)	2 (moderate)	3 (high)	
5a: Breastfeeding	<ul style="list-style-type: none"> Barriers to accessing social or cultural support for breastfeeding Lack of knowledge of breast care 	<ul style="list-style-type: none"> Some barriers to accessing social or cultural supports for breastfeeding Some knowledge of breast care 	<ul style="list-style-type: none"> No Barriers to accessing social or cultural supports for breastfeeding (LaLeche) Knowledge of breast care and S+S of infection 	
USE CLINICAL JUDGEMENT BASED ON ASSESSMENT OF INDICATORS AND COPING STRATEGIES USED				
Initial Postpartum CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
2 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
4 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
6 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
8 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
12 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
Additional Comments:				

KAA#6: Physical Activity				
Indicators	1 (low)	2 (moderate)	3 (high)	
6a: Maternal Physical Activity	<ul style="list-style-type: none"> Lack of knowledge of importance of physical activity for self/ family Lack of knowledge of interrelationship between diet and physical activity No regular physical activity, physician approves activity Watches television more than 2 hours a day No help for physical condition Has not lost pregnancy weight 	<ul style="list-style-type: none"> Some knowledge of importance of physical activity for self/family Some knowledge of interrelationship between diet and physical activity Sporadic physical activity upon physician approval Watches television about 2 hours a day Some help for physical condition Lost 50% of pregnancy weight 	<ul style="list-style-type: none"> Knowledge of importance of physical activity for self/family Knowledge of interrelationship between diet and physical activity Participates in regular physical activity upon physician approval Watches television less than 2 hours a day Adequate help for physical condition Has lost pregnancy weight 	
6b: Baby Physical Activity	<ul style="list-style-type: none"> Does not create safe environment and does not encourage activities which facilitate development of motor skills Lacks knowledge of role of physical activity in facilitating healthy weight, growth and development 	<ul style="list-style-type: none"> Working to create safe environment and encourages activities which facilitate development of motor skills Some knowledge of role of physical activity in facilitating healthy weight, growth and development 	<ul style="list-style-type: none"> Creates safe environment and encourages activities which facilitate development of motor skills Understands role of physical activity in facilitating healthy weight, growth and development 	
USE CLINICAL JUDGEMENT BASED ON ASSESSMENT OF INDICATORS AND COPING STRATEGIES USED				
Initial Postpartum CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
2 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
4 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
6 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
8 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
12 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
Additional Comments:				

KAA#7: Cognitive and Perceptual				
Indicators	1 (low)	2 (moderate)	3 (high)	
7a: Cognitive and Perceptual	<ul style="list-style-type: none"> Limited cognitive and perceptual abilities; difficulty understanding and using new information Unable to verbalize methods of learning that are most effective for her Verbalizes/demonstrates no confidence in problem solving abilities; unable to describe rationale behind decisions made 	<ul style="list-style-type: none"> May have limited cognitive and perceptual abilities; able to understand and use new information Verbalizes some methods of learning that are most effective for her Verbalizes/demonstrates some confidence in problem solving abilities; able to describe rationale behind decisions made 	<ul style="list-style-type: none"> Adequate cognitive and perceptual abilities; able to understand and use new information Verbalizes methods of learning that are most effective for her Verbalizes/demonstrates confidence in problem solving abilities/able to describe rationale behind decisions made 	
USE CLINICAL JUDGEMENT BASED ON ASSESSMENT OF INDICATORS AND COPING STRATEGIES USED				
Initial Postpartum CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
2 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
4 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
6 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
8 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
12 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
Additional Comments:				

KAA#8: Environmental Health and Safety			
Indicators	1 (low)	2 (moderate)	3 (high)
8a: Lead Poisoning	<ul style="list-style-type: none"> Lead in environment Denies dangers of lead poisoning. Precautions not taken Unable or unwilling to access annual lead testing of children 	<ul style="list-style-type: none"> Concern for lead in environment Verbalizes some of the dangers of lead poisoning and some precautions are taken Some barriers or has not accessed annual lead testing 	<ul style="list-style-type: none"> Lead free environment Verbalizes the dangers of lead poisoning and precautions are taken Knows need for annual lead testing of children to age 4
8b: Asthma	<ul style="list-style-type: none"> Asthmatic child present in home ; asthma not controlled Unable to verbalize asthma triggers including mold, pet dander, dust mites, second hand smoke, and cockroaches Precautions not taken 	<ul style="list-style-type: none"> Asthmatic child present in home; asthma somewhat controlled Verbalizes some asthma triggers including mold, pet dander, dust mites, second hand smoke, and cockroaches Some precautions taken 	<ul style="list-style-type: none"> No asthmatic child present in home or present with asthma in control Verbalizes asthma triggers including mold, pet dander, dust mites, second hand smoke, and cockroaches Precautions are taken
8c: Injury Prevention	<ul style="list-style-type: none"> Parent does not know basic First Aid/CPR Emergency numbers not available Does not use seat belt Does not verbalize knowledge of car seat safety and does not plan or cannot obtain approved car seat Does not verbalize knowledge of major childhood injury risks and does not practice prevention Does not verbalize knowledge of common safety hazards in home and does not practice prevention Does not use appropriate hand washing Weapons present and not secured 	<ul style="list-style-type: none"> Parent learning basic First Aid/CPR Some emergency numbers available Inconsistent use of seat belt Learning car seat safety and experiences some barriers in obtaining approved car seat Verbalizes some knowledge of major childhood injury risks and inconsistently practices prevention Verbalizes some knowledge of common safety hazards in home and inconsistently practices prevention Inconsistent hand washing Weapons present but secured appropriately 	<ul style="list-style-type: none"> Parent knows basic First Aid/CPR Emergency numbers available including poison control Consistent use of seat belt Verbalizes car seat safety and plans to obtain approved car seat Verbalizes knowledge of major childhood injury risks and consistently practices prevention Verbalizes knowledge of common safety hazards in home and consistently practices prevention Consistent, appropriate hand washing Weapons not present
8d: Housing	<ul style="list-style-type: none"> Major safety concerns with housing exists Neighborhood unsafe No smoke detectors in place or functioning 	<ul style="list-style-type: none"> Some safety concerns with housing exist Some concerns for safety of neighborhood Some smoke detectors in place and functioning 	<ul style="list-style-type: none"> No major safety concerns with housing Neighborhood is safe Smoke detectors are in place and functioning
8e: Occupational Hazards	<ul style="list-style-type: none"> Exposed to occupational hazards Precautions not taken 	<ul style="list-style-type: none"> Some knowledge of occupational hazards Precautions taken inconsistently 	<ul style="list-style-type: none"> No exposure to occupational hazards Precautions taken

USE CLINICAL JUDGEMENT BASED ON ASSESSMENT OF INDICATORS AND COPING STRATEGIES USED				
Initial Postpartum CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
2 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
4 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
6 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
8 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
12 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
Additional Comments:				

KAA#9: Alcohol, Tobacco, and Other Drugs				
<i>Assess substance use asking the following questions. Ask her each question and then circle her response.</i>				
1. Have you smoked any cigarettes in the past 3 months?	Yes	No	Decline	
2. Are you or your baby regularly exposed to secondhand smoke?	Yes	No	Decline	
3. Do/did your parents have any difficulties in their lives due to alcohol or drug use?	Yes	No	Decline	
4. Do/did any of your friends have any difficulties in their lives due to alcohol or drug use?	Yes	No	Decline	
5. Does/did your partner have any difficulties in his/her life due to alcohol or drug use?	Yes	No	Decline	
6. In the past, have you had any difficulties in your life due to alcohol or drug use?	Yes	No	Decline	
7. In the past month, have you used drugs?	Yes	No	Decline	
8. In the past month, have you had any alcoholic drinks	Yes	No	Decline	
9. On days when you drink alcohol, about how many drinks do you have?	0	1	2	3 4 5+
10. In a typical week, about how many days do you drink alcohol?	0	1	2	3 4 5+
11. In the past 6 months, have there been any times when you had 4 or more alcoholic drinks in a 2 hour span of time?	Yes	No	Decline	
12. Did you smoke, drink, or use drugs at any time during your pregnancy?	Yes	No	Decline	

Initial if done PP (Check one):

- positive screen, client use
- positive screen for peers, partners, past - no current use
- negative screen

2 Month (Check one):

- positive screen, client use
- positive screen for peers, partners, past - no current use
- negative screen

4 Month (Check one):

- positive screen, client use
- positive screen for peers, partners, past - no current use
- negative screen

6 Month (Check one):

- positive screen, client use
- positive screen for peers, partners, past - no current use
- negative screen

8 Month (Check one):

- positive screen, client use
- positive screen for peers, partners, past - no current use
- negative screen

12 Month (Check one):

- positive screen, client use
- positive screen for peers, partners, past - no current use
- negative screen

Indicators	1 (low)	2 (moderate)	3 (high)
9a: A/OD Screen	<ul style="list-style-type: none"> Positive screen, client use Use significantly impacts family functioning or precipitating family crisis 	<ul style="list-style-type: none"> Positive screen for Peers, Partners, Past- client has no current use Family use mildly impacts family functioning 	<ul style="list-style-type: none"> Negative Screen
9b: Tobacco	<ul style="list-style-type: none"> Mother smokes Smoking in home exposure to second hand smoke 	<ul style="list-style-type: none"> Hx of smoking Smokers in home smoke outside of the home 	<ul style="list-style-type: none"> No Hx of smoking No use in home

USE CLINICAL JUDGEMENT BASED ON ASSESSMENT OF INDICATORS AND COPING STRATEGIES USED				
Initial Postpartum CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
2 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
4 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
6 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
8 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
12 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)

Additional Comments:

KA#10: Violence		
<p>Complete this screening tool by asking EIPP Participant to think about her current or most recent relationship/partner. Ask her each question and then circle her response. If she answers no to the first two questions and/or yes to the last two questions below, then she screens positive on this screening tool.</p>		<p>Intervention:</p> <ul style="list-style-type: none"> ▪ Educate/support ▪ Thank her for disclosure ▪ Articulate concern for safety and well-being ▪ Offer referral/resources ▪ Follow-up soon ▪ Develop safety plan if appropriate, consult with DV Advocate at SAFELINK at 877-785-2020 ▪ Assess child safety – file 51A if needed, consult with DCF DV Unit at 617-748-2335
<p>Initial if done PP (Check one):</p> <ul style="list-style-type: none"> ▫ IPV endangering safety of family ▫ Positive IPV screen > one year ago and no current risk ▫ Negative IPV screen 		
1. Is my partner kind to me and respectful of my choices?	Yes No	
2. Does my partner help when the baby won't stop crying?	Yes No	
3. Am I ever afraid of my partner (of being hurt, shamed)?	Yes No	
4. Am I ever afraid to leave my baby alone with my partner?	Yes No	

2 Month (Check one):

- IPV endangering safety of family
- Positive IPV screen > one year ago and no current risk
- Negative IPV screen

4 Month (Check one):

- IPV endangering safety of family
- Positive IPV screen > one year ago and no current risk
- Negative IPV screen

6 Month (Check one):

- IPV endangering safety of family
- Positive IPV screen > one year ago and no current risk
- Negative IPV screen

8 Month (Check one):

- IPV endangering safety of family
- Positive IPV screen > one year ago and no current risk
- Negative IPV screen

12 Month (Check one):

- IPV endangering safety of family
- Positive IPV screen > one year ago and no current risk
- Negative IPV screen

Indicators	1 (low)	2 (moderate)	3 (high)
10a: Interpersonal and Family Violence	<ul style="list-style-type: none"> ▪ IPV endangering safety of family ▪ Positive screen within one year ▪ Current abuse 	<ul style="list-style-type: none"> ▪ Positive IPV screen > one year ago and no current risk ▪ No Hx of IPV with current partner ▪ Does not interfere with family functioning 	<ul style="list-style-type: none"> ▪ Negative IPV screen
10b: Child Abuse and Neglect	<ul style="list-style-type: none"> ▪ Children have been or will be placed outside home ▪ No communication or contact between one or both parents and child/ren ▪ Open conflict between parents ▪ Children have witnessed IPV ▪ Parents unable to distinguish between discipline and abuse ▪ Current DCF involvement 	<ul style="list-style-type: none"> ▪ Children may show some aggression or behavioral issues ▪ Parents may have tense relationship but address conflict productively ▪ Parents question ability to set limits consistently and provide structure. ▪ Hx of abuse but have received professional help. ▪ Hx of DCF involvement but resolved 	<ul style="list-style-type: none"> ▪ Children live with one or both parents in stable family ▪ Parents/partners communicate well with each other and with children ▪ Children appear happy and well-adjusted ▪ Parents confident in setting limits ▪ No Hx of abuse/neglect ▪ No Hx of DCF involvement

USE CLINICAL JUDGEMENT BASED ON ASSESSMENT OF INDICATORS AND COPING STRATEGIES USED

Initial Postpartum CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
2 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
4 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
6 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
8 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
12 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)

Additional Comments:

KAA#11: Emotional Health

EPDS Screen: If the EIPP Participant meets the EPDS cut-off score of equal to or greater than 13, a referral to a mental health professional may be warranted including a MBD referral when available.

Initial EPDS Score (if done postpartum): _____
 Offered a referral to MBD (circle one): Yes No MBD Not Available
 Accepted referral to MBD (circle one): Yes No
 If no, check below the reason why a referral was not accepted:

- Not interested
- Too many other services
- Already seeing a therapist
- Other medical priorities/needs
- Other, please explain: _____

2 Month EPDS Score: _____
 Offered a referral to MBD (circle one): Yes No MBD Not Available
 Accepted referral to MBD (circle one): Yes No
 If no, check below the reason why a referral was not accepted:

- Not interested
- Too many other services
- Already seeing a therapist
- Other medical priorities/needs
- Other, please explain: _____

4 Month EPDS Score: _____
 Offered a referral to MBD (circle one): Yes No MBD Not Available
 Accepted referral to MBD (circle one): Yes No
 If no, check below the reason why a referral was not accepted:

- Not interested
- Too many other services
- Already seeing a therapist
- Other medical priorities/needs
- Other, please explain: _____

6 Month EPDS Score: _____
 Offered a referral to MBD (circle one): Yes No MBD Not Available
 Accepted referral to MBD (circle one): Yes No
 If no, check below the reason why a referral was not accepted:

- Not interested
- Too many other services
- Already seeing a therapist
- Other medical priorities/needs
- Other, please explain: _____

8 Month EPDS Score: _____
 Offered a referral to MBD (circle one): Yes No MBD Not Available
 Accepted referral to MBD (circle one): Yes No
 If no, check below the reason why a referral was not accepted:

- Not interested
- Too many other services
- Already seeing a therapist
- Other medical priorities/needs
- Other, please explain: _____

12 Month EPDS Score: _____
 Offered a referral to MBD (circle one): Yes No MBD Not Available
 Accepted referral to MBD (circle one): Yes No
 If no, check below the reason why a referral was not accepted:

- Not interested
- Too many other services
- Already seeing a therapist
- Other medical priorities/needs
- Other, please explain: _____

Indicators	1 (low)	2 (moderate)	3 (high)
11a: Psychosocial or mental health issues including postpartum depression	<ul style="list-style-type: none"> ▪ Score \geq 13 on Emotional Health Screen ▪ Responds positively to Full Screen ▪ No Knowledge of PPD ▪ Changes in appetite, sleep, energy or activity that impact ADL ▪ Hx of depression or hospitalization ▪ Current use of antidepressants ▪ Current or Hx of suicide or expression of wanting to hurt others 	<ul style="list-style-type: none"> ▪ Score 10-12 on Emotional Health Screen ▪ Some knowledge of PPD ▪ Some changes in appetite, sleep, energy or activity, but no major impact on ADL ▪ Hx of depression – treated professionally ▪ No past hospitalization ▪ Past use of antidepressants ▪ Hx of suicidal ideation--treated professionally 	<ul style="list-style-type: none"> ▪ Score 0-9 on emotional health screen ▪ Knowledge of PPD ▪ No changes in appetite, sleep, energy or activity level except as related to normal PP changes ▪ No current or past Hx of depression ▪ No past hospitalizations ▪ No Hx/current use of antidepressant ▪ No Hx/current suicidal ideation or expressions of wanting to hurt others
11b: Maternal Stress and Anxiety	<ul style="list-style-type: none"> ▪ Physical or emotional stress – feeling overwhelmed ▪ Current unhealthy coping mechanism (drugs, unsafe sex, excessive food, tobacco use, uncontrolled anger/violence, rage) 	<ul style="list-style-type: none"> ▪ Some physical or emotional symptoms indicating stress ▪ Hx of poor coping strategies but recognizes/sought professional help 	<ul style="list-style-type: none"> ▪ No physical or emotional symptoms indicating stress response ▪ Current healthy coping strategies ▪ Avoids unhealthy coping mechanisms (drugs, unsafe sex, excessive food, tobacco, violence)

11c: Trauma and Loss	<ul style="list-style-type: none"> Current trauma or loss Poor coping in response to trauma or loss Hx/Current abuse (sexual, physical, verbal) not addressed Witnessed violence 	<ul style="list-style-type: none"> Hx/Current trauma/loss-sought professional help Hx of poor coping in response to trauma/loss-sought professional help Hx of abuse (sexual, physical, verbal)-professional help sought Witnessed violence-professional help sought 	<ul style="list-style-type: none"> No Hx of trauma or loss Healthy coping in response to trauma or loss No Hx of abuse (sexual, physical, verbal) No violence witnessed
11d: Self Concept/Self Perception	<ul style="list-style-type: none"> Poor sense of self/ low self-esteem Negative impact of parenthood on lifestyle 	<ul style="list-style-type: none"> Hx of low self-esteem Some negative impact of parenthood on lifestyle but positive impact outweighs negative impact 	<ul style="list-style-type: none"> Positive sense of self Positive impact of parenthood on lifestyle
11e: Relationships	<ul style="list-style-type: none"> Feels isolated from people, agencies, services or unable to access help with physical, emotional and social needs Feeling alienated/isolated from partner, family, and/or friends 	<ul style="list-style-type: none"> Identifies people, agencies, services for help with physical, emotional and social needs; not always accessed Relationship with partner, family and/or friends is sometimes strained 	<ul style="list-style-type: none"> Identifies people, agencies, services for help with physical, emotional and social needs Communicative, supportive and close relationship with partner, family & friends
11f: Sexual Orientation	<ul style="list-style-type: none"> Feels isolated from people, agencies, services or unable to access help with physical, emotional and social needs Feeling alienated/isolated from family, and/or friends 	<ul style="list-style-type: none"> Identifies people, agencies, services for help with physical, emotional and social needs; not always accessed Relationship with family and/or friends is sometimes strained 	<ul style="list-style-type: none"> Identifies people, agencies, services for help with physical, emotional and social needs Communicative, supportive and close relationship with family and friends

USE CLINICAL JUDGEMENT BASED ON ASSESSMENT OF INDICATORS AND COPING STRATEGIES USED

Initial Postpartum CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
2 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
4 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
6 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
8 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
12 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)

Additional Comments:

KAA#12: Healthy Parenting

Initial if done PP: Complete this screening tool by asking EIPP Participant each question and then circle her response.		
Do you feel that you are getting the support you need from others?	Yes	No
Do you have someone who you can count on to listen to you when you need to talk?	Yes	No
Do you have someone to call when you need someone to care for the baby?	Yes	No
2 Month: Complete this screening tool by asking EIPP Participant each question and then circle her response.		
Do you feel that you are getting the support you need from others?	Yes	No
Do you have someone who you can count on to listen to you when you need to talk?	Yes	No
Do you have someone to call when you need someone to care for the baby?	Yes	No

4 Month: Complete this screening tool by asking EIPP Participant each question and then circle her response.

Do you feel that you are getting the support you need from others? Yes No

Do you have someone who you can count on to listen to you when you need to talk? Yes No

Do you have someone to call when you need someone to care for the baby? Yes No

6 Month: Complete this screening tool by asking EIPP Participant each question and then circle her response.

Do you feel that you are getting the support you need from others? Yes No

Do you have someone who you can count on to listen to you when you need to talk? Yes No

Do you have someone to call when you need someone to care for the baby? Yes No

8 Month: Complete this screening tool by asking EIPP Participant each question and then circle her response.

Do you feel that you are getting the support you need from others? Yes No

Do you have someone who you can count on to listen to you when you need to talk? Yes No

Do you have someone to call when you need someone to care for the baby? Yes No

12 Month: Complete this screening tool by asking EIPP Participant each question and then circle her response.

Do you feel that you are getting the support you need from others? Yes No

Do you have someone who you can count on to listen to you when you need to talk? Yes No

Do you have someone to call when you need someone to care for the baby? Yes No

Indicators	1 (low)	2 (moderate)	3 (high)
12a: Parent's Transition to Parenting	<ul style="list-style-type: none"> Has unrealistic expectations of baby's care needs Unable to make necessary lifestyle changes/ adaptations to meet baby's needs Negative statements about own ability to parent Negative perception of parenting Inadequate nutrition and minimal rest to care for baby Barriers to learning about parenting, child development, nurturing, and bonding 	<ul style="list-style-type: none"> Uncertain expectations of baby's care needs Inconsistently makes necessary lifestyle changes/adaptations to meet baby's needs Lacks confidence in ability to parent Some ambivalence of parenting Inconsistent nutrition and sporadic rest to care for baby Some barriers to learning about parenting, child development, nurturing, and bonding 	<ul style="list-style-type: none"> Realistic expectations of baby's care needs Able to make necessary lifestyle changes/adaptations to meet baby's needs Positive statements about ability to parent Positive perception of parenting Adequate nutrition and rest to care for baby No barriers to learning more about parenting, child development, nurturing, and bonding
12b: Parent-Baby Attachment	<ul style="list-style-type: none"> Early bonding experiences were negative/abusive Lethargic, withdrawn baby Barriers to accurately interpreting baby's signals and cues 	<ul style="list-style-type: none"> Early bonding experiences were indistinct Baby inconsistently alert/responsive Difficulty with accurately interpreting baby's signals and cues 	<ul style="list-style-type: none"> Early bonding experiences were positive Alert, responsive baby Accurately interprets baby's signals and cues

USE CLINICAL JUDGEMENT BASED ON ASSESSMENT OF INDICATORS AND COPING STRATEGIES USED

Initial Postpartum CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
2 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
4 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
6 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
8 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
12 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)

Additional Comments:

KAA#13: Neonatal and Developmental Assessment Using Ages & Stages Questionnaire				
USE CLINICAL JUDGEMENT BASED ON ASSESSMENT OF INDICATORS AND COPING STRATEGIES USED				
Initial Postpartum CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)

2 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
4 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
6 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
8 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
12 Month CHA	0 (Unable to Assess)	1 (low)	2 (moderate)	3 (high)
Additional Comments:				

KAA#14: Medical History and Physical Assessment												
Medical History:												
Allergies:			Current Medications:			Risk: Average?		Date of Last CBE:		Date of Last Mammogram:		
Medical Conditions: CAD HTN Diabetes Seizures Asthma Urinary Anemia/Coag Thyroid Dx Osteoporosis Mental Illness Surgeries Cancers Other: _____												
Physical Assessment:												
Age:		GR:	P:	EDC:		Height:	Weight:	B/P (if indicated):		HR:	RR:	Temp:
Skin: NP Edema Lesions Erythema Pruritus					GI: NP Anorexia Nausea Vomiting Constipation			GU: NP Burning Frequency Incont. Painful Color Clarity				
Diet:				Birth Defects/Genetic Susceptibility:				BC Plans:				
Mobility Concerns:						Any Risk Factors:						
Postpartum Maternal:												
C-Section: N/A NP Erythema Drainage Painful				Episiotomy: N/A NP Erythema Drainage Painful			Breasts: NP Supportive Bras Engorged Erythema Painful Cracked					
Lochia: Rubra (1-4 days) Serosa(5-7 days) Alba(1-3wks)			Repro: NP Increased or changed discharge Odor Bleeding			Concerns:						
2 – 12 months Postpartum:												
Date Postpartum F/U complete:			Repro: NP Increased or changed discharge Odor			Contraception Method: w/o problems						
Infant Assessment:												
AHR:		RR:	Temp (if indicated):		Weight gain since birth: Normal At-risk		Tone: WNL	General appearance: WNL		Tone: WNL	Reflexes:	
Skin: NP Well Hydrated			Cap refill/Temp/Turgor: WNL		Umbilicus: N/A NP Erythema Drainage Edema			Circumcision: N/A NP Erythema Drainage Edema				
GI: NP Colic Vomiting Constipation Jaundice				GU: Voiding WNL Stool WNL			Risk factors for sensorineural hearing impairment:					
USE CLINICAL JUDGEMENT BASED ON ASSESSMENT OF INDICATORS AND COPING STRATEGIES USED												
Initial Postpartum CHA				0 (Unable to Assess)		1 (low)		2 (moderate)		3 (high)		
2 Month CHA				0 (Unable to Assess)		1 (low)		2 (moderate)		3 (high)		
4 Month CHA				0 (Unable to Assess)		1 (low)		2 (moderate)		3 (high)		
6 Month CHA				0 (Unable to Assess)		1 (low)		2 (moderate)		3 (high)		
8 Month CHA				0 (Unable to Assess)		1 (low)		2 (moderate)		3 (high)		
12 Month CHA				0 (Unable to Assess)		1 (low)		2 (moderate)		3 (high)		
Additional Comments:												