

Sharps Injuries among Hospital Workers in Massachusetts, 2004

Findings from the Massachusetts Sharps Injury Surveillance System

Deval L. Patrick, Governor
Timothy P. Murray, Lieutenant Governor
JudyAnn Bigby, MD, Secretary, Executive Office of Health and Human Services
John Auerbach, Commissioner of Public Health

Letitia K. Davis, Sc.D., Director, Occupational Health Surveillance Program
**Alfred DeMaria, Jr., M.D., Assistant Commissioner, Bureau of Communicable Disease
Control**

April 2007

Acknowledgements

This report was prepared by Angela K. Laramie, MPH, Letitia K. Davis, Sc.D., Natalia Firsova, MA, and James Laing, BS, of the Occupational Health Surveillance Program, Alfred DeMaria, Jr., MD, of the Bureau of Communicable Disease Control, and Laurie M. Robert, MS, of John Snow Inc., who serves as a consultant to this project. Special acknowledgement goes to the members of the Massachusetts Department of Public Health Sharps Injury Prevention Advisory Committee who have dedicated substantial time and effort to guide the development of the Massachusetts Sharps Injury Surveillance System and the preparation of this report. In addition to Alfred DeMaria, Jr., MD, these include: Marie-Eileen Onieal, RNC, Gail Palmeri, RN, Phillip Adamo, MD, Evelyn Bain, RN, Karen Daley, RN, Anuj Goel, Margaret Quinn, Sc.D., and James Ryan, MD, MPH. Additional thanks to Helene Bednarsh, RDH, Catherine Galligan, MS, and Liz O'Connor, RN; who provided invaluable technical expertise and practical insights. Finally, special thanks go to the infection control, employee health department and other staff in Massachusetts hospitals who collected and provided the data on which this report is based. Many hospital staff have provided helpful input in developing the reporting system and continued input is welcome.

This report is dedicated to the memory of Dr. James Ryan, Occupational Medicine Physician, for his passionate work to protect the health and safety of workers, particularly those in the healthcare field.

This work was funded in part through a cooperative agreement with the National Institute for Occupational Safety and Health (U60/OH008490) of the Centers for Disease Control and Prevention.

To obtain additional copies of this report, contact:

Massachusetts Department of Public Health
Center for Health Information, Statistics, Research and Evaluation
Occupational Health Surveillance Program
250 Washington Street, 6th Floor
Boston, MA 02108

617-624-5632

Sharps.Injury@state.ma.us

This report is also available on line at MDPH's website:
www.mass.gov/dph/ohsp

Suggested citation:

Massachusetts Department of Public Health Occupational Health Surveillance Program. (2007)
Sharps Injuries among Hospital Workers in Massachusetts, 2004: Findings from the
Massachusetts Sharps Injury Surveillance System.

Contents

	Page
Executive Summary	1
Introduction	3
Methods	6
Results	9
Work Status of Injured Worker	10
Occupation of Injured Worker	11
Department where Injury Occurred	12
Occupation of Injured Worker by Department where Injury Occurred	13
Procedure for which Sharp was Used or Intended	14
Device Involved in the Injury	15
Device Involved in the Injury by Occupation of Injured Worker	16
Device Involved in the Injury by Department where Injury Occurred	17
Safety Devices	18
Standard v. Safety Device by Type of Device Involved in the Injury	19
Manufacturer/Brand of Device Involved in the Injury	20
When the Injury Occurred: Before, During, After Use of Device	21
How the Injury Occurred	22
Experimental Sharps Injury Rates by Number of Licensed Hospital Beds	23
Discussion	24
References	25
Appendices	
A. Members of the MDPH Sharps Injury Prevention Advisory Committee	27
B. Massachusetts General Laws: An Act Relative to Needle Stick Injury Prevention	28
C. Code of Massachusetts Regulations	30
D. List of Data Elements required on Sharps Injuries Logs in Massachusetts Hospitals and reportable to MDPH	33
E. MDPH Bloodborne Pathogen Exposure Incident Recording Form and Instructions	34
F. MDPH form: Annual Summary of Sharp Injuries Reporting Form and Instructions	40
G. Detailed Tables of Sharps Injuries among Hospital Workers, All Hospitals	
Table G-1. Work Status of Injured Worker	44
Table G-2. Occupation of Injured Worker	44
Table G-3. Department where Incident Occurred	45
Table G-4. Procedure for which Device was Used	47
Table G-5. Device Involved in the Injury	48
Table G-6. Safety Device	49
Table G-7. When the Injury Occurred	49
Table G-8. How the Injury Occurred	50
H. Detailed Tables of Sharps Injuries among Hospital Workers by Number of Licensed Hospital Beds, All Hospitals	52
I. Detailed Tables of Sharps Injuries among Hospital Workers by Teaching Status, All Hospitals	54
J. List of Selected Resources about Bloodborne Pathogen Exposures for Health Care Workers	56

Executive Summary

Health care worker exposures to bloodborne pathogens as a result of injuries caused by needles and other sharp devices are a significant public health concern. The U.S. Centers for Disease Control and Prevention (CDC) estimate that, nationwide, between 600,000 and 800,000 percutaneous injuries from contaminated sharp devices occur each year in health care; approximately half are sustained by hospital workers.

Sharps injuries are preventable, and health care facilities are required by state and federal regulations to implement comprehensive plans to reduce these injuries. Elements of a successful sharps injury prevention program (as outlined by the CDC) include: promoting an overall culture of safety in the workplace, eliminating the unnecessary use of needles and other sharp devices, using devices with sharps injury prevention features (safety devices), employing safe workplace practices, and training health care personnel. Sharps injury surveillance is also a key component of a comprehensive program.

Prior to 2000, while some national data had been collected, little was known about the extent and distribution of sharps injuries among health care workers at the state level. In 2001, pursuant to the Massachusetts law – An Act Relative to Needlestick Injury Prevention (MGL Chapter 111 §53D) – the Massachusetts Department of Public Health (MDPH) promulgated regulations requiring hospitals to report sharps injury data to MDPH. This led to the establishment of the Massachusetts Sharps Injury Surveillance System.

This third annual report from the Massachusetts Sharps Injury Surveillance System provides information about sharps injuries among Massachusetts hospital workers that occurred in 2004. For all hospitals combined, patterns of sharps injuries by a) occupation of the injured worker, b) department in which the injury occurred, c) procedure for which the device was used, and d) device involved are described. Sharps injury rates¹ (defined as number of sharps injuries per 100 licensed hospital beds) are presented for the state overall and for three hospital size categories (small, medium and large hospitals). The report also provides feedback to hospitals regarding data quality. Results stratified by hospital size and by teaching status are included at the end of the report.

Under-reporting of sharps injuries by employees has been well documented in the literature, and varies by occupation and by hospital. Hospitals with well established sharps injury surveillance programs and strong safety cultures may identify and report more injuries than hospitals with less well developed programs. Under-reporting must be taken into account in interpreting the findings presented in this report. Hospitals, in evaluating their own data, should do so within the context of their own sharps injury surveillance and prevention programs. Assessment of under-reporting should be an integral part of sharps injury prevention activities.

The Massachusetts Sharps Injury Surveillance System is intended to provide information that can assist Massachusetts hospitals and health care workers in targeting and evaluating efforts to reduce the incidence of sharps injuries and the associated human and economic costs. This report illustrates ways in which surveillance data can be used within hospitals to identify prevention priorities. Input from hospitals and health care workers regarding the surveillance activities and the content of this report is welcome. MDPH looks forward to continued collaboration in building an effective sharps injury surveillance system to improve the health and safety of health care workers in Massachusetts.

¹ Rates based on the number of licensed beds have a number of limitations, and should be interpreted with caution. Alternative approaches to calculating rates are being explored for future reports.

Findings:

Overview

- A total of 3,279 sharps injuries among hospital health care workers in Massachusetts were reported for the surveillance period January 1 to December 31, 2004. Ninety-seven percent (3,180) of the injuries were reported by acute care hospitals.
- Eighty-five percent of workers (2,776) who sustained injuries were hospital employees, 9% (310) were non-employee practitioners, 3% (83) were students, and 1% (35) were temporary or contract employees.

Occupation and Department

- Nurses sustained more injuries (1,279, 39%) than any other occupational group followed by physicians, who sustained 33% (1,068) of all reported sharps injuries. Close to half of the injuries in the physician category were sustained by interns and residents. Physicians accounted for proportionately more injuries in large hospitals (> 300 licensed beds).
- Technicians and support service workers were also at risk for sharps injuries. Technicians, such as surgical technicians and phlebotomists, accounted for 657 (20%) injuries. Support service workers sustained 123 (4%) injuries; 78 (2%) were sustained by housekeepers.
- Injuries occurred most frequently in operating and procedure rooms (1,460, 45%) and inpatient units (excluding intensive care units) (732, 22%).

Type of Device

- Hollow bore needles as a group accounted for 56% (1,836) of all injuries reported and proportionately more injuries among nurses (77%) than physicians (33%). More than half of the injuries involving hollow bore needles occurred with hypodermic needles.
- The type of device involved in the incident varied by occupation. Hypodermic needles accounted for the greatest number of injuries (583, 46%) among nurses, whereas suture needles accounted for the greatest number of injuries (456, 43%) among physicians.
- Over half of the injuries (1,773, 54%) involved standard devices that were reported as not having engineered sharps injury prevention features. Hypodermic needles accounted for 23% (416) of the injuries involving devices without safety features, even though hypodermic needles with safety features have been available on the market for the past 12 years.

Procedure for which the Device was Used and When the Injury Occurred

- Devices involved in injuries were most frequently used for injections (754, 23%) and suturing (729, 22%). Proportionately more of the injuries in large hospitals were related to suturing, while in medium hospitals, more injuries were related to injections. In small hospitals, blood procedures resulted in the most injuries.
- Injuries occurred during the use of devices in 42% (1,363) of the cases. After use of the device was a more dangerous time to handle a device. About half (1,603, 49%) of the injuries occurred after use of the device, including injuries sustained after use but before disposal of devices (1,105, 34%) and injuries occurring during or after disposal (498, 15%).

Introduction

Health care worker exposures to bloodborne pathogens as a result of injuries from needles and other sharp devices are a significant public health concern. The U.S. Centers for Disease Control and Prevention (CDC) estimate that, nationwide, there are between 600,000 and 800,000 percutaneous injuries from contaminated needles and other sharp devices (referred to as "sharps injuries" in this report) each year in the health care industry, approximately half of which are sustained by hospital-based health care workers (NIOSH, 1999). This averages out to more than 1,000 percutaneous injuries each day in US hospitals (Panlilio, Cardo, Campbell, Srivastava, Jagger, Orelie, et al., 2000). As a measure of the likelihood of injury among hospital workers, it has been estimated that annually there are 22 sharps injuries for every 100 occupied hospital beds (Perry, Parker & Jagger, 2003).

Sharps injuries have been associated with occupational transmission of hepatitis B (HBV), hepatitis C (HCV) and human immunodeficiency virus (HIV), as well as other pathogens. As of 2000, 25 million individuals in the general population are infected with HBV, 4 million are infected with HCV, and 900,000 with HIV (OSHA, 1998). For many, infection status is not known. The estimated risk of a health care worker developing HCV after each percutaneous exposure to blood or body fluids from an infected patient is estimated to be between 0.4-1.8% (OSHA, 1998). For HIV, the calculated risk is 0.3% (OSHA, 1998). The risk of developing HBV after percutaneous exposure is estimated to be between 6-30% among those workers who have not received HBV vaccinations (OSHA, 1998). HBV vaccination lowers this risk and has been shown to be 80-95% effective in preventing the disease (MMWR, 1982). Since 1992, when the Occupational Safety and Health Administration (OSHA) promulgated the Bloodborne Pathogen Standard, employers have been required to offer the HBV vaccine to employees who may be exposed to blood or potentially bloody body fluids in the course of their jobs. As a result, HBV vaccination rates have increased in recent years, and rates of HBV infection have dropped significantly among health care providers (OSHA, 1998).

The U.S. Public Health Service has recommended guidelines for post-exposure management of all workers who have sustained occupational exposures to bloodborne pathogens (MMWR, 2001; MMWR, 2005). These guidelines provide information for determining when post-exposure prophylaxis is appropriate. Preventive medical treatment following exposure may decrease the likelihood of seroconversion for HIV (Cardo, Culver, Ciesielski, Srivastava, Marcus, Abiteboul, et al., 1997).

While the risk of developing disease after a sharps injury is low, the economic and human costs associated with these injuries are substantial. These include the costs for baseline and follow-up testing of the exposed worker, testing the source patient if serostatus is not already known, and the costs of post-exposure prophylaxis. The costs are estimated to range from \$500 to \$3,000 per incident depending on the treatment provided (Jagger, Bentley & Julliet, 1998). Other direct costs include health care costs when workers develop infection and disease as a result of exposure, overtime to make up for any staffing changes that may result from the injury, and increases in workers' compensation costs. In addition to these direct costs, there are indirect costs, that are more difficult to quantify; including the emotional costs to workers and their families associated with the anxiety about the possible consequences of sharps injuries, as well as other human costs when workers become infected. Also difficult to quantify are the effects of sharps injuries on morale of workers, turnover, and perceptions of quality of care within the hospital.

Sharps injuries are preventable, and the OSHA Bloodborne Pathogen Standard requires all health care facilities to have comprehensive plans in place to reduce sharps injuries and other bloodborne pathogen exposures. According to the CDC, sharps injuries can be prevented by: promoting a culture of safety in the work environment; eliminating the unnecessary use of needles and other

sharps devices; using devices with sharps injury prevention features; using safe work practices; and educating and training health care personnel (CDC, 2004). Surveillance of sharps injuries sustained by workers is also a critical component of a comprehensive prevention strategy. Information about the types of devices and procedures associated with sharps injuries, the departments in which the injuries occurred, and the occupations at risk is essential to developing effective prevention programs in health care facilities, and at the state and national levels.

Surveillance of Sharps Injuries among Health Care Workers

Currently, there are two national surveillance systems for tracking sharps injuries to health care workers: The National Surveillance System for Health Care Workers (NaSH), operated by the Centers for Disease Control and Prevention (<http://www.cdc.gov/ncidod/hip/SURVEILL/nash.htm>); and EpiNet which is operated by the International Health Care Worker Safety Center at the University of Virginia (www.med.virginia.edu/epinet).

NaSH is a voluntary reporting system with approximately 20 (mostly teaching) hospitals throughout the country. NaSH has collected data since 1995 on vaccine preventable diseases, bloodborne pathogen exposures, and tuberculosis exposures. EpiNet is also voluntary and has collected data regarding occupational bloodborne pathogen exposures since 1992; approximately 70 hospitals in three geographic regions report exposure data through EpiNet.

Surveillance of sharps injuries is limited by the fact that health care workers often fail to report sharps injuries to their employers. NaSH and EpiNet have estimated the under-reporting rate for sharps injuries to be 56% and 39% respectively (Perry, 2000). There are many reasons why health care workers may not report sharps injuries; they may perceive that the injuries or the source patients are low risk; they may fear the diseases to which they have potentially been exposed; they may have concerns about job security or the extra paperwork and time involved in follow-up (Tandberg, Stewart & Doezema, 1991). In addition, they may lack information and training about appropriate reporting procedures or the reporting procedures themselves may be inadequate. Under-reporting should be taken into account in interpreting sharps injury surveillance data.

Although these two national reporting systems are in place, and a few states in addition to Massachusetts and California have released data, there is still a lack of comprehensive information about sharps injuries among health care workers at the state level. State specific data are important to inform state prevention activities and promote action at the local level. Statewide surveillance of sharps injuries can provide important information about trends in sharps injuries and the devices, procedures, and departments associated with sharps injuries. It can identify health care facilities where increased intervention efforts are needed. Statewide surveillance can also identify facilities where prevention efforts have been effective, and facilitate sharing of information about successful programs and practices.

The Massachusetts Sharps Injury Surveillance System

Work-related sharps injuries potentially affect the lives of many individuals in Massachusetts. The health care industry employs almost 390,000 people in the Commonwealth, more than any other industrial sector (Massachusetts Department of Workforce Development, 2004). Forty-four percent of Massachusetts health care workers are employed in hospitals, including over 80,000 health care practitioners and technical occupations, as well as thousands of others who perform other important functions in the hospital setting (Massachusetts Department of Workforce Development, 2005). Notably, the risk of sharps injury is not limited to direct care providers, but also affects support staff such as maintenance and environmental service workers. When sharps devices are improperly disposed of, many people, including patients and visitors, are placed at risk.

In 2000, Massachusetts joined a growing number of states that have enacted state laws to prevent sharps injuries among health care workers. The Massachusetts law--An Act Relative to Needlestick Injury Prevention (MGL Chapter 111 §53D)--requires all Massachusetts hospitals licensed by the Massachusetts Department of Public Health (MDPH) (except state facilities) to (See Appendix B for MGL Chapter 111 §53D):

- Utilize sharps with engineered sharps injury prevention features to the extent feasible;
- Develop written exposure control plans that include effective procedures for identifying and selecting existing sharps injury prevention technology;
- Record percutaneous exposure incidents in sharp injury logs (including information about the type and brand of device involved in the incidents);
- Use this information for continuous quality improvement in reducing sharps injuries through education and procurement of improved products; and
- Report information from sharps injury logs annually to MDPH.

The Massachusetts law also calls for the formation of an advisory committee at MDPH to address sharps injuries, and the compilation of a list of safer sharps devices to be maintained by MDPH. (See Appendix A for current Advisory Committee membership.)

Shortly after the enactment of MGL Chapter 111 §53D, Congress mandated OSHA to amend the existing Bloodborne Pathogens Standard (29 CFR 1910.1030) to include provisions explicitly requiring employers to use safer sharps devices, to record percutaneous injuries on Sharps Logs and to utilize this information for quality improvement.

In 2001, regulations pursuant to MGL Chapter 111 §53D went into effect requiring hospitals to record sharps injuries (also referred to as "reportable exposure incidents" as defined below) on Sharps Injury Logs starting October 1, 2001 (See Appendix C for 105 CMR 130.1001 *et seq.*). The MDPH regulations implementing the state law mirror federal law regarding use of safe devices and recording sharps injuries, and they add the requirement that MDPH licensed hospitals submit the data from their Sharps Injury Logs annually to the Department. The initial reporting period was defined as October 1, 2001 – December 31, 2001. The first Annual Summaries of Sharps Injuries, to include data from this period, were due at MDPH on February 1, 2002. The subsequent reporting periods cover the full calendar year.

This third annual report from the Massachusetts Sharps Injury Surveillance System uses data reported by licensed hospitals (except state facilities) and provides a look at sharps injuries among Massachusetts hospital workers from January 1, 2004 through December 31, 2004. This picture will be augmented in the future, as more data become available. This report includes information regarding the devices and procedures associated with sharps injuries in Massachusetts hospitals as well as the departments in which these injuries occurred and the occupations involved. Findings are presented by hospital bed-size categories and by teaching status as well as for the state as a whole to allow hospitals to compare their individual experiences with those in similar facilities. Several data quality issues are discussed. Data from the Sharps Injury Surveillance System are intended to assist hospitals and health care workers in targeting and evaluating their efforts to prevent sharps injuries. Feedback from hospitals and health care workers regarding the content and format of this report is welcome, and it will be taken into account in preparing future reports.

Methods

Reportable Exposure Incident: A reportable exposure incident is a bloodborne pathogen exposure incident that is the result of events involving a contaminated sharp device that pierces the skin or mucous membranes. It is also referred to in this report as a “reportable sharps injury”. Bloodborne pathogen exposure is defined more broadly as a specific eye, mouth or other mucous membrane, non-intact skin, or parenteral contact with blood or other potentially infectious materials that result from the performance of an employee’s duties. A sharp is defined as any object that can penetrate the skin or any part of the body and result in an exposure incident, including but not limited to needle devices, scalpels, lancets, broken glass, and broken capillary tubes.

Population Under Surveillance: All health care workers in acute and non-acute care hospitals licensed by MDPH, as well as any satellite units (e.g., community health centers, ambulatory care centers) operating under a hospital license, are included in the population under surveillance. These health care workers include hospital employees, employees of other agencies working in the hospital, those providing patient services without compensation such as students, and anyone providing care within the facility, regardless of the source of their compensation.

Surveillance Period: The surveillance period is defined as January 1 through December 31, 2004. MDPH regulations require that sharps injury data be submitted by licensed hospitals to MDPH (except for state facilities) by February 1 for the previous calendar year.

Definitions:

Sharps Injury Prevention Technology: Sharps injury prevention technology is defined as devices or other technology that minimize the risk of injury to health care workers from hypodermic syringes, needles or other sharps. OSHA refers to non-needle sharps and needle devices used for withdrawing body fluids, accessing a vein or artery, or administering medications or other fluids, with built-in safety features or mechanisms that effectively reduce the risk of an exposure incident as “sharps with engineered sharps injury protections” (SESIPs). They are referred to in this report as “safety devices”.

Teaching hospital: Teaching hospitals are defined by the Medicare Payment Advisory Commission as hospitals with at least 25 medical residents per 100 hospital beds. Hospitals self-designate as teaching hospitals when reporting financial records to the Massachusetts Division of Health Care Finance and Policy.

Data collection methods: Prior to implementing the record-keeping and annual reporting requirements, MDPH worked with members of its Sharps Injury Prevention Advisory Committee to develop effective mechanisms for collecting and reporting sharps injury data. MDPH identified data elements to be recorded on the sharps injury log, consistent with OSHA record-keeping requirements. Several additional data elements were recommended (Appendix D). To facilitate collection of standard data and reduce the need for coding narrative text at both the hospital and state levels, MDPH developed a recommended Bloodborne Pathogen Exposure Incident Recording Form that includes lists of device types, procedures, clinical practice settings, occupations, and how the injury occurred (Appendix E). Based on lists developed for NaSH, these standard lists allow data from Massachusetts to be compared with national data. Use of the Bloodborne Pathogen Exposure Incident Recording Form was voluntary. MDPH also developed a mandatory Annual Summary of Sharps Injury reporting form (referred to in this report as the Annual Summary) that included the same standard lists (Appendix F). Hospitals have the option of submitting this form either as hard copy or electronically.

For most hospitals, information from Sharps Injury Logs was submitted to the Occupational Health Surveillance Program at MDPH by infection control practitioners or employee health staff. In some

hospitals, reports were submitted by staff in risk management or human resources. Data from the Annual Summaries were entered at MDPH into an MSEXcel spreadsheet, imported into MSAccess and coded as needed using the standard lists developed for NaSH (See Appendix F). Expert clinicians assisted in making coding decisions, and data were then imported into SAS for analysis.

Experimental Sharps Injury Rates: Sharps injury rates indicate the probability or risk of a worker sustaining a sharps injury within the surveillance period. Numbers are the counts of sharps injury cases. A large hospital may have many workers who sustain sharps injuries but the rate of injury may be low. Conversely, in a smaller hospital, relatively few workers may sustain sharps injuries but the risk may be high. Both rates and numbers of injuries must be considered when targeting and evaluating prevention efforts.

Sharps injury rates presented in this report are defined as the number of reported sharps injuries divided by the number of licensed hospital beds. Information regarding bed numbers for each hospital was obtained from the MDPH Division of Health Care Quality that licenses hospitals. Rates were calculated for all hospitals combined, as well as by hospital size. Hospitals were divided into three groups based on the number of licensed beds - small (0-100 beds), medium (101-300 beds) and large (301+ beds) for this analysis. Rates by hospital size were calculated by adding all injuries reported in each category (small, medium, and large hospitals) and dividing by the total number of licensed beds in the respective category. The use of licensed beds as a denominator is a preliminary step in the process to determine the most appropriate denominator to be used for rate calculations. Currently, additional denominators, such as number of occupied beds, number of procedures, number of patients, and number of full-time equivalent employees, are being evaluated for use in rate calculations.

Limitations

There are a number of data limitations that need to be taken into account when interpreting sharps injury rates. Optimally, sharps injury rates would be calculated using information on the total number of hours worked, sharps devices purchased or used, or procedures performed at the hospitals in the denominator. This information, however, was not available. Rates based on numbers of licensed beds are approximate measures of risk, and are included in this report to allow hospitals to compare their injury experience with that of other hospitals in same size categories. However, it should be recognized that the number of licensed beds is neither an accurate reflection of the average daily census, nor does it take into account the number of inpatient or outpatient procedures performed in a hospital or satellite facilities. These rates, for example, may overestimate the risks of sharps injuries in facilities in which large numbers of procedures are performed. For these reasons, these rates are considered experimental and should be interpreted with caution. MDPH welcomes input on the usefulness of these rates, and is exploring alternative rate calculations for future reports.

There are also other limitations to be considered in interpreting the findings presented in this report. In order for an injury to be included on the Annual Sharps Summary, hospitals rely on health care workers to report sharps injuries. As discussed previously, there are many reasons why health care workers may choose not to report sharps injuries, and under-reporting by health care workers has been well documented. Thus the surveillance findings presented in this report should be considered conservative estimates of the burden of sharps injuries among hospital workers in Massachusetts.

Also, there is evidence that the likelihood of reporting varies by occupation and completeness of reporting varies by hospital (CDC, 1999). Hospitals with well established sharps injury surveillance programs and strong safety cultures may identify and report more injuries than hospitals with less

well developed employee health programs. Hospitals, in evaluating their own data, should do so within the context of their own sharps injury surveillance and prevention program.

Assessment of under-reporting should be an integral part of the sharps injury prevention activities in hospitals. Caution is advised in comparing experiences among hospitals. Hospitals with high numbers or rates of reported sharps injuries are not necessarily hospitals with the highest risks of sharps injury, but, rather, may have stronger internal reporting systems. This, however, should not detract from the need to address real problems in these facilities.

For the most part, the information about reported injuries provided by hospitals was complete. However, there was some missing information, and for several data elements (such as department where injury occurred and brand of device) there was some confusion about what information should be submitted. MDPH is working with hospitals to clarify these outstanding issues.

Results

Overview


All 99 hospitals licensed by MDPH submitted Annual Sharps Injury Reports for 2004. A total of 3,279 sharps injuries were sustained and reported by Massachusetts hospital workers from January 1 through December 31, 2004; these injuries were then reported by the hospitals to MDPH. The number of sharps injuries reported by individual hospitals ranged from 0 to 317. Over half of the hospitals reported fewer than 20 injuries. The extent to which high numbers of reported injuries in some hospitals reflect a truly higher incidence of injuries in these hospitals compared to those with low numbers or better sharps injury reporting practices is not known. MDPH plans to work with hospitals over time to better understand injury patterns, and improve reporting and prevention practices.

There are 76 acute care hospitals in Massachusetts. These hospitals reported 97% (3,180) of all sharps injuries. The 18 teaching hospitals in Massachusetts reported 59% (1,923) of all sharps injuries. Teaching status is strongly correlated with hospital size; more than half of the teaching hospitals (12, 67%) have over 300 beds. The 23 non-acute care hospitals are chronic care hospitals.

Key findings for all hospitals combined are presented in the following sections. When the pattern of sharps injuries varied markedly by hospital size, this is noted in the text. Detailed tables, including findings by hospital size categories and teaching status, are provided in Appendices G, H and I.

Comments on data quality are offered to assist hospital staff responsible for compiling the required information for reported injuries. These comments do not address under-reporting of sharps injuries to the surveillance system, which cannot be evaluated without additional sources of information.


Work Status of Injured Worker



State reporting regulations require hospitals to report sharps injuries to all workers in the hospital and satellite sites, regardless of the source of compensation for these workers. Eighty-five percent (2,776) of all sharps injuries reported were sustained by employees, followed by non-employee practitioners, with 9% (310) of the injuries (Figure 1). Three percent (83) of those injured were students. Non-employee practitioners include, but are not limited to, physicians with admitting privileges at a particular hospital and nurse practitioners or physicians assistants from private medical practices who are checking on patients from those practices.

Data quality: Information about work status was provided for 98% of the cases.

Occupation of Injured Worker



Nursing department staff sustained more sharps injuries than any other occupational category, accounting for 39% (1,279) of the injuries (Figure 2). Of these, 9 were among nursing students and 59 were among nursing assistants. Physicians followed nurses with 33% (1,068) of the sharps injuries. Close to half of the injuries in this category (476) were sustained by interns and residents. The physician category also included 66 injuries among medical students. Technicians comprised the third leading occupational group accounting for 20% (657) of sharps injuries. This group included individuals in a wide variety of technical occupations; the most frequently reported were operating room/surgical technicians (252) and phlebotomists (144) and clinical laboratory technicians (69). Of the 123 injuries (4%) sustained by workers in support services, 78 were among housekeepers.


The occupational distribution of the cases varied by hospital size. Most notably, physicians comprised 43% of the injuries in the large hospitals whereas they comprised 22% and 19% in the small and medium sized hospitals respectively (See Appendix H).

Recent studies indicate that the likelihood of workers reporting sharps injuries to employee health departments in hospitals varies by occupation. However, findings are not consistent among studies. In one study, nurses were found to be more likely than physicians to report needle stick injuries (Tandberg, et al., 1991). The CDC found, however, that while nurses were more likely to report needle stick injuries than surgeons, they were less likely to report than other physicians (CDC, 1999). This variation needs to be taken into account in interpreting the findings throughout this report.

Data quality: Information about occupation was provided for almost 100% (3278) of the cases.

Department or Work Area where the Injury Occurred

Figure 3. Sharps Injuries among Hospital Workers by Department where Injury Occurred, Massachusetts, 2004, N=3,279



Data Source: Annual Summary of Sharps Injuries, 2004

The greatest number of reported sharps injuries (1,460; 45%) occurred in operating or procedure rooms (Figure 3); of these, almost three-quarters (1,038) occurred in operating rooms.

In-patient units accounted for the second largest number of cases with 732 (22%) of the injuries. Of these, 570 occurred on medical surgical units, 29 in pediatrics, 24 in Ob/Gyn units, and 20 in psychiatry. For 8 of the injuries that occurred on in-patient units, hospitals reported hospital specific unit identifiers (such as 2 East) that could not be coded to more specific standard locations (See Table 1, page 13 for findings regarding occupation by department).

Data Quality: Some information on location where injuries occurred was provided for almost 100% (3,278) of the cases. However, in a number of cases, hospitals reported hospital specific unit identifiers that could not be coded to standard locations or departments without additional information from the hospitals. MDPH is interested in the department or clinical practice area (physical location) where the injury occurred. Hospitals are encouraged to use the standard department list provided on the Annual Summary of Sharps Injuries reporting form rather than hospital specific nomenclature.


Occupation of Injured Worker by Department where Injury Occurred

Table 1. Sharps Injuries among Hospital Workers by Occupation and Department, Massachusetts, 2004, N=3,279														
Department Where Injury Occurred														
	Operating/ Procedure Room		In-patient Unit		Emergency Department		Intensive Care Unit		Laboratory		Other or Unknown		Total	
	N	%*	N	%*	N	%*	N	%*	N	%*	N	%*	N	%*
Occupation														
Nurse	337	26 %	512	40 %	112	9 %	142	11 %	2	-- %	174	14 %	1,279	100%
Physician	688	64	110	10	89	8	66	6	35	3	80	7	1,068	100%
Technician	360	55	67	10	43	7	27	4	101	15	59	9	657	100%
Support Svcs	28	23	29	24	5	4	11	9	8	7	42	34	123	100%
All others/Unk	47	31	14	9	14	9	2	--	10	7	65	43	152	100%
Total	1,460	45	732	22	263	8	248	8	156	5	420	13	3,279	100%
* Percentages calculated are row percents; percentages for frequencies less than 5 were not calculated														
Data Source: Annual Summary of Sharps Injuries, 2004														

Physicians were most frequently injured in operating and procedure rooms (688, 64%) (Table 1). In contrast, nurses were most frequently injured on in-patient units (512, 40%). Of the 123 support staff who were injured, 78 were housekeepers, of whom 25 were injured on in-patient units.

Within operating and procedure rooms, physicians sustained more injuries than any other occupation group, accounting for 47% (688 of 1,460) of the injuries, followed by nurses with 23% (337 of 1,460) of the injuries. Nurses accounted for by far the greatest number of injuries - 512 of 732 or 70% - in in-patient units. In emergency departments, similar numbers of physicians and nurses were injured. Sixty-five percent (101 of 156) of the injuries in laboratories were sustained by technicians, followed by physicians who accounted for 22% (35 of 156).

Procedure for which Sharp was Used or Intended




Twenty-three percent (754) of the injuries reported involved devices used for injections (Figure 4). Information about type of injection was provided for 444 of these injuries. Of these, 370 involved needles used for subcutaneous injections and 63 for intramuscular injections. In another 22% (729) of the injuries, workers were injured with devices used for suturing. Devices used for blood procedures accounted for 19% (624) of the injuries. The majority of blood procedures (418, 13% of the total) involved devices used for percutaneous venous punctures. Blood procedures are those procedures which involve drawing blood; line procedures involve the insertion or removal of intravenous lines.

The distribution of sharps injuries by procedure varied by hospital size, with 25% of the injuries in large hospitals associated with suturing compared to 19% in both small and medium sized hospitals. In turn, 22% of the injuries in small hospitals and 23% in medium sized hospitals were associated with devices used for blood procedures, compared to 16% in large hospitals. (See Appendix H.)

Data quality: For 7% (228) of the injuries, the procedure for which the device was used or intended was reported as unknown. Most of these cases with unknown procedure (192 of 228) occurred after use of the device, either before, during or after disposal.

Device Involved in the Injury



Injuries from hollow bore needles, particularly those used in procedures accessing a vein or artery and those where residual blood is visible, are associated with increased risk of transmission of HIV when compared to other sharps devices (Cardo, et al., 1997).

As a group, hollow bore needles accounted for the majority – 56% (1,836) - of the sharps injuries reported (Figure 5). These included 1009 (31%) injuries from hypodermic needles, 335 injuries (10%) from butterfly needles, and 143 (4%) from vacuum tube needles. An additional 349 (11%) injuries were associated with “other hollow bore needle”, including IV stylets (152 injuries), epidural needles (21 injuries) and biopsy needles (19 injuries).

Suture needles accounted for 22% (722) of sharps injuries. Information as to whether these were straight or curved needles was provided for only 201 of these injuries; of these, 175 involved curved needles. Consistent with findings for procedures for which devices were used, suture needles accounted for proportionately more injuries in the larger hospitals (25%), as compared to small (19%) and medium (18%) size hospitals.

Data Quality: Information about device type available was not provided for 65 of the injuries. In 57 cases device type was reported as unknown and in 8 cases the question was not answered.

Device Involved in the Injury by Occupation of Injured Worker

Table 2. Sharps Injuries among Hospital Workers by Device and Occupation, Massachusetts, 2004, N=3,279

	Device Type										Total N	%*				
	Hollow Bore				Other Devices											
	Hypodermic Needle	Butterfly Needle	Vacuum Tube	Other Hollow Bore	Suture Needle	Scalpel	All Other/ Unknown									
N	%*	N	%*	N	%*	N	%*	N	%*	N	%*	N	%*			
Occupation																
Nurse	583	46	155	12	66	5	184	14	108	8	43	3	140	11	1,279	100%
Physician	238	22	11	1	7	1	99	9	456	43	116	11	141	13	1,068	100%
Technician	110	17	147	22	59	9	45	7	129	20	53	8	114	17	657	100%
Support Svcs	33	27	6	5	4	--	9	7	10	8	3	--	58	47	123	100%
All others/Unk	45	30	16	11	7	5	12	8	19	13	14	9	39	26	152	100%
Total	1,009	31	335	10	143	4	349	11	722	22	229	7	492	15	3,279	100%

* Percentages calculated are row percents; percentages for frequencies less than 5 were not calculated.
Data Source: Annual Summary of Sharps Injuries, 2004

The type of device involved in the incident varied by occupation (Table 2). Hollow bore needles, as a group, accounted for 77% of injuries sustained by nurses compared to 33% of injuries sustained by physicians. Hypodermic needles accounted for the greatest number of injuries (583; 46%) among nurses, whereas suture needles accounted for the greatest number of injuries (456; 43%) among physicians. The technicians with sharps injuries worked in a wide variety of technical occupations, such as operating room / surgical technicians, phlebotomists, and clinical laboratory technicians. No single device type stood out among the technicians who sustained sharps injuries.

Device Involved in the Injury by Department where Injury Occurred

Table 3. Sharps Injuries among Hospital Workers by Device and Department, Massachusetts, 2004, N=3,279


	Device Type															
	Hollow Bore Needles							Other Devices			Total					
	Hypodermic Needle	Butterfly Needle	Vacuum Tube		Other Hollow Bore	Suture Needle	Scalpel	All Other/Unknown								
N %*	N %*	N %*	N %*	N %*	N %*	N %*	N %*	N %*	N %*							
Department																
OR/Procedure Rm	272	19	50	3	11	1	139	10	597	41	155	11	236	16	1,460	100%
In-patient Units	366	50	118	16	43	6	93	13	28	4	20	3	64	9	732	100%
Emergency Dept	83	32	51	19	23	9	37	14	41	16	8	3	20	8	263	100%
Intensive Care	107	43	37	15	15	6	34	14	27	11	8	3	20	8	248	100%
Outpatient Areas	64	37	23	13	14	8	19	11	6	4	5	3	40	23	171	100%
Laboratories	11	7	39	25	26	17	6	4	4	--	26	17	44	28	156	100%
All Other/Unknown	106	43	17	7	11	4	21	8	19	8	7	3	68	27	249	100%
Total	1,009	31	335	10	143	4	349	11	722	22	229	7	492	15	3,279	100%

* Percentages calculated are row percents; percentages for frequencies less than 5 were not calculated.
Data Source: Annual Summary of Sharps Injuries, 2004

As expected, the type of device associated with sharps injuries varied by department. Within operating and procedure rooms, suture needles accounted for the largest number of injuries reported (597, 41%) followed by hypodermic needles (272, 19%). Suture needles also accounted for a substantial proportion of the injuries in emergency departments (41, 16%) and intensive care units (27, 11%). On in-patient units, hypodermic needles accounted for the greatest number of injuries (366, 50%), followed by butterfly needles (118, 16%) and “other hollow bore needles” (93, 13%). Almost one third of the injuries in laboratory settings involved non-needle devices including scalpels (26, 17%), and glass (20, 13%), which is included in the “all other” category.

Safety Devices

Figure 6. Sharps Injuries among Hospital Workers by Safety Devices, Massachusetts, 2004, N = 3,279



Data Source: Annual Summary of Sharps Injuries, 2004

On the Annual Summary, for each injury, hospitals were encouraged to answer the question “Was it (the device) a safety device?”. For the majority of injuries reported (1,773, 54%), the answer to this question was “No”; the devices involved were not safety devices (Figure 6).


Thirty-three percent (1,072) of injuries were reported to have involved safety devices, underscoring the need to evaluate these devices and to train health care workers in their appropriate use. These data alone cannot be used to assess the efficacy of safety devices, as there is no information regarding the prevalence of safety devices in hospitals. Likewise, these data do not reveal the number of injuries that were prevented by using safety devices. Recent findings from EPINet demonstrate a marked decline in the rate of sharps injuries among nurses in teaching hospitals from 1993 – 2001 (Jagger & Perry, 2003). During this period, there was a substantial increase in the adoption of safety devices. As the number of safety devices increased, there was a rise in the proportion of injuries associated with them, as would be expected. However, the overall injury rate declined.

For some sharps devices, such as suture needles and scalpels, there are only a limited number of alternative devices with engineered sharps injury prevention features available on the market. Excluding injuries involving suture needles and scalpels changed the distribution of safety devices only slightly. The proportion of injuries with standard devices decreased by 12%, while the proportion of injuries with safety devices increased by 12%. This finding highlights the need for increased efforts to meet the federal and state requirements for use of sharps devices with sharps injury prevention features where feasible. Documentation of these situations, as required by MDPH and OSHA, is important to promote effective work-practice controls and the development of new technologies.

The proportion of injuries associated with safety devices was highest in medium hospitals (43%), followed by small sized hospitals (36%) and large hospitals (25%). (See Appendix H) The extent to which this can be explained by the variation in the types of devices used in different sized hospitals is not known.

Standard v. Safety Device by Type of Device Involved in the Injury


Figure 7. Sharps Injuries among Hospital Workers by Device – Standard v. Safety Device, Massachusetts, 2004, N=3,279



Data Source: Annual Summary of Sharps Injuries, 2004

Information as to whether or not the device involved in the injury was a safety device was provided for 2,845 of the 3,279 injuries reported (87%). Among injuries associated with suture needles where safety device information was provided, 621 (100%) occurred with standard devices (Figure 7). Regarding injuries involving scalpel blades, 180 of 195 injuries (92%) occurred with standard devices. Among the 918 injuries from hypodermic needle for which safety device information was reported, 45% involved devices reported as standard devices (416 of 918 injuries). This is the first time the proportion of injuries from standard hypodermic needles is less than the proportion of injuries from hypodermic needles with safety features (Massachusetts Department of Public Health, 2004 and 2006). Among “other hollow-bore needles”, 53% (160 of 302 with information) of injuries involved standard devices. In contrast, 92% (303 of 328 with information) of injuries involving butterfly needles and 72% (93 of 130 with information) of injuries involving vacuum tubes occurred with devices reported as safety devices. It should be noted that safety devices are widely available for many of the device categories shown, such as hypodermic needles, butterfly needles, vacuum tube collection holders and needles, as well as many other types of hollow bore needles. There are, however, a few specific devices currently on the market for which there are no alternative devices with engineered sharps injury prevention features. As previously noted, MDPH and OSHA require employers to maintain documentation of situations where alternative devices are not utilized. This is essential in the promotion of effective work-practice controls and the development of new technologies.

Manufacturer/Brand of Device Involved in the Injury




Information about the manufacturer of the device involved in the injury was provided or was able to be ascertained from the name of the product line in 77% (2,532) of the injuries reported (Figure 8). In 12% (396) of the injuries, the manufacturer of the product was not known, and in 11% (351) there was no response.

Because information about the market share of different manufacturers and product lines was not available, it is not possible to use these data to make judgments about a particular manufacturer's products and the efficacy of the products with respect to safety.

Data quality: Both OSHA and MDPH regulations pertaining to sharps injuries require facilities to collect and record information about the "brand" of the devices involved in the incidents. There is some legitimate confusion about whether "brand" means the name of the manufacturer or name of the product line. Technically "brand" means name of the product line. This distinction was not made clear in previous instructions to hospitals. MDPH is interested in the name of the product line as well as the manufacturer, and will clarify this on forms for the future.

When the Injury Occurred: Before, During, After Use of Device

Figure 9. Sharps Injuries among Hospital Workers by When the Injury Occurred, Massachusetts, 2004, N=3,279




Data Source: Annual Summary of Sharps Injuries, 2004

Injuries occurred at various points in the course of handling needles or other sharp devices (Figure 9). After use was a dangerous time: about half of the injuries (1,603, 49%) occurred either after use and before disposal (1,105, 34%) or during or after disposal (498, 15%) of the device. Forty-two percent (1,363) occurred during use of the item. The 53 injuries (2%) that happened before use of the item involved sharps devices penetrating contaminated gloves.

How the Injury Occurred

Figure 10. Sharps Injuries among Hospital Workers by How the Injury Occurred, Massachusetts, 2004, N = 3,279



Data Source: Annual Summary of Sharps Injuries, 2004

The largest number of injuries reported (751, 23%) fell into the broadly defined category of “collision with sharp or other person”. Another 12% of injuries (387) occurred while suturing (Figure 10).


Improper disposal of sharps accounted for 8% (250) of the injuries. These included cases in which the contaminated sharps were left on the floor, in the trash, or in beds. Others cases involved sharps found in linens or laundry, in clothing, or on tables or trays.

Eight percent (247) of the injuries occurred during disposal. A majority of these (133, 4%) were reported as involving sharps containers. In 2% (71) of the injuries, the health care worker was injured by the sharp being disposed of while placing it in the sharps container. In 13 cases (<1%), the health care worker was injured by a sharp already in the container.

Five percent of the injuries (156) occurred during activation of safety devices. No information was collected regarding the failure rate of safety features; therefore it is not possible to use this information to assess the efficacy of the safety features on those devices.


It should be noted that the OSHA Bloodborne Pathogen standard states that contaminated needles shall not be recapped. In 3% (100) of the cases, injuries occurred while recapping devices.

Experimental Sharps Injury Rates by Number of Licensed Hospital Beds



The statewide rate of sharps injuries among hospital workers for this twelve month surveillance period was 18.3 sharps injuries per 100 licensed hospital beds. The annual rate of sharps injuries varied by hospital size (Figure 11). Large hospitals had the highest annual rate of 24.9 sharps injuries per 100 licensed hospital beds, followed by small and medium sized hospitals, which had annual sharps injury rates of approximately 15 sharps injuries and 13 sharps injuries per 100 licensed hospital beds,

respectively. As discussed on page 7, given the limitations of hospital bed size as a denominator for assessing risks, these rates should be interpreted with caution. In comparing experience among hospitals, under-reporting must be taken into account. The extent to which high rates of reported injuries in some hospitals reflect a true higher incidence of injuries in these hospitals or better sharps injury reporting practices compared to those with low rates is not known. Comparison of rates among facilities is of limited usefulness (CDC, 2004; Perry, et. al., 2003). Hospitals evaluating their own rates should do so within the context of their own sharps injury surveillance and prevention programs.



Injuries reported by acute care hospitals accounted for 97% of all injuries reported. However, acute care hospitals account for only 79% of all licensed hospital beds. Therefore, sharps injury rates which include all licensed hospitals underestimate the risk for acute care hospitals. Sharps injury rates for acute care hospitals are presented below in order to more accurately reflect the injury rates in those settings.

Discussion

Sharps injuries are preventable and the overall goal should be their elimination. As a step in that direction, the U.S. Public Health Service has called for the reduction of sharps injuries among health care workers by 30% as a national health objective for 2010 (DHHS, 2000). Preventing sharps injuries requires the combined efforts of government agencies, employers, and equipment manufacturers, as well as health care workers. The Massachusetts Sharps Injury Surveillance System is intended to provide information to guide and evaluate these efforts in Massachusetts.

Nearly 3,300 sharps injuries were reported by Massachusetts hospitals in 2004, underscoring the need for continued efforts to reduce the incidence of these injuries. Given previously documented underreporting of sharps injuries to employee health by health care workers, this figure likely underestimates the full extent of the problem. While overall patterns are similar to NaSH and EpiNET, findings highlight a number of specific issues to be addressed in Massachusetts:

- Almost 50% of the injuries reported occurred after use of devices, including 23% which occurred either during clean up or disposal or as a result of improper disposal. Elimination of these preventable injuries will have a large impact on the incidence of sharps injuries in hospitals. (These injuries are entirely preventable.) Prevention strategies include the purchase and appropriate placement of sharps containers that allow staff to determine when containers should be emptied before they are dangerously full. It is also crucial to implement systems to regularly check containers to identify those that need to be replaced. Increased training and supervision to avoid improper disposal is needed in addition to appropriate sharps containers, and is essential to protect health care providers, support service workers and patients.
- Close to a third of the injuries reported were associated with hypodermic needles, and of these, more than 41% involved devices without sharps injury prevention features. There are a wide variety of hypodermic needles with engineered sharps injury prevention features on the market. Hospitals should evaluate their device inventory and aggressively identify, evaluate and implement use of alternative devices with engineered sharps injury prevention features.
- Blood procedures continue to account for about 20% of all injuries reported. This is may be related to the device used, particularly whether or not the device was one with engineered sharps injury prevention features. As previously shown, almost a third of all injuries with vacuum tube collection holders and needles occurred with standard devices. There are a wide variety of products with safety features available for these types of devices. As mentioned earlier, injuries with hollow bore needles, particularly those used for blood procedures, are associated with a higher risk of transmission of bloodborne pathogens.
- It may be more difficult to institute change in some areas than others: the operating room setting, for example, poses unique challenges. Some devices, such as suture needles, have fewer options for engineering controls; to date, safer options for suture needles have been blunt needles, which are not appropriate for all situations. In this instance, exploring alternative methods of closing wounds may be more appropriate than finding alternative devices. The use of neutral zones in the operating room to minimize hand-to-hand transfer of sharps is an effective work practice control to reduce sharps injuries.

The Massachusetts Sharps Injury Surveillance System is a collaborative effort between the MDPH, hospitals, professional associations and community advocates. The success of the program in collecting data is a result of this collaboration. MDPH will continue to work with these groups to conduct surveillance, review exposure control activities in hospitals, and facilitate the exchange of information among hospitals about successful prevention strategies.

References

- Cardo, D.M., Culver, D.H., Ciesielski, C.A., Srivastava, P.U., Marcus, R., Abiteboul, D., et al. (1997). A case-control study of HIV seroconversion in health care workers after percutaneous exposure. Centers for Disease Control and Prevention Needlestick Surveillance Group. *New England Journal of Medicine*, 337(21), 1485-90.
- The Centers for Disease Control and Prevention National Center for Infectious Diseases Division of Health Quality Promotion. (2004) Workbook for Designing, Implementing, and Evaluating a Sharps Injury Prevention Program. Retrieved November 1, 2006, <http://www.cdc.gov/sharpssafety>.
- The Centers for Disease Control and Prevention Hospital Infections Program. (1999). The National Surveillance System for Hospital Health Care Workers Summary Report for Data Collected from June 1995 through July 1999. Retrieved November 1, 2006, <http://www.cdc.gov/ncidod/hip/NASH/report99.PDF>.
- The Centers for Disease Control and Prevention. (2001). Updated U.S. Public Health Service Guidelines for the Management of Occupational Exposures to HBV, HCV, and HIV and Recommendations for Postexposure Prophylaxis. *Morbidity and Mortality Weekly Report*, 50, (No. RR-11).
- The Centers for Disease Control and Prevention. (2005). Updated U.S. Public Health Service Guidelines for the Management of Occupational Exposures to HIV and Recommendations for Postexposure Prophylaxis. *Morbidity and Mortality Weekly Report*, 54, (No. RR-9).
- The Centers for Disease Control and Prevention. (1982). Recommendation of the Immunization Practices Advisory Committee (ACIP) Inactivated Hepatitis B Virus Vaccine. *Morbidity and Mortality Weekly Report*, 31(24),317-22, 327-8.
- U.S. Department of Health and Human Services. (2000). *Healthy People 2010*. 2nd ed. With Understanding and Improving Health and Objectives for Improving Health. 2 vols. Washington, DC: U.S. Government Printing Office.
- Clarke, S.P., Rockett, J.L., Sloane, D.M., Aiken, L.H. (2002). Organizational climate, staffing, and safety equipment as predictors of needlestick injuries and near-misses in hospital nurses. *American Journal of Infection Control*, 30:207-16.
- Exposure Prevention Information Network (EPINet). (1998). Annual Number of Occupational Percutaneous Injuries and Mucocutaneous Exposures to Blood or Potentially Infective Biological Substances. International Health Care Worker Safety Center, University of Virginia. Retrieved April 9, 2002, <http://www.med.virginia.edu/medcntr/centers/epinet/estimates.html>.
- Jagger, J., Bentley, M., Juliet, E. (1998). Direct Cost of Follow-up for Percutaneous and Mucocutaneous Exposures to At-Risk Body Fluids: Data from Two Hospitals. *Advances in Exposure Prevention*, 3(3).
- Jagger, J. and Perry, J. (2003). Marked Decline in Needlestick Injury Rates. *Advances in Exposure Prevention*, 6(3).

- Massachusetts Department of Workforce Development. (2005). Regional LMI Profile Series Massachusetts Industry Staffing Patterns: Volume 5 of 5: Education and Health Services, Leisure and Hospitality, Other Services, and Government. Retrieved November 14, 2006 from <http://lmi2.detma.org/lmi/pdf/naicVol5.pdf>.
- Massachusetts Department of Workforce Development. (2004). Employment and Wages (ES-202). Retrieved November 14, 2006, <http://lmi2.detma.org>.
- National Institute for Occupational Safety and Health (NIOSH). (1999). *NIOSH Alert: Preventing Needlestick Injuries in Health Care Settings, DHHS (NIOSH) Publication No. 2000-18*. Washington, DC: U.S. Government Printing Office.
- Occupational Safety and Health Administration. (1998). Occupational Exposure to Bloodborne Pathogens: Request for information. *Federal Register*, 63,48250-48252.
- Panlilio AL, Cardo DM, Campbell S, Srivastava PU, Jagger H, Orelie JG et al. Estimate of the annual number of percutaneous injuries in U.S. health care workers [Abstract S-T2-01]. In: Program and abstracts of the 4th International Conference on Nosocomial and Health Care-Associated Infections; Atlanta, March 5-9, 2000:61.
- Perry, J. (2000). CDC Releases National Needlestick Estimates. *Advances in Exposure Prevention*, 5(2),19.
- Perry, J., Parker, G., Jagger, J. (2003). 2001 Percutaneous Injury Rates. *Advances in Exposure Prevention*, 6(3),32-36.
- Tandberg, D., Stewart, K.K., Doezema, D. (1991). Under-reporting of contaminated needlestick injuries in emergency health care workers. *Annals of emergency Medicine*, 20(1),66-70.

APPENDIX A

Massachusetts Department of Public Health Sharps Injury Prevention Advisory Committee

Members

Philip Adamo M.D., MPH	Board of Directors, New England College of Occupational and Environmental Medicine
Bob Alconada	Manager, State Government Affairs Massachusetts Dental Society
Evelyn Bain, M Ed, RN, COHN-S	Associate Director/Coordinator - Health and Safety Program Massachusetts Nurses Association
Karen Daley, RN, MPH	Consumer
Anuj Goel, JD, MPH	Director, Regulatory Compliance Massachusetts Hospital Association
Margaret Quinn, Sc.D., CIH	Associate Professor, Department of Work Environment, University of Massachusetts Lowell
James Ryan, M.D., MPH	Committee on, Occupational & Environmental Medicine Massachusetts Medical Society

Staff

Letitia Davis, Sc.D.	Director Occupational Health Surveillance Program, MDPH
Alfred DeMaria, Jr., M.D.	Assistant Commissioner, Bureau of Communicable Disease Control State Laboratory Institute, MDPH
Natalia Firsova, MA	Research Analyst, Occupational Health Surveillance Program, MDPH
Angela Laramie, MPH	Project Coordinator, Sharps Injury Surveillance Program Occupational Health Surveillance Program, MDPH
Marie-Eileen Onieal, Ph.D.(c), MMHS, RNC, PNP	Health Policy Coordinator Bureau of Health Quality Management, MDPH
Gail Palmeri, RN	Program Manager, Hospitals Division of Health Care Quality, MDPH

APPENDIX B

NOTE: This is an unofficial copy.

Chapter 252 of the Acts of 2000

AN ACT RELATIVE TO NEEDLESTICK INJURY PREVENTION.

Be it enacted by the Senate and House of Representatives in General Court assembled, and by the authority of the same, as follows:

SECTION 1. Chapter 111 of the General Laws is hereby amended by inserting after section 53C the following section:

Section 53D. (a) Any acute or non-acute hospital licensed under this chapter shall ensure the provision of services to individuals through the use of hollow-bore needle devices or other technology that minimize the risk of injury to health care workers from hypodermic syringes or needles, in accordance with rules and regulations promulgated pursuant to subsection (b).

(b) The department shall promulgate rules and regulations requiring the use, at all acute and non-acute hospitals, of only such devices which minimize the risk of injury to health care workers from needlestick and sharps, so-called. Such rules and regulations promulgated by the department shall include the following requirements:

- (1) Written exposure control plans shall be developed by each acute and non-acute hospital that include an effective procedure for identifying and selecting existing sharps prevention technology, so-called, of the types specified by the department.
- (2) Sharps injury prevention technology shall be included as engineering or work practice controls, except in cases where the employer or other appropriate party can demonstrate circumstances in which the technology does not promote employee or patient safety or interferes with a medical procedure. Those circumstances shall be specified by the employer and shall include, but not be limited to, circumstances where the technology is medically contraindicated or not more effective than alternative measures used by the employer to prevent exposure incidents. In all cases the department shall make the final determination as to whether an employer or other appropriate party has demonstrated in a satisfactory manner circumstances which warrant an exemption from the inclusion of sharps injury prevention technology.
- (3) Information concerning exposure incidents shall be recorded in a sharps injury log to be kept within such acute and non-acute hospitals and reported annually to the department, including but not limited to, the type and brand of device involved in the incident. Such logs shall be used as the basis for continuing quality improvement in reducing sharps injuries through the provision of education and the procurement of improved products. Such logs shall be kept confidential and shall be used only for the intended purposes of this section.
- (4) Written exposure control plans shall be updated when necessary to reflect progress in sharps prevention technology as determined by the department.

(c) The department shall promulgate all rules and regulations pursuant to this section in consultation with an advisory committee composed of, but not limited to: the department's director of infectious disease, a consumer to be selected by the commissioner, a technical expert to be selected by the commissioner, and a representative from the Massachusetts Nurses Association, the Massachusetts Association of Occupational and Environmental Medicine, the Massachusetts Medical Society and the Massachusetts Hospital Association.

APPENDIX B

The department, in consultation with the advisory committee, shall compile and maintain a list of needless systems, needles and sharps, so-called, with engineered injury protections meeting the purposes of this section. The list shall be available to assist employers in complying with rules and regulations promulgated in accordance with this section.

SECTION 2. The department of public health shall promulgate the rules and regulations required by section 53D of chapter 111 of the General Laws no later than November 1, 2000.

Approved August 17, 2000.

APPENDIX C

NOTE: This is an unofficial copy.

105 CMR 130.000 Hospital Licensure Regulations

105 CMR 130.000 is amended by adding the following new sections:

130.1001: Definitions

As used in 105 CMR 130.1001 through 130.1008 the following definitions shall apply:

“Advisory committee,” means a committee composed of, but not limited to the Department’s director of infectious disease; a consumer to be selected by the commissioner; a technical expert to be selected by the commissioner; and a representative from the Massachusetts Nurses Association, the New England Association of Occupational and Environmental Medicine, the Massachusetts Medical Society and the Massachusetts Hospital Association.

“Commissioner” means the Commissioner of the Massachusetts Department of Public Health.

“Department” means the Massachusetts Department of Public Health.

“Engineering and work practice controls” mean controls such as, but not limited to, sharps disposal containers, needleless systems, and sharps with engineered injury protections, that isolate or remove the bloodborne pathogens hazard from the workplace.

“Exposure Control Plan” means a plan that includes an effective procedure for identifying and selecting existing sharps injury prevention technology.

“Exposure Incident” means a specific eye, mouth, other mucous membrane, non-intact skin, or parenteral contact with blood or other potentially infectious materials that result from the performance of an employee’s duties.

“Health care worker” means all workers employed by the hospital, working within the hospital but employed by other agencies, those providing patient care services without pay such as students, or providers who are delivering care but receiving compensation from sources other than the hospital.

“Hospital” means any hospital licensed by the Department pursuant to M.G.L. c.111, § 51.

“Reportable Exposure incident” means an exposure incident a result of events that pierce the skin or mucus membranes.

“Sharp” means any object that can penetrate the skin or any part of the body, and result in an exposure incident, including, but not limited to, needle devices, scalpels, lancets, broken glass, broken capillary tubes and exposed ends of dental wires.

“Sharps injury log” means a log to be kept within acute and non-acute hospitals that records information concerning exposure incidents, including but not limited to, the type and brand of device involved in the incident.

“Sharps injury prevention technology” means devices or other technology that minimizes the risk of injury to health care workers from hypodermic syringes, needles or other sharps.

130.1002: Minimizing Risk of Injury

APPENDIX C

Every hospital shall:

- (A) Ensure the provision of services to individuals through the use of safe needle devices or other technology that minimizes the risk of injury to health care workers from hypodermic syringes, needles, and sharps; and
- (B) Except as provided in 105 CMR 130.1005; use only such devices designed to reduce risk of percutaneous exposure to bloodborne pathogens.

130.1003: Written Exposure Control Plans

Hospitals shall develop written exposure control plans that include an effective procedure for identifying and selecting existing sharps injury prevention technology consistent with the federal regulations concerning occupational exposure to bloodborne pathogens, 29 CFR 1910.1030 *et seq.* the Occupational Safety & Health Administration's (OSHA) Occupational Exposure to Bloodborne Pathogens standards. Written exposure control plans shall be updated when necessary to reflect progress in sharps injury prevention technology as determined by the Department.

130.1004: Engineering and Work Practice Controls

Hospitals shall include sharps injury prevention technology as engineering and work practice controls to isolate or remove the bloodborne pathogens hazard from the workplace consistent with the federal regulations concerning occupational exposure to bloodborne pathogens, 29 CFR 1910.1030 *et seq.*

130.1005: Exemption from the Inclusion of Sharps Injury Prevention Technology

- (A) Sharps injury prevention technology may be excluded as engineering and work practice controls in cases where the hospital or other appropriate party can demonstrate circumstances in which the technology does not promote employee or patient safety or interferes with a medical procedure.
- (B) Where sharps injury prevention technology is not utilized, the hospital shall specify those circumstances, which shall include but not be limited to, situations where the technology is medically contraindicated or not more effective than alternative measures used by the hospital to prevent exposure incidents.
- (C) In all cases the Department shall make the final determination as to whether a hospital or other appropriate party has demonstrated in a satisfactory manner those circumstances which warrant an exemption from the inclusion of sharps injury prevention technology.

130.1006: Sharps Injury Log

- (A) Information concerning exposure incidents shall be recorded in a sharps injury log that includes, but is not limited to, the type and brand of device involved in the incident, the department or work area where the exposure incident occurred, and an explanation of how the incident occurred.
- (B) Sharps injury logs shall be kept within the hospital and shall be used as the basis for continuing quality improvement in reducing sharps injuries through the provision of education and the procurement of improved products; and,

APPENDIX C

(C) Sharps injury logs shall be kept confidential.

130.1007: Reporting

Every licensed acute and non-acute care hospital shall report annually to the Department information from its sharps injury logs and such other information as the Department may require concerning exposure incidents. The Department shall supply each reporting hospital with guidelines indicating the specific data elements to be submitted.

130.1008: Advisory Committee

The Department shall convene an advisory committee composed of, but not limited to the Department's director of infectious disease; a consumer to be selected by the commissioner; a technical expert to be selected by the commissioner; and a representative from the Massachusetts Nurses Association, the New England Association of Occupational and Environmental Medicine, the Massachusetts Medical Society and the Massachusetts Hospital Association.

130.1009: List of Needleless Systems

The Department, in consultation with the advisory committee, shall compile, maintain and periodically update a list of needleless systems, with engineered injury protections meeting the purposes set forth in M.G.L. c. 111, § 53D. The list shall be available as a resource to assist hospitals in complying with these regulations.

APPENDIX D

MDPH Data Elements to be Recorded for each Exposure Incident

Those items in bold are required to be recorded by both OSHA and MDPH. The additional items are strongly recommended by MDPH to be recorded. The checks in the left-hand column identify the subset of data elements that should be reported annually to MDPH for each exposure incident. See also Annual Summary of Sharps Injuries. The items in bold are those data elements which employers are required to collect as outlined in the OSHA Bloodborne Pathogen Standard (29 CFR 1910.1030) and the OSHA Recordkeeping Standard (29 CFR 1904).

To be reported to MDPH annually	Data elements
√	Employer
√	Unique Incident Number
√	Employment status of exposed health care worker (temp, agency employee, pool nurse, contractor, employee)
√	Date of incident
	Time of incident
	Time work shift began
√	Occupation
√	Department or work area in which the exposure incident occurred
√	Device or item that was involved in the injury
√	Brand and model of device
√	Was the device a safety device?
√	Purpose or procedure for which the sharp was intended or used
√	How the incident occurred
	Health care worker's recommendations to prevent similar injuries

APPENDIX E

This form meets the requirements of recording sharps injuries under M.G.L. 105 CMR 130.1001 *et seq.*
 Please complete this form with the exposed health care worker. ***REQUIRED DATA ELEMENTS FOR RECORDING**

Massachusetts Department of Public Health Bloodborne Pathogen Exposure Incident Recording Form			
EMPLOYER:*		UNIQUE EXPOSURE INCIDENT NUMBER:*	
EXPOSED WORKER'S NAME: (or unique ID number)		OSHA RECORDABLE: <input type="checkbox"/> YES <input type="checkbox"/> NO <input type="checkbox"/> UNKOWN	
STATUS of EXPOSED WORKER: <input type="checkbox"/> EMPLOYEE <input type="checkbox"/> VOLUNTEER <input type="checkbox"/> STUDENT <input type="checkbox"/> NON EMPLOYEE PRACTITIONER <input type="checkbox"/> TEMP / CONTRACT <input type="checkbox"/> OTHER _____		TIME WORK am SHIFT BEGAN:* : pm	
DATE OF INCIDENT:* / /	TIME of INCIDENT:* : am pm	DATE REPORTED: / /	TIME REPORTED: : am pm
TYPE OF EXPOSURE:* <input type="checkbox"/> Percutaneous <input type="checkbox"/> Mucous membrane <input type="checkbox"/> Skin <input type="checkbox"/> Was skin intact?: YES NO UNKNOWN <input type="checkbox"/> Bite	TYPE OF FLUID: <input type="checkbox"/> Blood / blood products <input type="checkbox"/> Visibly bloody body fluid <input type="checkbox"/> Non-visibly bloody body fluid <input type="checkbox"/> Visibly bloody solution (iv fluid, etc.) <input type="checkbox"/> Non-visibly bloody solution <input type="checkbox"/> Other _____ (specify) <input type="checkbox"/> Unknown	FOR PERCUTANEOUS INJURIES: DEPTH OF INJURY: <input type="checkbox"/> Superficial <input type="checkbox"/> Moderate <input type="checkbox"/> Deep <input type="checkbox"/> Unknown BLOOD VISIBLE ON DEVICE BEFORE EXPOSURE? <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Unknown	
BODY PART INJURED: <input type="checkbox"/> Arm <input type="checkbox"/> Mouth / nose <input type="checkbox"/> Hand <input type="checkbox"/> Leg <input type="checkbox"/> Finger <input type="checkbox"/> Other _____ (specify)		PERSONAL PROTECTIVE EQUIPMENT WORN BY WORKER AT TIME OF EXPOSURE: <input type="checkbox"/> Gloves (single pair) <input type="checkbox"/> Eye protection <input type="checkbox"/> Mask <input type="checkbox"/> Gloves (double pair) <input type="checkbox"/> Face shield <input type="checkbox"/> Other _____ (specify) <input type="checkbox"/> Gloves (triple pair) <input type="checkbox"/> Gown/Garment <input type="checkbox"/> None of the above	
OCCUPATION:* <input type="checkbox"/> Attendant / orderly <input type="checkbox"/> Fellow <input type="checkbox"/> Medical student <input type="checkbox"/> Physical therapist <input type="checkbox"/> Attending physician <input type="checkbox"/> Fireperson / First responder <input type="checkbox"/> Nurse Anesthetist <input type="checkbox"/> Public health worker <input type="checkbox"/> Central supply <input type="checkbox"/> Food service <input type="checkbox"/> Nursing Assistant <input type="checkbox"/> Psychiatric technician <input type="checkbox"/> Clerical / administrative <input type="checkbox"/> Hemodialysis technician <input type="checkbox"/> Nurse Midwife <input type="checkbox"/> Radiologic technician <input type="checkbox"/> Clinical lab technician <input type="checkbox"/> Housekeeper <input type="checkbox"/> Nurse Practitioner <input type="checkbox"/> Registered Nurse <input type="checkbox"/> Counselor / social worker <input type="checkbox"/> Intern / resident <input type="checkbox"/> Nursing student <input type="checkbox"/> Researcher <input type="checkbox"/> Dentist <input type="checkbox"/> Laundry staff <input type="checkbox"/> OR / surgical technician <input type="checkbox"/> Respiratory Therapist / Tech <input type="checkbox"/> Dental assistant / tech <input type="checkbox"/> Law enforcement officer <input type="checkbox"/> Patient care technician <input type="checkbox"/> Safety / security <input type="checkbox"/> Dental hygienist <input type="checkbox"/> Licensed Practical Nurse <input type="checkbox"/> Pharmacist <input type="checkbox"/> Transport / messenger <input type="checkbox"/> Dental student <input type="checkbox"/> Maintenance <input type="checkbox"/> Phlebotomist <input type="checkbox"/> Volunteer <input type="checkbox"/> Dietician <input type="checkbox"/> Morgue technician <input type="checkbox"/> Physician assistant <input type="checkbox"/> Other _____ (specify)			
DEPARTMENT OR WORK AREA WHERE EXPOSURE INCIDENT OCCURRED:* <i>Select all that apply</i> Identify specific location (room number, floor etc): _____ <input type="checkbox"/> Ambulance <input type="checkbox"/> Endoscopy / <input type="checkbox"/> Blood bank bronchoscopy /cytосcopy <input type="checkbox"/> Intensive care unit <input type="checkbox"/> Obstetrics / gynecology ward <input type="checkbox"/> Central sterile supply <input type="checkbox"/> Exam room <input type="checkbox"/> Jail unit <input type="checkbox"/> Operating room <input type="checkbox"/> Central trash area <input type="checkbox"/> Hematology <input type="checkbox"/> Labor and delivery <input type="checkbox"/> Pediatrics <input type="checkbox"/> Clinical chemistry <input type="checkbox"/> Histology / pathology <input type="checkbox"/> Laundry room <input type="checkbox"/> Procedure room <input type="checkbox"/> Dialysis <input type="checkbox"/> Home health visit (home) <input type="checkbox"/> Medical / surgical ward <input type="checkbox"/> Psychiatry ward <input type="checkbox"/> Dental Clinic <input type="checkbox"/> Hospital grounds <input type="checkbox"/> Microbiology <input type="checkbox"/> Radiology department room <input type="checkbox"/> Emergency Department <input type="checkbox"/> Morgue / autopsy room <input type="checkbox"/> Other location _____ (specify)			
IS THIS THE DEPARTMENT TO WHICH THE WORKER IS REGULARLY ASSIGNED? <input type="checkbox"/> YES <input type="checkbox"/> NO <input type="checkbox"/> N/A			
IF NO, TO WHICH DEPARTMENT IS THE WORKER REGULARLY ASSIGNED?			

APPENDIX E

WHAT DEVICE OR ITEM WAS INVOLVED IN THE INJURY?*		
<p>Hollow bore needle</p> <input type="checkbox"/> Biopsy needle <input type="checkbox"/> IV stylet <input type="checkbox"/> Hollow-bore needle, type unknown <input type="checkbox"/> Huber needle <input type="checkbox"/> Hypodermic needle attached to a disposable syringe <input type="checkbox"/> Hypodermic needle attached to IV tubing <input type="checkbox"/> Prefilled cartridge syringe <input type="checkbox"/> Spinal or epidural needle <input type="checkbox"/> Unattached hypodermic needle <input type="checkbox"/> Winged steel needle <input type="checkbox"/> Winged steel needle attached to a vacuum tube collection holder <input type="checkbox"/> Winged steel needle attached to IV tubing <input type="checkbox"/> Vacuum tube collection holder / needle <input type="checkbox"/> Other type of hollow bore needle _____ (Specify)	<p>Other sharp object</p> <input type="checkbox"/> Bone chip / chipped tooth <input type="checkbox"/> Bone cutter <input type="checkbox"/> Bovie electrocautery device <input type="checkbox"/> Bur <input type="checkbox"/> Explorer <input type="checkbox"/> Histology cutting blade <input type="checkbox"/> Lancet <input type="checkbox"/> Laser <input type="checkbox"/> Pin <input type="checkbox"/> Razor <input type="checkbox"/> Retractor <input type="checkbox"/> Scaler / curette <input type="checkbox"/> Scalpel blade <input type="checkbox"/> Scissors <input type="checkbox"/> Sharp object, type unknown <input type="checkbox"/> Tenaculum <input type="checkbox"/> Trocar <input type="checkbox"/> Wire <input type="checkbox"/> Other type of sharp object _____ (specify)	<p>Suture needle</p> <input type="checkbox"/> Curved suture needle <input type="checkbox"/> Straight suture needle
<p>Glass</p> <input type="checkbox"/> Capillary tube <input type="checkbox"/> Medication ampule / vial / IV bottle <input type="checkbox"/> Pipette <input type="checkbox"/> Slide <input type="checkbox"/> Specimen / test / vacuum tube <input type="checkbox"/> Other glass item _____ (specify) <p>Additional dental / surgical devices</p> <input type="checkbox"/> Hypodermic needle attached to non-disposable syringe <input type="checkbox"/> Elevator <input type="checkbox"/> Extraction forceps <input type="checkbox"/> Root canal file <input type="checkbox"/> Rod (orthopaedic) <input type="checkbox"/> Other device or item _____ (specify)		
BRAND / MODEL OF DEVICE:*		
WAS IT A SAFETY DEVICE? <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Unknown		
IF YES, WHEN DID THE INJURY OCCUR?		
<input type="checkbox"/> Before activation of safety feature <input type="checkbox"/> Safety feature failed; after activation <input type="checkbox"/> Other _____ (specify) <input type="checkbox"/> During activation of safety feature <input type="checkbox"/> Safety feature not activated <input type="checkbox"/> Safety feature improperly activated <input type="checkbox"/> Passive safety feature, activation not required <input type="checkbox"/> Unknown		
IF YES, WAS THE WORKER TRAINED IN THE PROPER USE OF THIS SAFETY DEVICE? <input type="checkbox"/> Yes <input type="checkbox"/> No		Describe training:
PURPOSE OR PROCEDURE FOR WHICH SHARP WAS USED OR INTENDED:*		
<p>Line procedures:</p> <input type="checkbox"/> To insert a peripheral IV line or set up a heparin lock <input type="checkbox"/> To insert a central IV line <input type="checkbox"/> To insert an arterial line <input type="checkbox"/> To connect IV line (intermittent IV / piggy back / IV infusion / other IV line connection) <input type="checkbox"/> To flush heparin / saline <input type="checkbox"/> Other injection into IV injection site or IV port _____ (specify) <input type="checkbox"/> Other line procedure _____ (specify) <p>Blood procedures:</p> <input type="checkbox"/> Percutaneous venous puncture (e.g. phlebotomy) <input type="checkbox"/> Percutaneous arterial puncture <input type="checkbox"/> Central or peripheral IV line or port <input type="checkbox"/> Arterial line <input type="checkbox"/> Dialysis / AV fistula site <input type="checkbox"/> Umbilical vessel <input type="checkbox"/> Fingerstick / heel stick <input type="checkbox"/> Other blood sampling _____ (specify)	<p>Other procedures:</p> <input type="checkbox"/> Cutting (e.g. surgery / autopsy) <input type="checkbox"/> During disposal <input type="checkbox"/> Epidural / spinal anesthesia <input type="checkbox"/> Intramuscular (IM) injection <input type="checkbox"/> Subcutaneous / intradermal injection / skin test placement <input type="checkbox"/> Suturing <input type="checkbox"/> Transferring blood / body fluid to another container <input type="checkbox"/> To obtain a body fluid or tissue sample (CFS / amniotic / biopsy) <input type="checkbox"/> To obtain laboratory specimens <input type="checkbox"/> Other procedure (not a line procedure or blood sampling procedure) _____ (specify) <input type="checkbox"/> Unknown	<p>Dental procedure:</p> <input type="checkbox"/> During disposal <input type="checkbox"/> Hygiene (prophy, root plane, curettage) <p>Oral surgery</p> <input type="checkbox"/> Simple Extraction <input type="checkbox"/> Surgical Extraction <input type="checkbox"/> Fracture Reduction <input type="checkbox"/> Other _____ (specify) <input type="checkbox"/> Unknown <p>Orthodontic procedure</p> <input type="checkbox"/> Periodontal surgery <input type="checkbox"/> Restorative (amalgam, composite, crown) <input type="checkbox"/> Root canal <input type="checkbox"/> Other _____ (specify) <input type="checkbox"/> Unknown <p>Where did the injury occur?</p> <input type="checkbox"/> Inside the patient's mouth <input type="checkbox"/> Outside the patient's mouth <input type="checkbox"/> Unknown

APPENDIX E

INSTRUCTIONS FOR MDPH BLOODBORNE PATHOGEN EXPOSURE INCIDENT RECORDING FORM

The Bloodborne Pathogen Exposure Incident Recording Form shall be completed with the exposed health care worker at the time that post-exposure care is given following a percutaneous injury resulting in an exposure to blood and potentially infectious bodily fluids.

Health care workers are defined as: all workers employed in the hospital, working within the hospital but employed by other agencies, those providing patient care services without pay such as students, or providers who are delivering care but receiving compensation from sources other than the hospital.

This form shall be kept in a place that protects the confidentiality of the exposed health care worker. If this information is to be shared with committees within the hospital, all measures that protect the privacy of the exposed health care worker shall be taken.

The **name of the employer** shall be recorded. If incident occurred in a satellite site, note site here.

A **unique exposure incident number** shall be assigned to each incident. This number along with the ID number should be used when referring to this incident on subsequent reports. There should be only one location where the connection is made between the ID number, incident number and the health care worker's name. This information shall be kept confidential.

The **exposed health care worker's name** or **unique ID number** shall be recorded. An **ID number**, unique to the exposed health care worker should be assigned. A social security number or employee ID number should not be used. If this form is shared with other departments, then the health care worker's name should not be used, in order to maintain confidentiality.

Indicate if this is an **OSHA recordable** incident.

The **employment status** shall be given. If the health care worker is a paid employee of the organization, then indicate that the health care worker is an employee. If the health care worker is from an outside agency, (e.g., staffing agency) then indicate that the health care worker is a temp or a contract employee. An attending physician employed by a group practice would be classified as a non-employee practitioner.

Indicate the time that the health care worker began the **work shift** in which the incident occurred.

Indicate the **date and time of the incident**, and the **date and time that the incident was reported**.

APPENDIX E

Indicate the **type of exposure**.

Percutaneous – punctured or broke the skin

Mucous membrane – contact with mouth, eyes or other mucous membranes

Skin – contact with unprotected skin

Bite – bite where the skin was broken

Identify the **type of fluid** involved in the exposure. If the fluid type is not listed, describe in OTHER.

Describe the **depth of the injury**.

Superficial – injuries such as a scratch

Moderate – those injuries that are more serious than scratches, but not so serious that they would be considered to be deep (e.g., superficial laceration or tissue avulsion)

Deep – injuries that touched bone or muscle contracted

Indicate whether there was **blood visible on the device** before the incident occurred.

Indicate the **body part injured**. If it is not listed, describe in OTHER.

Indicate the type of **personal protective equipment** worn by the exposed health care worker at the time of exposure. If the type of protection is not listed, describe in OTHER.

Indicate the usual **occupation** of the exposed health care worker. If the occupation is not listed, provide the occupation in OTHER.

Indicate the **department or work area where the incident occurred**. This may be different from the department in which the health care worker is regularly assigned. If the department is not listed, indicate the department in OTHER. In the space provided, indicate the specific location of the incident, such as the room number, or the floor in which the incident occurred.

Indicate whether the department in which the exposure occurred is the department to which the health care worker is regularly assigned. If the answer is no, please indicate the department to which the employee is regularly assigned.

Indicate which **device or item was involved in the injury**. If the device is not listed, indicate the type of device in the space for OTHER in the category of devices provided.

Identify the **brand and or model of the device**. It may be helpful to have the samples or pictures of the types of devices available, with the sharp covered, so that the injured employee can identify the device.

Indicate whether the **device was a safety device**. If yes, indicate **when the injury occurred**, relative to the activation of the safety feature.

APPENDIX E

If the device was a safety device, indicate whether the health care worker was **trained in the use of the device**. Describe the training provided (e.g., printed instructions, on the job, in-service demonstration, hands on in-service).

Identify the **purpose or procedure for which the sharp was used or intended**. If the purpose is not listed, indicate the purpose in OTHER within the category of uses provided.

Choose up to two items describing **how the injury occurred**.

Provide a narrative description of the incident, identifying the events that led up to the incident, as well as if anyone else was involved. Describe the nature of the injury and the body part injured, along with any other information about the incident.

Ask the injured health care worker to suggest ways to prevent this type of injury from occurring in the future. Suggestions may range from increased training, to changing the devices that are utilized within the facility.

The name and title of the individual filling out the recording form as well as the date it is completed shall be recorded.

APPENDIX F

INSTRUCTIONS for MDPH ANNUAL SUMMARY OF SHARPS INJURIES

*This form shall be either typed or computer generated. This form shall cover sharps injuries occurring in the previous calendar year and shall be sent to MDPH-OHSP by **February 1**.*

Provide the **name and license number** of the **hospital** facility.

Provide the **calendar year** of the data.

Provide a **contact name** and **phone number** for the person who is responsible for the data at the hospital.

A **unique exposure incident number** shall be assigned to each exposure event.

Indicate the usual **occupation** of the exposed health care worker. Use of the list of occupations provided on the attached list is encouraged.

Identify the **department or work area** where the incident occurred. Use of the list of departments provided on the attached list is encouraged.

Identify the **device or item** involved in the exposure incident. Use of the list of devices provided on the attached list is encouraged.

Indicate whether the device involved was a **safety device**.

Identify the **brand and model** of the device involved in the exposure incident.

Identify the **purpose or procedure for which the sharp was used or intended**, that is what the device was being utilized for at the time of the exposure incident. Use of the list of procedures provided on the attached list is encouraged.

Indicate **how** the exposure incident occurred. Use of the list of possible causes provided on the attached list is encouraged.

Insert **page numbers** and total number of pages.

Sharps Injury Surveillance Project
Occupational Health Surveillance Program
Massachusetts Department of Public Health
250 Washington Street, 6th floor
Boston, MA 02108

For information on reporting, contact:
Occupational Health Surveillance Program
at Sharps.Injury@state.ma.us or 617-624-5625

APPENDIX F

OCCUPATION				
Attendant / orderly	Dietician	Licensed Practical Nurse	Patient care technician	Respiratory Therapist / tech
Attending physician	EMT / paramedic	Maintenance	Pharmacist	Safety / security
Central supply	Fellow	Morgue technician	Phlebotomist	Transport / messenger
Clerical / administrative	Fireperson / First responder	Medical student	Physician assistant	Volunteer
Clinical lab technician	Food service	Nurse Anesthetist	Physical therapist	Other ancillary staff (specify)
Counselor / social worker	Hemodialysis technician	Nursing Assistant	Public health worker	Other dental worker (specify)
Dentist	Housekeeper	Nurse Midwife	Psychiatric technician	Other medical staff (specify)
Dental assistant / tech	Intern / resident	Nurse Practitioner	Radiologic technician	Other student (specify)
Dental hygienist	Laundry staff	Nursing student	Registered Nurse	Other (specify)
Dental student	Law enforcement officer	OR / surgical technician	Researcher	Other technician (specify)
DEPARTMENT OR WORK AREA WHERE EXPOSURE INCIDENT OCCURRED				
Ambulance	Emergency Department	Home health visit (home)	Medical / surgical ward	Pediatrics
Blood bank	Endoscopy / bronchoscopy	Hospital grounds	Microbiology	Procedure room
Central sterile supply	/cytology	Intensive care unit	Morgue / autopsy room	Psychiatry ward
Central trash area	Exam room	Jail unit	Nursery	Radiology department room
Clinical chemistry	Hematology	Labor and delivery	Obstetrics / gynecology ward	Other laboratory (specify)
Dialysis	Histology / pathology	Laundry room	Operating room	Other outpatient area (specify)
Dental Clinic				Other location (specify)
WHAT DEVICE OR ITEM WAS INVOLVED IN THE INJURY?				
Hollow bore needle	Winged steel needle attached to	Bur	Tenaculum	Suture Needle
Biopsy needle	a vacuum tube collection	Explorer	Trocar	Curved suture needle
IV stylet	holder	Histology cutting blade	Wire	Straight suture needle
Hollow-bore needle, type unknown	Winged steel needle attached to	Lancet	Other type of sharp object	
Huber needle	IV tubing	Laser	(specify)	Additional dental / surgical devices
Hypodermic needle attached to a disposable syringe	Vacuum tube collection holder / needle	Pin	Glass	Hypodermic needle attached to non-disposable syringe
Hypodermic needle attached to IV tubing	Bone chip / chipped tooth	Razor	Capillary tube	Elevator
Prefilled cartridge syringe	Bone cutter	Retractor	Medication ampule / vial / IV bottle	Extraction forceps
Spinal or epidural needle	Bovie electrocautery device	Scaler / curette		Root canal file
Unattached Hypodermic needle		Scalpel blade	Pipette	Rod (orthopaedic)
Winged steel needle		Scissors	Slide	Other device or item (specify)
		Sharp object, type unknown	Specimen / test / vacuum tube	
			Other glass item (specify)	

APPENDIX F

PURPOSE OR PROCEDURE FOR WHICH SHARP WAS USED OR INTENDED:		
<p>Line procedures:</p> <p>To insert a peripheral IV line or set up a heparin lock</p> <p>To insert a central IV line</p> <p>To insert an arterial line</p> <p>To connect IV line (intermittent IV / piggy back / IV infusion / other IV line connection)</p> <p>To flush heparin / saline</p> <p>Other injection into IV injection site or IV port (specify)</p> <p>Other line procedure (specify)</p> <p>Blood procedures:</p> <p>Percutaneous venous puncture (e.g. phlebotomy)</p> <p>Percutaneous arterial puncture</p> <p>Central or peripheral IV line or port</p> <p>Arterial line</p> <p>Dialysis / AV fistula site</p> <p>Umbilical vessel</p> <p>Finger stick / heel stick</p> <p>Other blood sampling (specify)</p>	<p>Other procedures:</p> <p>Cutting (e.g. surgery / autopsy)</p> <p>During disposal</p> <p>Epidural / spinal anesthesia</p> <p>Intramuscular (IM) injection</p> <p>Subcutaneous / intradermal injection / skin test placement</p> <p>Suturing</p> <p>Transferring blood / body fluid to another container</p> <p>To obtain a body fluid or tissue sample (CFS / amniotic / biopsy)</p> <p>To obtain laboratory specimens</p> <p>Other procedure (not a line or blood sampling procedure) (specify)</p> <p>Unknown</p>	<p>Dental procedure:</p> <p>During disposal</p> <p>Hygiene (prophy, root plane, curettage)</p> <p>Oral surgery</p> <p>Simple Extraction</p> <p>Surgical Extraction</p> <p>Fracture Reduction</p> <p>Other (specify)</p> <p>Unknown</p> <p>Periodontal surgery</p> <p>Restorative (amalgam, composite, crown)</p> <p>Root canal</p> <p>Other (specify)</p> <p>Unknown</p> <p><i>Where did the injury occur??</i></p> <p>Inside the patient's mouth</p> <p>Outside the patient's mouth</p> <p>Unknown</p>
<p>HOW DID THE INJURY OCCUR? Choose up to two.</p>		
<p><i>Before use of the item</i></p> <p><i>During use of the item</i></p> <p>Collided with co-worker or other person</p> <p>Collided with sharp</p> <p>Incising</p> <p>Manipulating suture needle in holder</p> <p>Palpating / Exploring</p> <p>Passing or receiving equipment</p> <p>Transferring equipment</p> <p>Patient moved and jarred device</p> <p>Sharp object dropped</p> <p>Suturing</p> <p>Tying sutures</p> <p>While inserting needle in line</p> <p>While inserting needle in patient</p> <p>While manipulating needle in line</p> <p>While manipulating needle in patient</p> <p>While withdrawing needle from line</p> <p>While withdrawing needle from patient</p> <p>Other (specify)</p> <p>Unknown</p>	<p><i>After use, before disposal</i></p> <p>Activating safety device</p> <p>Cap fell off after recapping</p> <p>Collided with co-worker or other person</p> <p>Collided with sharp after procedure</p> <p>Disassembling device or equipment</p> <p>Decontamination / processing of used equipment</p> <p>During clean-up</p> <p>Handling equipment on a tray or stand</p> <p>In transit to disposal</p> <p>Opening / breaking glass containers</p> <p>Processing specimens</p> <p>Passing or transferring equipment</p> <p>Recapping (missed or pierced cap)</p> <p>Sharp object dropped after procedure</p> <p>Struck by detached I.V. line needle</p> <p>Transferring blood / bodily fluids into specimen container</p> <p>Other (specify)</p> <p>Unknown</p>	<p><i>During or after disposal of item</i></p> <p>Collided with co-worker or other person</p> <p>Collided with sharp during / after disposal</p> <p>In trash</p> <p>In linen / laundry</p> <p>In pocket / clothing</p> <p>Left on table / tray</p> <p>Left in bed / mattress</p> <p>On floor</p> <p>Over-filled sharps container</p> <p>Punctured sharps container</p> <p>Protruding from opened container</p> <p>Sharp object dropped during / after disposal</p> <p>Struck by detached I.V. line needle during / after disposal</p> <p>While manipulating container</p> <p>While placing sharp in container, injured by sharp being disposed</p> <p>While placing sharp in container, injured by sharp already in container</p> <p>Other (specify)</p> <p>Unknown</p>

APPENDIX G

TABLE G-1

WORK STATUS OF INJURED WORKER	N	%
Employee	2,776	85
Non-Employee Practitioner	310	9
Student	83	3
Temp / Contract	35	1
Volunteer	1	--
Other	3	--
Not answered	71	2
STATE TOTAL	3,279	100%

TABLE G-2

OCCUPATION OF INJURED WORKER	N	%
Nurse	1,279	39%
RN or LPN	1,128	34
Nursing Assistant	59	2
Patient Care Technician	41	1
Nurse Practitioner	19	<1
Nurse Anesthetist	9	<1
Nursing Student	9	<1
Home Health Aide	9	<1
Nurse Midwife	5	<1
Physician	1,068	33%
Intern / Resident	476	15
MD	372	11
Medical Student	66	2
Fellow	60	2
Physician Assistant	44	1
Surgeon	37	1
Anesthesiologist	9	<1
Radiologist	4	--
Technician	657	20%
OR / Surgical Technician	252	8
Phlebotomist	144	4
Clinical Lab Technician	69	2
Radiologic Technician	60	2
Respiratory Therapist / Tech	22	1
Hemodialysis Technician	6	<1
Other Technician	104	3
Support Services	123	4%
Housekeeper	78	2
Central Supply	27	1
Maintenance	8	<1
Attendant / Orderly	6	<1
Food Service	1	--
Safety / Security	1	--
Transport / Messenger / Porter	1	--
Other Ancillary Staff	1	--

APPENDIX G

TABLE G-2

OCCUPATION OF INJURED WORKER	N	%
Other Medical Staff	31	1%
Medical Assistant	30	1
Physical Therapist	1	--
Dental Staff	14	<1%
Dentist	6	<1
Dental Assistant / Tech	4	--
Dental Student	3	--
Dental Hygienist	1	--
Other	106	3%
EMT / Paramedic	7	--
Clerical / Administrative	5	--
Counselor / Social Worker	2	--
Pharmacist	1	--
Researcher	1	--
Other student	15	<1
Other	75	2
Not answered	1	--%
STATE TOTAL	3,279	100%

TABLE G-3

DEPARTMENT WHERE INCIDENT OCCURRED	N	%
Operating and Procedure Rooms	1,460	45%
Operating room	1,038	32
Labor and delivery	151	5
Radiology	139	4
Cardiac catheterization laboratory	35	1
Hematology / Oncology	29	1
Dialysis	27	1
Endoscopy / Bronchoscopy / Cytoscopy	19	1
Phlebotomy room	5	<1
Procedure room, not specified	17	1
Inpatient units	732	22%
Medical / Surgical ward	570	17
Pediatrics	29	1
Obstetrics / Gynecology	24	1
Psychiatry ward	20	1
Nursery	16	<1
Specific ward, type unknown**	8	<1
Patient room, ward unspecified	65	2
Emergency Department	263	8%

APPENDIX G

TABLE G-3

DEPARTMENT WHERE INCIDENT OCCURRED

	N	%
Intensive Care Units	248	8%
Intensive care unit	231	7
Post anesthesia care unit	17	1
Outpatient areas	171	5%
Home health visit	24	1
Dental Clinic	19	1
Ambulatory care clinic	9	<1
Other outpatient areas	119	4
Laboratory	156	5%
Histology / Pathology	22	1
Morgue / Autopsy room	12	<1
Clinical chemistry	9	<1
Blood bank	6	<1
Microbiology	1	--
Other laboratory	36	1
Other areas	248	8%
Rehabilitation unit	34	1
Central Sterile Supply	30	1
Exam room	17	1
Dermatology	15	<1
Anesthesia	9	<1
Pain clinic	7	<1
Employee health / Infection control	5	<1
Pharmacy	5	<1
Detox unit	3	--
Central trash area	2	--
Ambulance	1	--
Jail unit	1	--
Laundry room	1	--
Long term care	1	--
Other location	113	3
Unknown / Not answered	1	--%
STATE TOTAL	3,279	100%

APPENDIX G

TABLE G-4

PROCEDURE FOR WHICH DEVICE WAS USED

	N	%
Injection	754	23%
Subcutaneous injection	370	11
Intramuscular injection	63	2
Epidural / Spinal anesthesia	7	<1
Injection, unspecified	314	10
Suturing	729	22%
Suturing	725	22
Suture removal	4	--
Blood procedures	624	19%
Percutaneous venous puncture	418	13
Percutaneous arterial puncture	63	2
Finger stick / Heel stick	57	2
Draw blood from central or peripheral IV line or port	21	1
Dialysis / AV fistula site	14	<1
Draw blood from arterial line	8	<1
Draw blood from umbilical vessel	8	<1
Other blood sampling	35	1
Line procedures	290	9%
To insert a peripheral IV line or set up a heparin lock	116	4
Other injection into IV site / port	45	1
To flush heparin / saline	28	1
To insert a central IV line	20	1
To insert an arterial line	20	1
To connect IV line	4	--
Other line procedure	57	2
Making the incision	242	7%
To obtain body fluid or tissue sample	74	2%
Dental procedures	6	<1%
Restorative	3	--
Dental drilling	1	--
Hygiene	1	--
Other dental	1	--
Other	309	9%
Transferring blood / body fluid to another container	23	1
Drilling	17	1
To obtain lab specimens	16	<1
Shaving	9	<1
Other procedure	244	7
Unknown / Not answered	251	8%
STATE TOTAL	3,279	100%

APPENDIX G

TABLE G-5

DEVICE INVOLVED IN THE INJURY

	N	%
Hypodermic needle	1,009	31%
Hypodermic needle attached to a disposable syringe	837	26
Prefilled cartridge syringe	59	2
Hypodermic needle attached to a non-disposable syringe	48	1
Unattached hypodermic needle	48	1
Hypodermic needle attached to IV tubing	4	--
Suture needle	722	22%
Curved suture needle	175	5
Straight suture needle	26	1
Suture needle, unspecified	521	16
Other hollow bore needle	349	11%
IV stylet	152	5
Huber needle	46	1
Spinal or epidural needle	21	1
Biopsy needle	19	1
Hollow bore needle, type unknown	101	3
Other type of hollow bore needle	10	<1
Butterfly needle	335	10%
Winged steel needle	253	8
Winged steel needle attached to a vacuum tube collection holder	69	2
Winged steel needle attached to IV tubing	13	<1
Scalpel blade	229	7%
Vacuum tube collection holder / needle	143	4%
Vacuum tube collection holder / needle	117	4
Phlebotomy needle (other than butterfly)	26	1
Glass	35	1%
Specimen / Test / Vacuum tube	19	1
Medication ampule / Vial / IV bottle	5	<1
Pipette	2	--
Slide	2	--
Other glass item	7	<1
Dental device or item	11	<1%
Dental explorer	3	--
Dental bur	2	--
Dental needle	2	--
Scaler / Curette	1	--
Other dental device or item	3	--

APPENDIX G

Other	381	12%
Lancet	55	2
Wire	54	2
Scissors	28	1
Pin	21	1
Forceps	20	1
Retractor	20	1
Razor	17	1
Drill bit	15	<1
Bovie electrocautery device	13	<1
Trocar	13	<1
Bone cutter	7	<1
Bone chip / chipped tooth	2	--
Elevator	1	--
Tenaculum	1	--
Other needle	47	1
Other type of sharp object	65	2
Unknown / Not answered	65	2%
STATE TOTAL	3,279	100%

TABLE G-6

SAFETY DEVICE	N	%
No	1,773	54
Yes	1,072	33
Unknown / Not answered	434	13
STATE TOTAL	3,279	100%

TABLE G-7

WHEN THE INJURY OCCURRED	N	%
During Use of the Item	1,363	42
After Use and Before Disposal	1,105	34
During or After Disposal of the Item	498	15
Before Use of the Item	53	2
Unknown / Not answered	260	8
STATE TOTAL	3,279	100%

APPENDIX G

TABLE G-8

HOW THE INJURY OCCURRED

	N	%
Collision with worker or sharp	751	23%
Collided with sharp	332	10
Collided with sharp after procedure	272	8
Collided with coworker or other person	147	4
Suturing	387	12%
Suturing	345	11
Manipulating suture needle in holder	31	1
Tying suture	11	<1
Improper disposal	250	8%
In trash	78	2
Left on table / tray	58	2
Left in bed / mattress	36	1
On floor	20	1
In pocket / clothing	10	<1
In linen / laundry	3	--
Improper disposal, unspecified	45	1
During sharps disposal	247	8%
Collided with sharp during / after disposal	81	2
While placing sharp in container, injured by sharp being disposed	71	2
In transit to disposal	33	1
Protruding from opened container	21	1
Overfilled sharps container	14	<1
While placing sharp in container, injured by sharp already in container	13	<1
While manipulating container	10	<1
Punctured sharps container	4	--
During clean-up	224	7%
During clean-up	113	3
Disassembling device or equipment	74	2
Decontamination / Processing of used equipment	37	1
Handle / pass equipment	210	6%
Receiving / Passing / Transferring equipment	115	4
Handling equipment on tray or stand	80	2
Opening / breaking glass containers	15	<1
Patient moved and jarred device	175	5%
Activating safety device	156	5%
Manipulate needle in patient	121	4%
While withdrawing needle from patient	85	3
While inserting needle in patient	22	1
While manipulating needle in patient	14	<1

APPENDIX G

TABLE G-8

HOW THE INJURY OCCURRED

	N	%
Recap needle	107	3%
Recapping	100	3
Cap fell off after recapping	7	<1
Access IV line	82	3%
While withdrawing needle from line	50	2
While inserting needle in line	14	<1
While manipulating needle in line	11	<1
Struck by detached IV line needle	7	<1
Failure to activate safety device	66	2%
Device malfunction	36	1%
Before use of the item	43	1%
Other	376	11%
Incising	66	2
Sharp object dropped	30	1
Processing specimens	23	1
Transferring blood / bodily fluids into specimen container	17	1
Sharp object dropped after procedure	9	<1
Palpating / Exploring	1	--
Other	230	7
Unknown / Not answered	48	1%
STATE TOTAL	3,279	100%

** Hospital specific nomenclature provided, without specifying department

Percentages for frequencies less than 5 were not calculated; Percentages calculated are column percents.

APPENDIX H

Sharps Injuries among Hospital Workers by Number of Licensed Hospital Beds, Massachusetts, 2004

	Number of Licensed Hospital Beds							
	0-100 Beds		101-300 Beds		300+ Beds		All Hospitals	
	32 hospitals		51 hospitals		16 hospitals		99 hospitals	
	N	%*	N	%*	N	%*	N	%*
STATE TOTAL	277	100%	1,182	100%	1,820	100%	3,279	100%

WORK STATUS OF INJURED WORKER

Employee	220	79 %	990	84 %	1,566	86 %	2,776	85 %
Non-Employee Practitioner	22	8	101	9	187	10	310	9
Student	1	--	29	2	53	3	83	3
Temp / Contract	7	3	16	1	12	1	35	1
Volunteer	0	--	0	--	1	--	1	--
Other	3	--	0	--	0	--	3	--
Unknown / Not answered	24	9	46	4	1	--	71	2
STATE TOTAL	277	100%	1,182	100%	1,820	100%	3,279	100%

OCCUPATION

Nurse	122	44 %	518	44 %	639	35 %	1,279	39 %
Physician	60	22	228	19	780	43	1,068	33
Technician	70	25	329	28	258	14	657	20
Support Services	11	4	58	5	54	3	123	4
Other Medical Staff	5	2	16	1	10	1	31	1
Dental Staff	1	--	1	--	12	1	14	<1
Other	8	3	32	3	66	4	106	3
Unknown / Not answered	0	--	0	--	1	--	1	--
STATE TOTAL	277	100%	1,182	100%	1,820	100%	3,279	100%

DEPARTMENT WHERE INJURY OCCURRED

Operating and procedure rooms	122	44 %	464	39 %	874	48 %	1,460	45 %
Inpatient units	59	21	291	25	382	21	732	22
Emergency Department	20	7	109	9	134	7	263	8
Intensive Care Units	11	4	88	7	149	8	248	8
Laboratories	19	7	62	5	75	4	156	5
Outpatient areas	25	9	51	4	95	5	171	5
Other areas	21	8	117	10	110	6	248	8
Unknown / Not answered	0	--	0	--	1	--	1	--
STATE TOTAL	277	100%	1,182	100%	1,820	100%	3,279	100%

PROCEDURE FOR WHICH DEVICE WAS USED

Injection	56	20 %	310	26 %	388	21 %	754	23 %
Suturing	53	19	221	19	455	25	729	22
Blood procedures	62	22	267	23	295	16	624	19
Line procedures	30	11	95	8	165	9	290	9
Making the incision	18	7	77	7	147	8	242	7
To obtain body fluid or tissue sample	4	--	31	3	39	2	74	2
Dental procedures	0	--	3	--	3	--	6	<1
Other	36	13	72	6	201	11	309	9
Unknown / Not answered	18	7	106	9	127	7	251	8
STATE TOTAL	277	100%	1,182	100%	1,820	100%	3,279	100%

Percentages calculated are column percents; Percentages for frequencies less than 5 were not calculated.

APPENDIX H

Sharps Injuries among Hospital Workers by Number of Licensed Hospital Beds, Massachusetts, 2004

	Number of Licensed Hospital Beds							
	0-100 Beds		101-300 Beds		300+ Beds		All Hospitals	
	32 hospitals		51 hospitals		16 hospitals		99 hospitals	
	N	%*	N	%*	N	%*	N	%*
STATE TOTAL	277	100%	1,182	100%	1,820	100%	3,279	100%
DEVICE INVOLVED IN THE INJURY								
Hypodermic needle	73	26 %	399	34%	537	30%	1,009	31 %
Suture needle	54	19	217	18	451	25	722	22
Butterfly needle	32	12	164	14	139	8	355	10
Scalpel blade	13	5	70	6	146	8	229	7
Vacuum tube collection holder / needle	20	7	57	5	66	4	143	4
Glass	3	--	11	1	21	1	35	1
Dental device or item	2	--	3	--	6	<1	11	<1
Other hollow bore needle	29	10	114	10	206	11	349	11
Other	42	15	124	10	215	12	381	12
Unknown / Not answered	9	3	23	2	33	2	65	2
STATE TOTAL	277	100%	1,182	100%	1,820	100%	3,279	100%
SAFETY DEVICE								
No	146	53 %	584	49%	1,043	57%	1,773	54 %
Yes	101	36	508	43	463	25	1,072	33
Unknown / Not answered	30	11	90	8	314	17	434	13
STATE TOTAL	277	100%	1,182	100%	1,820	100%	3,279	100%
WHEN THE INJURY OCCURRED								
During Use of the Item	94	34 %	423	36%	846	46%	1,363	42 %
After Use / Before Disposal	107	39	444	38	554	30	1,105	34
During or after disposal of the item	50	18	209	18	239	13	498	15
Before use of the item	6	2	15	1	32	2	53	2
Unknown / Not answered	20	7	91	8	149	8	260	8
STATE TOTAL	277	100%	1,182	100%	1,820	100%	3,279	100%
HOW THE INJURY OCCURRED								
Collision with worker or sharp	45	16 %	244	21%	462	25%	751	23 %
Suturing	30	11	87	7	270	15	387	12
Improper disposal	20	7	109	9	121	7	250	8
During sharps disposal	29	10	99	8	119	7	247	8
During clean-up	33	12	86	7	105	6	224	7
Handle / pass equipment	18	7	76	6	116	6	210	6
Patient moved / jarred device	19	7	82	7	74	4	175	5
Activate safety device	15	5	78	7	63	3	156	5
Manipulate needle in patient	5	2	47	4	69	4	121	4
Recap needle	7	3	40	3	60	3	107	3
Access IV line	13	5	26	2	43	2	82	3
Failure to activate safety device	8	3	41	3	17	1	66	2
Before use of item	4	--	12	1	27	1	43	1
Device malfunctioned	4	--	21	2	11	1	36	1
Other	23	8	117	10	236	13	376	11
Unknown / Not answered	4	--	17	1	27	1	48	1
STATE TOTAL	277	100%	1,182	100%	1,820	100%	3,279	100%

Percentages calculated are column percents; Percentages for frequencies less than 5 were not calculated.

APPENDIX I

Sharps Injuries among Hospital Workers by Teaching Status, Massachusetts, 2004

	Teaching Status					
	Teaching		Non-teaching		All Hospitals	
	18 hospitals		81 hospitals		99 hospitals	
	N	%*	N	%*	N	%*
STATE TOTAL	1,923	100%	1,356	100%	3,279	100%

WORK STATUS OF INJURED WORKER

Employee	1,662	86%	1,114	82%	2,776	85%
Non-Employee Practitioner	180	9	130	10	310	9
Student	57	3	26	2	83	3
Temp / Contract	15	1	20	1	35	1
Volunteer	1	--	0	0	1	--
Other	3	--	0	0	3	--
Unknown / Not answered	5	<1	66	5	71	2
STATE TOTAL	1,923	100%	1,356	100%	3,279	100%

OCCUPATION

Nurse	650	34%	629	46%	1,279	39%
Physician	837	44	231	17	1,068	33
Technician	280	15	377	28	657	20
Support Services	56	3	67	5	123	4
Other Medical Staff	15	1	16	1	31	1
Dental Staff	13	1	1	--	14	<1
Other	71	4	35	3	106	3
Unknown / Not answered	1	--	0	0	1	--
STATE TOTAL	1,923	100%	1,356	100%	3,279	100%

DEPARTMENT WHERE INJURY OCCURRED

Operating and procedure rooms	930	48%	530	39%	1,460	45%
Inpatient units	380	20	352	26	732	22
Emergency Department	134	7	129	10	263	8
Intensive Care Units	157	8	91	7	248	8
Outpatient areas	119	6	52	4	171	5
Laboratories	81	4	75	6	156	5
Other areas	121	6	127	9	248	8
Unknown / Not answered	1	--	0	0	1	--
STATE TOTAL	1,923	100%	1,356	100%	3,279	100%

PROCEDURE FOR WHICH DEVICE WAS USED

Injection	400	21%	354	26%	754	23%
Suturing	484	25	245	18	729	22
Blood procedures	316	16	308	23	624	19
Line procedures	166	9	124	9	290	9
Making the incision	159	8	83	6	242	7
To obtain body fluid or tissue sample	37	2	37	3	74	2
Dental procedures	5	<1	1	--	6	<1
Other	217	11	92	7	309	9
Unknown / Not answered	139	7	112	8	251	8
STATE TOTAL	1,923	100%	1,356	100%	3,279	100%

Percentages calculated are column percents; Percentages for frequencies less than 5 were not calculated.

APPENDIX I

Sharps Injuries among Hospital Workers by Teaching Status, Massachusetts, 2004

	Teaching Status					
	Teaching		Non-teaching		All Hospitals	
	18 hospitals		81 hospitals		99 hospitals	
	N	%*	N	%*	N	%*
STATE TOTAL	1,923	100%	1,356	100%	3,279	100%
DEVICE INVOLVED IN THE INJURY						
Hypodermic needle	545	28%	464	34%	1,009	31%
Suture needle	478	25	244	18	722	22
Butterfly needle	160	8	175	13	335	10
Scalpel blade	158	8	71	5	229	7
Vacuum tube collection holder / needle	68	4	75	6	143	4
Glass	20	1	15	1	35	1
Dental device or item	8	<1	3	--	11	<1
Other hollow bore needle	213	11	136	10	349	11
Other	239	12	142	10	381	12
Unknown / Not answered	34	2	31	2	65	2
STATE TOTAL	1,923	100%	1,356	100%	3,279	100%
SAFETY DEVICE						
No	1,116	58%	657	48%	1,773	54%
Yes	483	25	589	43	1,072	33
Unknown / Not answered	324	17	110	8	434	13
STATE TOTAL	1,923	100%	1,356	100%	3,279	100%
WHEN THE INJURY OCCURRED						
During Use of the Item	887	46%	476	35%	1,363	42%
After Use / Before Disposal	596	31	509	38	1,105	34
During or after disposal of the item	248	13	250	18	498	15
Before use of the item	34	2	19	1	53	2
Unknown / Not answered	158	8	102	8	260	8
STATE TOTAL	1,923	100%	1,356	100%	3,279	100%
HOW THE INJURY OCCURRED						
Collision with worker or sharp	487	25%	264	19%	751	23%
Suturing	287	15	100	7	387	12
Improper disposal	123	6	127	9	250	8
During sharps disposal	126	7	121	9	247	8
During clean-up	122	6	102	8	224	7
Handle / pass equipment	121	6	89	7	210	6
Patient moved / jarred device	78	4	97	7	175	5
Activate safety device	71	4	85	6	156	5
Manipulate needle in patient	70	4	51	4	121	4
Recap needle	69	4	38	3	107	3
Access IV line	48	3	34	3	82	3
Failure to activate safety device	18	1	48	4	66	2
Before use of the item	27	1	16	1	43	1
Device malfunctioned	14	1	22	2	36	1
Other	238	12	138	10	376	11
Unknown / Not answered	24	1	24	2	48	1
STATE TOTAL	1,923	100%	1,356	100%	3,279	100%

Percentages calculated are column percents; Percentages for frequencies less than 5 were not calculated.

APPENDIX J

Resources Sharps Injury Surveillance and Prevention

MDPH Occupational Health Surveillance Program

<http://www.mass.gov/dph/ohsp>

Sharps Injury Surveillance and Prevention Project - e-mail: Sharps.Injury@state.ma.us

OSHA Subject Page for Needle Sticks

Includes Bloodborne Pathogens Standard and compliance directive

<http://www.osha.gov/SLTC/bloodbornepathogens/index.html>

CDC-MMWR September 30, 2005 / Vol. 54 / RR-9

Updated U.S. Public Health Service Guidelines for the Management of Occupational Exposures to HIV and Recommendations for Post Exposure Prophylaxis

<http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5409a1.htm>

CDC-MMWR June 29, 2001 / Vol. 50 / RR-11

Updated U.S. Public Health Service Guidelines for the Management of Occupational Exposures to HBV, HCV and HIV and Recommendations for Post Exposure Prophylaxis

<http://www.cdc.gov/mmwr/PDF/rr/rr5011.pdf>

CDC Division of Healthcare Quality Promotion

Workbook for Designing, Implementing, and Evaluating a Sharps Injury Prevention Program

<http://www.cdc.gov/sharpsafety/>

CDC Division of Healthcare Quality Promotion, Issues in Healthcare

Information related to bloodborne pathogens

<http://www.cdc.gov/ncidod/hip/Blood/blood.htm>

CDC Division of Healthcare Quality Promotion, National Surveillance System for Health care Workers

<http://www.cdc.gov/ncidod/hip/SURVEILL/nash.HTM>

National Surveillance System for Health care Workers,

Summary report for data collected from June 1995 through July 1999

<http://www.cdc.gov/ncidod/hip/NASH/report99.PDF>

NIOSH Alert – Preventing Needlestick Injuries in Health care settings

<http://www.cdc.gov/niosh/2000-108.html>

JCAHO Sentinel Event Alert, Issue 22 August 2001

Preventing Needlestick and Sharps Injuries

http://www.jcaho.org/edu_pub/sealert/sea22.html

EPINet, International Health Care Worker Safety Center, University of Virginia

<http://www.med.virginia.edu/medcntr/centers/epinet/>

Training for Development of Innovative Control Technologies (TDICT) Project, San Francisco General Hospital

<http://www.tdict.org/>

Sustainable Hospitals Project, Lowell Center for Sustainable Production, University of Massachusetts Lowell

<http://sustainablehospitals.org>

