


Massachusetts 2018 Air Quality Report



Department of Environmental Protection
Bureau of Air and Waste
Division of Air and Climate Programs

Air Assessment Branch
Wall Experiment Station
37 Shattuck Street
Lawrence, Massachusetts 01843

August 2019

ACKNOWLEDGEMENTS

This 2018 Air Quality Report was prepared by the Massachusetts Department of Environmental Protection (MassDEP), Air Assessment Branch (AAB), which collects representative samples of ambient air for a number of pollutants at monitoring stations located across the Commonwealth. All samples are collected in a precise and scientifically sound manner in order to properly characterize the quality of the air in the Commonwealth.

The photo on the cover is a view of the near-road air monitoring station on Manning Road in Chelmsford, MA.

This report is available on MassDEP's web site at

www.mass.gov/eea/agencies/massdep/air/quality/air-monitoring-reports-and-studies.html

Questions about this report may be directed to:

Sean Dunn

Air Assessment Branch

Wall Experiment Station

Lawrence, MA 01843-1343

(978) 242-1335

email: sean.dunn@mass.gov

TABLE OF CONTENTS

LIST OF FIGURES	ii
-----------------------	----

LIST OF ABBREVIATIONS	iii
-----------------------------	-----

SECTION I – AMBIENT AIR MONITORING PROGRAM

Program Overview	1
National Ambient Air Quality Standards.....	4
Pollutant Health Effects and Sources.....	5
Monitoring Network Description.....	7

SECTION II – ATTAINMENT OF AIR QUALITY STANDARDS

Attainment Status Summary	9
2018 Ozone Season.....	11
Daily Ozone and PM Forecast	13

SECTION III – MASSACHUSETTS AIR QUALITY DATA SUMMARIES

Ozone Summary.....	14
Sulfur Dioxide (SO ₂) Summary	18
Nitrogen Dioxide (NO ₂) Summary	20
Carbon Monoxide (CO) Summary.....	22
Particulate Matter 10 Microns (PM ₁₀) Summary.....	24
Particulate Matter 2.5 Microns (PM _{2.5}) Summary	26
Speciation.....	31
IMPROVE.....	31
Quality Control and Quality Assurance	31

SECTION IV – PAMS/AIR TOXICS MONITORING

PAMS Monitoring	32
Air Toxics Monitoring	33

APPENDIX A – 2018 Monitoring Stations.....	35
--	----

List of Figures

Section II – Attainment of Air Quality Standards

Figure 1	8-hour Ozone Exceedance Days and Total Exceedances 2009-2018.....	12
Figure 2	O ₃ NAAQS Levels 2016-2018	

Section III – Massachusetts Air Quality Data Summaries

Figure 3	8-hour Ozone Exceedance Day Trends 2009-2018	16
Figure 4	Sulfur Dioxide Trends 2009-2018	19
Figure 5	Nitrogen Dioxide Trends 2009-2018	21
Figure 6	Carbon Monoxide Trends 2009-2018.....	23
Figure 7	Particulate Matter 10 Microns (PM ₁₀) Trends 2009-2018.....	25
Figure 8	PM ₁₀ Trends 2009-2018	
Figure 9	Particulate Matter 2.5 Microns (PM _{2.5}) FRM Trends 2009-2018.....	28
Figure 10	Particulate Matter 2.5 Microns (PM _{2.5}) FEM Trends 2009-2018.....	30

Section IV – PAMS/Air Toxics Monitoring

Figure 11	Lynn Toxics VOC Summary 2009-2018.....	33
-----------	--	----

List of Abbreviations

AAB	Air Assessment Branch
AQS	Air Quality System
AQI.....	Air Quality Index
BAM	Beta Attenuation Monitor
BC	Black Carbon
BP.....	Barometric Pressure
CAA	Clean Air Act
CFR.....	Code of Federal Regulations
CO.....	Carbon Monoxide
CO ₂	Carbon Dioxide
FEM	Federal Equivalent Method
FRM	Federal Reference Method
EPA.....	United States Environmental Protection Agency
IMPROVE	Interagency Monitoring of Protected Visual Environments
MassDEP	Massachusetts Department of Environmental Protection
NAAQS.....	National Ambient Air Quality Standards (for criteria pollutants)
NATTS.....	National Air Toxics Trends Station
NCore	National Core Monitoring Network
NO.....	Nitric Oxide
NO _x	Nitrogen Oxides
NO _y	Total Reactive Oxidized Nitrogen
NO ₂	Nitrogen Dioxide
NO ₃	Nitrate
O ₃	Ozone
PAH	Polycyclic Aromatic Hydrocarbon
PAMS.....	Photochemical Assessment Monitoring Stations
Pb	Lead
pH.....	Concentration of hydrogen cations (H ⁺) in solution (an indicator of acidity)
ppb	parts per billion by volume
ppm	parts per million by volume
PM _{2.5}	Particulate matter ≤ 2.5 microns aerodynamic diameter
PM ₁₀	Particulate matter ≤ 10 microns aerodynamic diameter
QA/QC	Quality Assurance and Quality Control
RH.....	Relative Humidity
SIP.....	State Implementation Plan
SO ₂	Sulfur Dioxide
SO ₄	Sulfate
SR.....	Solar Radiation
SVOC.....	Semi-Volatile Organic Compounds
TEMP.....	Temperature
TSA.....	Technical Systems Audit
TSP.....	Total Suspended Particulates
µg/m ³	micrograms per cubic meter
VOCs	Volatile Organic Compounds
WS/WD.....	Wind Speed/Wind Direction
WSv/WDv.....	Wind Speed Vector/Wind Direction Vector

Section I

Ambient Air Monitoring Program

Program Overview

Introduction

The Massachusetts Department of Environmental Protection (MassDEP) is the state agency responsible for monitoring outdoor air quality in Massachusetts and developing plans and regulatory programs to reduce emissions of pollutants that adversely affect public health, welfare, and the environment.

MassDEP's Air Assessment Branch (AAB) operates a network of air monitoring stations throughout the Commonwealth. During 2018, MassDEP operated 22 monitoring stations located in 17 cities and towns. MassDEP also received data from the Wampanoag Tribe of Gay Head (Aquinnah), which operates an air monitoring station on Martha's Vineyard.

MassDEP submits ambient air quality data to the national Air Quality System (AQS) database that is administered by the U.S. Environmental Protection Agency (EPA). Continuous monitoring data is sent to the AIRNOW website, which reports data from all U.S. monitoring stations. MassDEP's MassAir Online website provides air quality information and allows users to point and click on a map of the state to find current air quality data from the MassDEP continuous air monitoring network. MassAir Online is found at www.mass.gov/eea/agencies/massdep/air/quality/. EPA also makes historical AQS data for all U.S. monitoring stations available at <https://www.epa.gov/outdoor-air-quality-data>.

Why is Air Quality Data Collected?

Ambient air quality data is used for a number of purposes, including to:

- Provide information about air quality to the public;
- Provide short-term and long-term information regarding air pollution and public health;
- Verify compliance with National Ambient Air Quality Standards (NAAQS);
- Assess the effectiveness of current air pollution control regulations and initiatives;
- Support development of policies and regulations aimed at reducing air pollution;
- Support long-term trend analysis and special research; and
- Fulfill requirements to report ambient air quality data to EPA.

What is Monitored?

MassDEP monitors parameters in the following categories:

Criteria pollutants for which EPA has established NAAQS. The criteria pollutants monitored are:

- sulfur dioxide (SO₂)
- ozone (O₃)
- carbon monoxide (CO)
- nitrogen dioxide (NO₂)
- lead (Pb)
- particulate matter ≤10 microns (PM₁₀)
- particulate matter ≤ 2.5 microns (PM_{2.5})

Non-criteria pollutants do not have NAAQS, but can contribute to the formation of ozone and particulate matter and/or be toxic. The non-criteria pollutants monitored include:

- nitric oxide (NO)
- total nitrogen oxides (NO_x)
- total reactive oxidized nitrogen (NO_y)
- volatile organic compounds (VOCs) – ozone precursors and reaction product chemicals
- black carbon (i.e., soot)
- toxics – health-relevant VOCs, semi-volatile organic compounds (SVOCs), carbonyls and metals

Meteorological parameters monitored include:

- wind speed/wind direction (WS/WD) and vector (WSv/WDv)
- relative humidity (REL)
- temperature (TEM)
- barometric pressure (BP)
- solar radiation (SUN)
- precipitation (PRECIP)


Monitoring Station Locations

Monitoring stations are sited to provide data for various purposes. Some are located where maximum pollutant concentrations are expected, while others are positioned in areas that will provide data that is representative of larger geographical areas. Local topography and pollutant source areas are factors that determine how well a particular monitor's location will represent a region.

Networks of monitors are located throughout the state. These networks are designed to reflect pollutant concentrations for all of Massachusetts. Section III of this report contains data summaries for each pollutant measured and maps showing the monitor locations for each network. Appendix A contains a list of monitoring stations.

The map on page 3 shows Massachusetts cities and towns where air monitors were located during 2018.

2018 Air Monitoring Network



National Ambient Air Quality Standards (NAAQS)

Below are the current NAAQS for criteria pollutants set by EPA. **Primary Standards** are designed to protect public health, including the health of sensitive populations such as asthmatics, children, and the elderly. **Secondary Standards** are designed to protect public welfare, including protection against decreased visibility, damage to crops, vegetation, and buildings.

National Ambient Air Quality Standards					
Pollutant		Primary/ Secondary	Averaging Time	Level	Form
Carbon Monoxide		Primary	8-hour	9 ppm	Not to be exceeded more than once per year
			1-hour	35 ppm	
Lead		Primary and secondary	Rolling 3 month average	0.15 µg/m ³	Not to be exceeded
Nitrogen Dioxide		Primary	1-hour	100 ppb	98 th percentile, averaged over 3 years
		Primary and secondary	Annual	0.053 ppm	Annual Mean
Ozone		Primary and secondary	8-hour	0.070 ppm	Annual fourth-highest daily maximum 8-hour concentration, averaged over 3 years
Particle Pollution	PM _{2.5}	Primary	Annual	12 µg/m ³	Annual mean, averaged over 3 years
		Secondary	Annual	15 µg/m ³	Annual mean, averaged over 3 years
		Primary and secondary	24-hour	35 µg/m ³	98 th percentile, averaged over 3 years
	PM ₁₀	Primary and secondary	24-hour	150 µg/m ³	Not to be exceeded more than once per year on average over 3 years
Sulfur Dioxide		Primary	1-hour	75 ppb	99 th percentile of 1-hour daily maximum concentrations, averaged over 3 years
		Secondary	3-hour	0.5 ppm	Not to be exceeded more than once per year

µg/m³ = micrograms per cubic meter; ppm = parts per million; ppb = parts per billion

Pollutant Health Effects and Sources

Ozone (O₃)

- Tropospheric O₃ (ground-level) and Stratospheric O₃ (upper atmosphere) are the same chemical compound, just found at different places in the atmosphere. Stratospheric O₃ found at greater than 30,000 feet above the surface of the earth is beneficial to all life because it filters out the sun's harmful UV radiation before it reaches the earth's surface. Ground-Level O₃ on the other hand is a health and environmental problem. This report pertains exclusively to ground-level O₃.
- O₃ is a respiratory irritant that can reduce lung function and resistance to infection. It can cause asthma attacks, nasal congestion, and throat irritation. It can inflame and damage (possibly permanently) cells that line the lungs, and aggravate chronic lung diseases. In addition, a number of studies have found a strong link between increases in ground-level O₃ and increased risk of premature death.
- O₃ is toxic to vegetation, inhibiting growth and causing leaf damage.
- O₃ deteriorates materials such as rubber and fabrics.
- Ground-level O₃ is unique in that it is formed by the reactions that occur between certain pollutants in the presence of intense, high-energy sunlight during the hot summer months. The complexity of the reactions and the amount of time needed to complete these reactions can result in the buildup of ground-level ozone concentrations far downwind from the original source of the precursors.
- Sources of ground-level O₃ precursors, i.e., nitrogen oxides and hydrocarbons, include motor vehicles, lawn and garden equipment, power plants and other industrial sources.

Carbon Monoxide (CO)

- CO binds with hemoglobin in the blood, reducing the amount of oxygen carried to organs and tissues.
- Symptoms of high CO exposure include shortness of breath, chest pain, headaches, confusion, and loss of coordination. The health threat is most severe for those with cardiovascular disease.
- Motor vehicle emissions are the largest source of CO, which is produced from incomplete combustion of carbon in fuels.
- Industrial processes and non-transportation fuel combustion (e.g., boilers, lawn and garden equipment) also are sources of CO.

Sulfur Dioxide (SO₂)

- SO₂ combines with water vapor to form acidic aerosols harmful to the respiratory tract, aggravating symptoms associated with lung diseases such as asthma and bronchitis.
- SO₂ is a primary contributor to acid deposition. Impacts of acid deposition include: acidification of lakes and streams, damage to vegetation, damage to materials, and diminution of visibility.
- SO₂ is a product of fuel combustion (e.g., the burning of coal and oil that contains sulfur). Sources include power plants and business and residential sources burning heating oil.

Nitrogen Dioxide (NO₂)

- NO₂ lowers resistance to respiratory infections and aggravates symptoms associated with asthma and bronchitis.
- NO₂ contributes to acid deposition. Impacts of acid deposition include: acidification of lakes and streams, damage to vegetation, damage to materials, and diminution of visibility.
- NO₂ and nitric oxide (NO) contribute to the formation of ozone.
- NO₂ is formed from the oxidation of NO. Major sources of NO are fuel combustion, space heating, power plants and motor vehicles.

Particulate Matter (PM₁₀ and PM_{2.5})

- Particulate matter is tiny airborne particles or aerosols, which include dust, dirt, soot, smoke, and liquid droplets. Fine particulate matter (mostly below 2.5 microns in size) are not only the result of direct emissions, but can be formed in the atmosphere by chemical reactions involving gaseous pollutants.
- The numbers 2.5 and 10 refer to the particle size (actually the particles equal to or less than that size), measured in microns, which are collected by the monitors. Several thousand PM_{2.5} particles could fit on the period at the end of this sentence.
- The small size of these particles allows easy entry into the human respiratory system. Long-term exposure causes the particles to accumulate in the lungs and affects breathing and produces respiratory symptoms. The small particles can migrate through the lungs and into the circulatory system and potentially produce cardio-vascular symptoms, as well as impacts from toxic components contained in the particulate matter.
- Particulate matter causes soiling and corrosion of materials.
- Particulate matter contributes to atmospheric haze that degrades visibility.
- Sources of particulates include industrial process emissions, motor vehicles, incinerators, power plants, and other fuel combustion sources.

Lead (Pb)

- Lead is an elemental metal that is found in nature.
- Exposure to lead can occur by inhalation or ingestion with food, water, soil or dust particles.
- Children, infants, and fetuses are the most susceptible to the effects of lead exposure.
- Lead causes intellectual disability, brain damage, and liver disease. It may be a factor in high blood pressure and damages the nervous system.
- Lead enters the atmosphere from the incineration of lead containing materials and from the manufacture and processing of lead containing products or materials like storage batteries, smelting and removal of lead-containing paint.

Monitoring Network Description

The following describes the ambient air monitoring network in 2018.

Network Size

- 22 monitoring stations
- 17 cities and towns with monitoring stations

Number of Continuous Monitors

Continuous monitors measure air quality 24 hours per day. The data are reported as hourly means.

- Criteria pollutant monitors measure pollutants for which NAAQS have been set.
 - ❑ 5 – Trace-level CO monitors
 - ❑ 11 – NO₂, NO and NO_x monitors
 - ❑ 16 – O₃ monitors
 - ❑ 7 – Trace-level SO₂ monitors
 - ❑ 18 – PM_{2.5} Beta Attenuation Monitors (BAMs)
- Meteorological monitors track weather conditions.
 - ❑ 13 – Barometric pressure
 - ❑ 13 – Relative humidity
 - ❑ 13 – Solar radiation
 - ❑ 13 – Temperature
 - ❑ 12 – Wind speed/wind direction
 - ❑ 1 – Wind speed vector/wind direction vector
 - ❑ 2 – Precipitation
- Other Monitors
 - ❑ 2 – Total Reactive Oxidized Nitrogen (NO/NO_y)
 - ❑ 2 – Photochemical assessment monitoring station (PAMS). PAMS monitors measure VOCs using automated gas chromatographs (GCs) on an hourly basis during the summer
 - ❑ 7 – Black Carbon

**Number of
Intermittent
Monitors**

Intermittent monitors collect discrete samples for a specific time period. The samples are collected every day, every third day, or every sixth day. The data are averaged in 3-hour or 24-hour intervals.

- Criteria pollutant monitors measure pollutants that have NAAQS.
 - ❑ 5 – PM₁₀ monitors
 - ❑ 14 – PM_{2.5} Federal Reference Method (FRM) monitors
- Non-criteria pollutant monitors measure pollutants that do not have NAAQS.
 - ❑ 2 – PAMS. PAMS monitors measure VOCs on a periodic schedule during the summer months.
 - ❑ 2 – Toxics. These monitors measure health-relevant VOCs.
 - ❑ 2 – Speciation. These monitors measure for PM_{2.5}, nitrates and organics
 - ❑ 1 – PM₁₀ for metals analysis

Section II

Attainment of Air Quality Standards

Attainment Status Summary

The federal Clean Air Act (CAA) contains timeframes and milestones for states to meet and maintain NAAQS for criteria pollutants, which include carbon monoxide, lead, nitrogen dioxide, ozone, particulate matter, and sulfur dioxide. EPA sets NAAQS at levels to protect public health and the environment. The EPA must review each NAAQS every five years and may update the standards based on new scientific information as well as establish new monitoring requirements. Each state is required to monitor the ambient air to determine whether it meets each standard.¹ If monitoring shows that the air quality does not meet a standard, the state must develop and implement pollution control strategies to attain that standard. Once air quality meets a standard, a state must develop a plan to maintain that standard while accounting for future economic and emissions growth. Taken together, these plans and control strategies constitute the State Implementation Plan (SIP).

Carbon Monoxide

Monitored levels of CO in Massachusetts meet the CO standards. Prior to the mid-1980s, Massachusetts was in nonattainment of the CO standards. However, with the adoption of numerous control programs, CO emissions significantly decreased and monitored levels of CO met the standards beginning in 1987. Massachusetts is designated as Unclassifiable/Attainment for the CO standards. Based on EPA's most recent review of the CO standards, in August 2011, EPA retained the existing primary CO standards of 9 ppm measured over 8 hours, and 35 ppm measured over 1 hour. Since EPA did not change the standards, no new designation process was triggered.

Lead

Monitored levels of lead in Massachusetts meet the lead standards. In October 2009, EPA lowered the lead standard from 1.5 µg/m³ to 0.15 µg/m³ averaged over a rolling 3-month period. In November 2011, EPA designated all of Massachusetts as Unclassifiable/Attainment for the 2009 standard. Based on EPA's most recent review of the lead standards, in September 2016, EPA retained the existing lead standards. Since EPA did not change the standards, no new designation process was triggered.

Nitrogen Dioxide

Monitored levels of NO₂ in Massachusetts meet the NO₂ standards. Based on EPA's most recent review of the NO₂ standards, in January 2010, EPA established a new 1-hour NO₂ standard of 100 ppb and new near-road monitoring requirements. In January 2012, EPA designated all of Massachusetts as Unclassifiable/Attainment for the 2010 standard.

¹ MassDEP develops an annual Ambient Air Monitoring Network Plan that describes recent and planned changes to the statewide monitoring network, available at www.mass.gov/eea/agencies/massdep/air/reports/annual-ambient-air-quality-monitoring-network-plan.html.

Sulfur Dioxide

Monitored levels of SO₂ in Massachusetts meet the SO₂ standards. Based on EPA's most recent review of the SO₂ standards, in June 2010, EPA established a new 1-hour SO₂ standard of 75 ppb. In December 2018, EPA designated all of Massachusetts as Unclassifiable/Attainment for the 2010 standard.

Particulate Matter

There are standards for two types of particulate matter: PM₁₀ and PM_{2.5}. Monitored levels of PM₁₀ and PM_{2.5} in Massachusetts meet the respective standards.

Based on EPA's most recent review of the PM_{2.5} standards, in December 2012, EPA lowered the primary annual PM_{2.5} standard to 12 µg/m³. In December 2014, EPA designated all of Massachusetts as Unclassifiable/Attainment for the 2012 standard.

Ozone

In 1979, EPA established an ozone standard (0.12 ppm) based on the maximum 1-hour ozone concentration that occurred each day during the ozone monitoring season. Massachusetts was designated as Nonattainment with this standard.

In 1997, EPA established new 8-hour ozone standards (0.08 ppm) that were designed to be more representative of exposure over time, rather than just the maximum concentration (the 1-hour standard was revoked in 2005). Massachusetts was designated as Nonattainment for these standards at that time. Through a combination of state and regional controls, Massachusetts' air quality attained the 1997 standards by the 2009 attainment deadline.

In 2009, EPA lowered the 8-hour ozone standards to 0.075 ppm. In April 2012, EPA designated Dukes County as Nonattainment for the 2009 ozone standards and designated the remainder of Massachusetts as Unclassifiable/Attainment. Dukes County attained the 2009 ozone standard by the 2015 attainment deadline.

Based on EPA's most recent review of the ozone standards, in October 2015, EPA lowered the 8-hour ozone standards to 0.070 ppm. EPA has designated all of Massachusetts as Unclassifiable/Attainment for the 2015 standard.

2018 Ozone Season

In 2018, there were 12 days when the 8-hour ozone standard of 0.070 ppm was exceeded in Massachusetts. Based on the most recent three years of data (2016–2018), two monitors (Chicopee and Fall River) monitored violations of the 0.070 ppm standard.

While ozone concentrations have trended downward over the past several decades due to air pollution control programs, ozone concentrations vary each year due to varying weather patterns. In general, the chemical reactions that produce elevated ozone concentrations occur when high energy sunlight (present on hot Summer days) facilitates the react of ozone “precursor” pollutants – VOCs and NO_x, which results in ozone formation. Typically, Massachusetts ozone exceedances occur when a high pressure area well south of New England creates a broad southwesterly airflow containing precursors from the upwind coastal urban corridor to New England, where reactions in the atmosphere result in elevated levels of ozone. This typical pattern also moves slowly, promoting heat wave conditions that can last several days, allowing pollutants to build up.

Difference Between Ozone Exceedances and Violations

An ozone exceedance occurs when monitored ozone concentrations exceed the ozone NAAQS. Ozone monitoring data is collected as an hourly average of continuous data which is then used to determine the highest 8-hour average value for the day. An exceedance of the 8-hour standard is an 8-hour averaged value that is greater than 0.070 ppm. An ozone exceedance occurs when a monitor records ambient levels of ozone above the standard. Monitoring an ozone exceedance does not mean that a violation of the ozone standard has occurred, because a violation of an ozone standard (as opposed to an exceedance) is based on three-year averages of data at each monitor.

An ozone violation of the 8-hour standard is determined using the annual 4th-highest daily maximum eight-hour ozone value at each monitor. A violation requires a three-year average of the annual 4th-highest daily maximum eight-hour value that is greater than 0.070 ppm. In other words, the eight-hour values for each day during a year for a specific monitor are ranked from highest to lowest. Then, the 4th-highest value for three consecutive years is averaged. If the three-year average is greater than 0.070 ppm, a violation of the 8-hour standard has occurred at that monitoring site.

Exceedance Days and Total Exceedance Trends

Figure 1 shows the trend in the number of 8-hour exceedance days and the total number of exceedances for the past ten years. Note that years 2009-2015 show what exceedances would have been had the 0.070 ppm 8-hour standards been in effect. Figure 2 shows the most recent ozone design values (i.e., the 4th highest 8-hour O₃ value averaged over three years) relative to the 2015 ozone NAAQS. Monitoring sites with less than three years of data, and therefore no three-year averages, are not included in Figure 2.

Figure 1
8-hour Ozone Exceedance Days and Total Exceedances 2009-2018
Exceedances Based on 0.070 ppm 8-hour Standard





Figure 2
Ozone Design Values (2016-2018)



Daily Ozone and PM Forecasts

MassDEP provides the public with daily air quality forecasts for ozone from April through September and for fine particles all year round using weather maps and meteorological factors to predict whether or not conditions will result in elevated pollution levels. The daily air quality forecasts are available from www.mass.gov/eea/agencies/massdep/air/quality/. EPA web sites that contain regional and national pollution forecasts using data that is provided by participating states are located at www.epa.gov/region01/airquality/forecast.html and <http://airnow.gov/>. The table below describes the ratings used in the daily air quality forecasts.

<u>Air Quality Index (AQI): Ozone</u>			<u>Air Quality Index (AQI): Particle Pollution</u>		
Index Values	Levels of Health Concern	Cautionary Statements	Index Values	Levels of Health Concern	Cautionary Statements
0-50	Good	None	0-50	Good	None
51-100	Moderate	Unusually sensitive people should consider reducing prolonged or heavy exertion outdoors.	51-100	Moderate	Unusually sensitive people should consider reducing prolonged or heavy exertion.
101-150	Unhealthy for Sensitive Groups	Active children and adults, and people with lung disease, such as asthma, should reduce prolonged or heavy exertion outdoors.	101-150	Unhealthy for Sensitive Groups	People with heart or lung disease, older adults, and children should reduce prolonged or heavy exertion.
151-200	Unhealthy	Active children and adults, and people with lung disease, such as asthma, should avoid prolonged or heavy exertion outdoors. Everyone else, especially children, should reduce prolonged or heavy exertion outdoors.	151-200	Unhealthy	People with heart or lung disease, older adults, and children should avoid prolonged or heavy exertion. Everyone else should reduce prolonged or heavy exertion.
201-300	Very Unhealthy	Active children and adults, and people with lung disease, such as asthma, should avoid all outdoor exertion. Everyone else, especially children, should avoid prolonged or heavy exertion outdoors.	201-300	Very Unhealthy	People with heart or lung disease, older adults, and children should avoid all physical activity outdoors. Everyone else should avoid prolonged or heavy exertion.

Section III

Massachusetts Air Quality Data Summaries

Ozone Summary

2018 Ozone Data Summary

A summary of the data collected during the 2018 ozone season (March 1 – September 30) is shown below (in parts per million). MassDEP operated 16 ozone monitors during 2018. The Wampanoag Tribe operated one monitor in Aquinnah on Martha's Vineyard.


CITY	COUNTY	ADDRESS	1ST MAX 8-HR	2ND MAX 8-HR	3RD MAX 8-HR	4TH MAX 8-HR	8-HR MAX>0.070 STD
Aquinnah (Tribal)	Dukes	Herring Creek Drive	0.073	0.073	0.071	0.068	3
Boston	Suffolk	Harrison Avenue	0.073	0.070	0.070	0.067	1
Brockton	Plymouth	Clinton Street	0.084	0.074	0.073	0.072	4
Chelmsford	Middlesex	Technology Drive	0.069	0.067	0.066	0.065	0
Chelmsford	Middlesex	Manning Road	0.051	0.049	0.049	0.049	0
Chicopee	Hampden	Anderson Road	0.078	0.074	0.070	0.070	2
Fairhaven	Bristol	School Street	0.074	0.070	0.069	0.067	1
Fall River	Bristol	Globe Street	0.080	0.080	0.080	0.075	10
Greenfield	Franklin	Barr Avenue	0.070	0.067	0.067	0.066	0
Haverhill	Essex	Washington Street	0.067	0.066	0.066	0.064	0
Lynn	Essex	Parkland Avenue	0.075	0.074	0.072	0.071	4
Milton	Norfolk	Canton Avenue	0.072	0.069	0.069	0.065	1
Pittsfield	Berkshire	Silver Lake Drive	0.059	0.055	0.051	0.051	0
Truro	Barnstable	Collins Road	0.080	0.076	0.071	0.071	4
Uxbridge	Worcester	E. Hartford Ave	0.079	0.074	0.071	0.070	3
Ware	Hampshire	Skyline Drive	0.077	0.071	0.069	0.065	2
Worcester	Worcester	Airport Drive	0.075	0.071	0.070	0.065	2

STANDARDS: 8-hour = 0.070 ppm

ABBREVIATIONS AND SYMBOLS USED IN TABLE

SITE ID = AIRS SITE IDENTIFICATION NUMBER; **1ST, 2ND, 3RD, 4TH MAX 8-HR** = MAXIMUM 8-HOUR VALUE FOR THE 1ST, 2ND, 3RD AND 4TH HIGHEST DAY; **8-HR MAX > .070 STD** = NUMBER OF MEASURED DAILY 8-HOUR MAXIMUM VALUES GREATER THAN THE 0.070 PPM 8-HOUR STANDARD

2018 Ozone Monitoring Network



8-hour Ozone Exceedance Trends

Figure 3 shows the trend for each monitor for the past ten years based on the 0.070 ppm 8-hour standard.

Figure 3
8-hour Ozone Exceedance Trends 2009 – 2018
 Based on the 0.070 ppm 8-hour Standard






Figure 3 cont.
8-hour Ozone Exceedance Trends 2009 – 2018
 Based on the 0.070 ppm 8-hour Standard



Sulfur Dioxide (SO₂) Summary

2018 SO₂ Data Summary


A summary of the 2018 SO₂ data is shown below (in parts per billion). MassDEP operated seven SO₂ monitors during 2018.

CITY	COUNTY	ADDRESS	1ST MAX 1-HR	2ND MAX 1-HR	99TH PCTL 1-HR	ARITH MEAN	1ST MAX 24-HR	2ND MAX 24-HR
Boston	Suffolk	Kenmore Square	4.2	3.6	3.3	0.49	2.2	2.0
Boston	Suffolk	Harrison Avenue	6.5	5.1	3.8	0.53	2.5	2.2
Fall River	Bristol	Globe Street	4.9	4.7	3.8	0.60	1.6	1.5
Springfield	Hampden	Liberty Street (1)	7.8	5.8	7.8	1.16	2.7	2.7
Springfield	Hampden	Liberty Street (2)	5.2	3.8	3.3	0.38	2.3	1.4
Ware	Hampshire	Skyline Drive	6.6	4.8	3.8	0.34	3.2	2.3
Worcester	Worcester	Summer Street	5.8	5.0	3.9	0.56	2.5	2.2

STANDARDS: 1-hour = 75 ppb 3-hour = 0.5 ppm (500 ppb)

ABBREVIATIONS AND SYMBOLS USED IN TABLE


1ST, 2ND MAX 1-HR = FIRST AND SECOND HIGHEST 1-HOUR VALUE; **99TH PCTL 1-HR** = 99th PERCENTILE OF THE 1-HOUR MAXIMUM VALUE;
ARITH MEAN = ANNUAL ARITHMETIC MEAN; **1ST, 2ND MAX 24-HR** = FIRST AND SECOND HIGHEST 24-HOUR VALUE



SO₂ Trends

Figure 4 shows the trend of the 1-hour 99th percentile for each SO₂ monitor for the past 10 years relative to the 1-hour standard of 75 ppb.

Figure 4
Sulfur Dioxide Trends 2009 – 2018
1-hour 99th Percentile Annual Average



Nitrogen Dioxide (NO₂) Summary

2018 NO₂ Data Summary


A summary of the 2018 NO₂ data is shown below (in parts per billion). MassDEP operated 11 NO₂ monitors during 2018.

CITY	COUNTY	ADDRESS	1ST MAX 1-HR	2ND MAX 1-HR	98TH PERCENTILE VALUE	ARITH MEAN
Boston	Suffolk	Kenmore Square	54	53	45	13.13
Boston	Suffolk	Harrison Avenue	54	53	48	11.19
Boston	Suffolk	Von Hillern Street	53	50	44	13.45
Chelmsford	Middlesex	Manning Road	36	34	33	9.67
Chicopee	Hampden	Anderson Road	42	41	37	5.24
Lynn	Essex	Parkland Avenue	39	37	34	4.36
Milton	Norfolk	Canton Avenue	77	42	32	4.09
Springfield	Hampden	Liberty Street (1)	52	45	43	12.98
Springfield	Hampden	Liberty Street (2)	37	35	31	8.57
Ware	Hampshire	Skyline Drive	30	26	23	2.27
Worcester	Worcester	Summer Street	47	45	42	11.44

STANDARDS: Annual Arithmetic Mean = 53 ppb 1-hour = 100 ppb

ABBREVIATIONS AND SYMBOLS USED IN TABLE


SITE ID = AIRS SITE IDENTIFICATION NUMBER; **1ST, 2ND MAX 1-HR** = FIRST AND SECOND HIGHEST 1-HOUR VALUE; **ARITH MEAN** = ANNUAL ARITHMETIC MEAN



NO₂ Trends

Figure 5 shows the trend of the 1-hour 98th percentile annual average for each NO₂ monitor over the past 10 years relative to the 1-hour standard of 100 ppb.

Figure 5
Nitrogen Dioxide Trends 2009 - 2018
 1-hour 98th Percentile Annual Average



Carbon Monoxide (CO) Summary

2018 CO Data Summary


A summary of the 2018 CO data is shown below (in parts per million). MassDEP operated five CO monitors during 2018.

CITY	COUNTY	ADDRESS	1ST MAX 1-HR	2ND MAX 1-HR	1ST MAX 8-HR	2ND MAX 8-HR
Boston	Suffolk	Harrison Avenue	1.315	1.105	0.9	0.7
Boston	Suffolk	Von Hillern Street	1.105	1.092	0.8	0.7
Springfield	Hampden	Liberty Street (1)	1.400	1.400	1.0	0.9
Springfield	Hampden	Liberty Street (2)	1.007	0.973	0.9	0.7
Worcester	Worcester	Summer Street	1.432	1.367	0.9	0.8

STANDARDS: 1-hour = 35 ppm 8-hour = 9 ppm

ABBREVIATIONS AND SYMBOLS USED IN TABLE


SITE ID = AIRS SITE IDENTIFICATION NUMBER; **1ST, 2ND MAX 1-HR** = FIRST AND SECOND HIGHEST 1-HOUR VALUE; **1ST, 2ND MAX 8-HR** = FIRST AND SECOND HIGHEST 8-HOUR VALUE



CO Trends

Figure 6 shows the trend of the 2nd maximum 8-hour average for each CO monitor over the past 10 years relative to the 8-hour standard of 9 ppm.

Figure 6
Carbon Monoxide Trends 2009-2018
2nd Maximum 8-hour Values



Particulate Matter 10 Microns (PM₁₀) Summary

2018 PM₁₀ Data Summary


A summary of the 2018 PM₁₀ data is shown below (in $\mu\text{g}/\text{m}^3$). MassDEP operated five PM₁₀ monitors in 2018.

CITY	COUNTY	ADDRESS	1ST MAX 24-HR	2ND MAX 24-HR	3RD MAX 24-HR	4TH MAX 24-HR	DAYS MAX >STD	ARITH MEAN
Boston	Suffolk	Harrison Avenue	24	23	22	20	0	9.0
Boston	Suffolk	Harrison Avenue (collocated)	24	23	22	19	0	9.4
Springfield	Hampden	Liberty Street (1)	27	21	20	19	0	8.9
Ware	Hampshire	Skyline Drive	17	16	13	11	0	5.1
Worcester	Worcester	Summer Street	46	32	30	25	0	13.4

STANDARD: 24-hour = 150 $\mu\text{g}/\text{m}^3$

ABBREVIATIONS AND SYMBOLS USED IN TABLE


SITE ID = AIRS SITE IDENTIFICATION NUMBER; **COLLOC** = COLLOCATED; **1ST, 2ND, 3RD, 4TH 24-HR MAX** = 1ST, 2ND, 3RD, AND 4TH HIGHEST 24-HOUR VALUES FOR THE YEAR; **ARITH MEAN** = ANNUAL ARITHMETIC MEAN



PM₁₀ Trends

Figure 7 shows the 2018 calendar year 24-hour maximum concentration for each PM₁₀ monitor relative to the 24-hour standard of 150 µg/m³.

Figure 7
PM₁₀ Trends 2009-2018
24-Hour Calendar Year Maximum



Particulate Matter 2.5 Microns (PM_{2.5}) Summary

During 2018, MassDEP operated 14 Federal Reference Method (FRM) filter-based PM_{2.5} monitors and 18 Federal Equivalence Method (FEM) continuous PM_{2.5} monitors.

FRM monitors require the manual set-up and collection of filters that measure 24-hour samples every three or six days. The filters are weighed prior to placement in the field and then weighed again after the sample is collected to determine the amount of PM_{2.5} collected on the filter.

FEM monitors measure PM_{2.5} on an hourly basis. MassDEP operates FRM and FEM monitors side-by-side at some locations for comparison. In general, FEM monitors provide slightly higher values than FRM monitors, which may be due to measurement of fresh hourly PM_{2.5} samples with FEM versus slightly aged 24-hour samples with the FRM filter method.

2018 PM_{2.5} FRM Data Summary

A summary of the 2018 PM_{2.5} FRM data is shown below (in µg/m³).


CITY	COUNTY	ADDRESS	1ST MAX 24-HR	2ND MAX 24-HR	3RD MAX 24-HR	4TH MAX 24-HR	98 TH PERCENTILE 24-HR	ARITH MEAN
Boston	Suffolk	Kenmore Square	17.5	14.4	13.3	13.1	14.4	5.69
Boston	Suffolk	Harrison Avenue	21.5	14.3	13.4	12.1	14.3	5.31
Boston	Suffolk	Von Hillern Street	16.5	13.5	13.0	12.3	13.5	6.21
Boston	Suffolk	North Street	15.9	12.3	9.7	9.5	15.9	5.60
Boston	Suffolk	North Street (collocated)	9.7	9.6	8.6	8.1	9.7	5.97
Brockton	Plymouth	Clinton Street	13.3	12.1	10.1	10.0	12.1	4.89
Chicopee	Hampden	Anderson Road	15.9	14.4	13.4	12.9	14.4	5.17
Chicopee	Hampden	Anderson Road (collocated)	18.2	14.4	13.5	13.4	14.4	5.38
Greenfield	Franklin	Barr Avenue	15.2	14.8	12.7	11.5	14.8	5.18
Haverhill	Essex	Washington Street	13.3	12.4	12.4	11.8	12.4	5.03
Pittsfield	Berkshire	Silver Lake Drive	44.0	17.5	16.0	15.3	17.5	5.93
Springfield	Hampden	Liberty Street (1)	15.4	9.9	9.8	9.7	15.4	4.53
Springfield	Hampden	Liberty Street (2)	16.2	15.0	14.5	13.4	15.0	6.67
Worcester	Worcester	Summer Street	15.0	14.4	12.9	12.8	14.4	5.58

STANDARDS: Annual Mean = 12.0 µg/m³ (primary) 24-hour (98th percentile) = 35 µg/m³

ABBREVIATIONS AND SYMBOLS USED IN TABLE

SITE ID = AIRS SITE IDENTIFICATION; **COLLOC** = COLLOCATED; **1ST, 2ND, 3RD, 4TH 24-HR MAX** = 1ST, 2ND, 3RD, AND 4TH HIGHEST 24-HOUR VALUES FOR THE YEAR; **98TH PERCENTILE 24-HR** = 98TH PERCENTILE VALUE FOR THE YEAR; **ARITH MEAN** = ANNUAL ARITHMETIC MEAN


2018 PM_{2.5} Monitoring Network



PM_{2.5} FRM Trends

Figure 8 shows trends of the annual arithmetic mean for each PM_{2.5} FRM monitor over the past 10 years relative to the standard of 12 $\mu\text{g}/\text{m}^3$.

Figure 8
PM_{2.5} Trends 2009-2018
FRM Annual Arithmetic Mean



2018 PM_{2.5} FEM Data Summary

A summary of the 2018 PM_{2.5} FEM data is shown below (in µg/m³).

CITY	COUNTY	ADDRESS	1ST MAX	2ND MAX	3RD MAX	4TH MAX	98TH PERCENTILE 24-HR	ARITH MEAN
Boston	Suffolk	Harrison Avenue	29.1	24.5	23.7	22.6	20.9	8.77
Boston	Suffolk	Von Hillern Street	26.2	24.9	23.4	23.0	21.0	10.11
Boston	Suffolk	Von Hillern Street (collocated)	27.0	22.5	21.8	21.3	23.0	8.78
Boston	Suffolk	North Street	22.0	19.4	17.8	17.4	19.4	7.57
Brockton	Plymouth	Clinton Street	27.3	24.2	22.3	20.6	17.5	6.03
Chelmsford	Middlesex	Manning Road	17.5	17.5	16.8	16.5	15.9	7.51
Chicopee	Hampden	Anderson Road	14.4	11.5	10.4	9.4	11.5	3.45
Fall River	Bristol	Globe Street	20.5	18.5	18.2	17.0	15.1	6.81
Greenfield	Franklin	Barr Avenue	24.9	20.7	20.3	19.5	17.5	5.78
Haverhill	Essex	Washington Street	19.2	15.9	15.8	15.1	13.1	4.91
Lynn	Essex	Parkland Avenue	23.9	23.5	21.6	21.6	19.7	6.33
North Adams	Berkshire	Holden Street	22.7	19.0	17.6	17.5	16.0	5.85
Pittsfield	Berkshire	Center Street	22.4	20.0	19.9	18.3	19.9	5.06
Pittsfield	Berkshire	Silver Lake Drive	19.3	18.9	18.8	18.1	18.1	7.63
Springfield	Hampden	Liberty Street (1)	22.7	21.9	21.8	21.7	12.8	7.66
Springfield	Hampden	Liberty Street (2)	18.4	18.1	17.9	16.7	16.2	6.95
Ware	Hampshire	Skyline Drive	22.9	21.6	19.9	18.3	16.4	5.71
Worcester	Worcester	Summer Street	22.6	22.6	21.6	20.5	17.1	6.51

STANDARDS: Annual Mean = 12.0 µg/m³ (primary) 24-hour (98th percentile) = 35 µg/m³


ABBREVIATIONS AND SYMBOLS USED IN TABLE

SITE ID = AIRS SITE IDENTIFICATION; **1ST, 2ND, 3RD, 4TH MAX** = 1ST, 2ND, 3RD, AND 4TH HIGHEST 24-HOUR VALUES FOR THE YEAR;
98TH PERCENTILE 24-HR = 98TH PERCENTILE VALUE FOR THE YEAR; **ARITH MEAN** = ANNUAL ARITHMETIC MEAN

PM_{2.5} FEM Trends

Figure 9 shows trends of the annual arithmetic mean for each PM_{2.5} FEM monitor over the past 10 years relative to the standard of 12 $\mu\text{g}/\text{m}^3$.

Figure 9
PM_{2.5} Trends 2009-2018
FEM Annual Arithmetic Mean



Speciation

MassDEP collects PM_{2.5} samples for speciation in Boston (Harrison Avenue) and Chicopee. Speciation involves analysis of particulate matter to determine its chemical composition and to identify air pollution sources that affect the area around the monitoring station. Pollutants analyzed include elements (e.g., metals), sulfates, nitrates, and carbon (total and organic).

IMPROVE (Interagency Monitoring of Protected Visual Environments)

IMPROVE is a nationwide program designed to assess air quality at rural locations where air pollution may affect visibility over long distances (e.g., mountain ranges or scenic vistas). During 2018, the National Park Service operated an IMPROVE sampler at the Truro monitoring site, and the Wampanoag Tribe operated an IMPROVE sampler at its Martha's Vineyard monitoring site. These samplers acquire PM_{2.5} filter samples for speciation analysis to determine effects on visibility. Data can be viewed at the IMPROVE web site at <http://vista.cira.colostate.edu/improve/Data/data.htm>.

Quality Control and Quality Assurance

In order to ensure that all air quality data is of acceptable and consistent quality, MassDEP has developed standard operating procedures (SOPs) based on federal requirements that include quality control and quality assurance protocols that systematically assess the entire sample collection and data handling system on an ongoing basis. Ambient air monitoring quality assurance requirements are contained in the Federal Regulations at 40 CFR Part 58, Appendix A – E. Each year MassDEP certifies that it is in compliance with the federal requirements. A few of the considerations that affect sample collection data quality are:

- Site Placement
- Intake Probe Material
- Intake Probe Height
- Spacing from roadways and trees

For data processing there are quantitative statistics and qualitative descriptors used to interpret the degree of acceptability and utility of data. Examples of these data quality indicators are:

- Representativeness
- Precision
- Bias
- Detectability
- Completeness
- Comparability

MassDEP's Air Assessment Branch maintains a Quality Assurance Group that ensures samples are collected correctly and conducts performance audits throughout the air monitoring network to verify data validity. There also is a Quality Control Group that reviews daily monitored and historical data for validity, tracks precision results, finalizes monthly values, and submits air quality data to EPA's database in a timely manner. Computer software tools, report queries, and "eyes on" data reviews all are used to validate data before it is submitted to EPA. EPA also conducts its own performance audits on MassDEP samplers and every three years conducts a thorough Technical Systems Audit (TSA). The latest TSA was conducted in 2016.

Section IV

PAMS/Air Toxics Monitoring

PAMS Monitoring

Ground-level ozone is a secondary pollutant and is not discharged directly to the atmosphere from a stack or tailpipe, but forms in the atmosphere from the photochemical reactions of other pollutants such as VOCs and NO_x. Ozone formation can occur many miles downwind from the source of the original emissions. These reactions occur in the presence of strong sunlight and are most pronounced during the hottest days of the summer. The PAMS program was established by the 1990 Clean Air Act Amendments as a way to collect data for assessing NAAQS attainment progress independent of the meteorological variation that occurs between years and for identifying appropriate pollution control strategies. In 2018 MassDEP operated PAMS stations in Lynn and Chicopee.

PAMS is a special designation for enhanced monitoring stations that are designed to gather information on the ozone formation process. Instruments at these sites measure pollutants and meteorological parameters that are specific to the photochemical processes by which ozone is created in the atmosphere at ground level. In addition to the pertinent NAAQS pollutants (ozone, NO₂, etc.), non-criteria pollutants, including VOCs, are measured at PAMS stations on either an hourly basis or at regular intervals during the hottest part of the summer (June, July and August). Meteorology is a critical component of ozone formation and each PAMS site has a full complement of meteorological sensors including wind speed, wind direction, temperature, relative humidity, barometric pressure, solar radiation and at some sites, total ultraviolet light and precipitation.


Air Toxics Monitoring

Toxic air pollutants are known or suspected to cause cancer or other serious health effects. Air toxics include certain VOCs and toxic metals (e.g., arsenic, cadmium).

The Boston (Harrison Avenue) monitoring site is designated as a National Air Toxics Trends Station (NATTS) designed to collect and quantify a number of toxic air pollutants, including VOCs, metals, carbonyls, black carbon and polycyclic aromatic hydrocarbons (PAHs). Data from this site is compared with data from a network of similar sites across the country to identify transport, trends and site-specific characteristics of these pollutants.

MassDEP also monitors VOCs as part of the PAMS monitoring program, some of which are classified as air toxics. Figure 10 summarizes concentrations of benzene, toluene, ethyl benzene, xylenes measured at the Lynn PAMS site for the past 10 years. Allowable Ambient Limit (AAL) values are presented for reference. AALs are health-based air toxics guidelines developed by MassDEP based on known or suspected carcinogenic and toxic health properties of individual compounds. AAL concentrations were developed for a 70-year lifetime exposure, but are used for comparison with annual averages.

Figure 10
Lynn Toxics VOC Summary 2009– 2018
24-hour samples



The table below summarizes 24-hour concentrations of target VOCs measured at the Boston (Harrison Ave) and Lynn sites for 2018. Harrison Avenue serves as the central city sampling location and Lynn serves as the area background site.

Compounds	Boston (Harrison Avenue)		Lynn	
	Max Value ppb	Mean ppb	Max Value ppb	Mean ppb
1,3-butadiene	0.050	0.016	0.028	0.009
1,1,1-trichloroethylene	0.000	0.000	0.000	0.000
tetrachloroethylene	0.050	0.000	0.050	0.000
trichloroethylene	0.000	0.000	0.000	0.000
benzene	0.300	0.130	0.267	0.098
toluene	0.957	0.214	0.414	0.133
ethylbenzene	0.075	0.030	0.050	0.019
xylene	0.300	0.115	0.188	0.071

Samples collected at the Harrison Avenue site are analyzed for a suite of metals that are known to be toxic in the environment. The table below summarizes the 2018 metals data.

Metal	Boston (Harrison Avenue)	
	Max Value ug/m3	Mean ug/m3
Antimony	0.00350	0.00154
Arsenic	0.00106	0.00036
Beryllium	0.00002	0.00001
Cadmium	0.00029	0.00007
Chromium	0.00630	0.00422
Cobalt	0.00024	0.00008
Lead	0.00670	0.00183
Manganese	0.01260	0.00409
Mercury	0.00003	0.00001
Nickel	0.00456	0.00092
Selenium	0.00077	0.00054

Appendix A 2018 Monitoring Stations

CITY/TOWN	ADDRESS	PARAMETERS MONITORED
AQUINNAH*	HERRING CREEK RD	O3, IMPROVE
BOSTON	KENMORE SQUARE	NO2, NO, NOx, SO2, PM2.5
BOSTON	1159 HARRISON AVENUE	O3, NO2, NO, NOx, NOy, SO2, CO, PM2.5, PM10, PM Coarse, PM2.5 Speciation, Black Carbon, Toxics, Carbonyls, WS/WD, WSv/WDv, TEMP, SUN, REL, BP
BOSTON	174 NORTH ST	PM2.5, Black Carbon
BOSTON	19 VON HILLERN ST	NO2, NO, NOx, CO, PM2.5, Black Carbon, WS/WD, TEMP, SUN, REL, BP
BROCKTON	170 CLINTON ST	O3, PM2.5
CHELMSFORD	11 TECHNOLOGY DR	O3
CHELMSFORD	5 MANNING RD	NO2, NO, NOx, PM2.5, O3, Black Carbon
CHICOPEE	ANDERSON RD	O3, NO2, NO, NOx, PM2.5, PM2.5 speciation, WS/WD, TEMP, SUN, REL, BP
FAIRHAVEN	30 SCHOOL ST	O3, WS/WD, TEMP, SUN, REL, BP
FALL RIVER	659 GLOBE ST	O3, SO2, PM2.5
GREENFIELD	16 BARR AVE	O3, PM2.5, Black Carbon, WS/WD, TEMP, SUN, REL, BP
HAVERHILL	685 WASHINGTON ST	O3, PM2.5, WS/WD, TEMP, SUN, REL, BP
LYNN	390 PARKLAND AVE	O3, NO2, NO, NOx, PM2.5, VOCs, Carbonyls, WS/WD, TEMP, SUN, REL, BP, PRECIP
MILTON	1904 CANTON AVENUE	O3, NO2, NO, NOx, WS/WD, TEMP, SUN, REL, BP
NORTH ADAMS	86 HOLDEN STREET	PM2.5, Black Carbon
PITTSFIELD	78 CENTER STREET	PM2.5
PITTSFIELD	25 SILVER LAKE DR	O3, PM2.5, Black Carbon
SPRINGFIELD (1)	165 LIBERTY STREET	NO2, NO, NOx, SO2, CO, PM2.5, PM10, Black Carbon
SPRINGFIELD (2)	600 LIBERTY STREET	NO2, NO, NOx, SO2, CO, PM2.5, Black Carbon
TRURO	6 COLLINS ROAD	O3, WS/WD, TEMP, SUN, REL, BP, IMPROVE
UXBRIDGE	366 E. HARTFORD AVE	O3, WS/WD, TEMP, SUN, REL, BP
WARE	36 SKYLINE DRIVE	O3, NO2, NO, NOx, NOy, SO2, PM2.5, PM10, WS/WD, TEMP, SUN, REL, BP, PRECIP
WORCESTER	375 AIRPORT DRIVE	O3, WS/WD, TEMP, SUN, REL, BP
WORCESTER	SUMMER STREET	NO2, NO, NOx, SO2, CO, PM2.5, PM10

* Wampanoag Tribal Site