

Environmental Monitoring Report

For 2018

Pilgrim and Seabrook
Nuclear Power Station
Emergency Planning Zones

Prepared by:

Environmental Toxicology Program
Bureau of Environmental Health
Massachusetts Department of Public Health
250 Washington Street
Boston, Massachusetts 02108

March 2020

EXECUTIVE SUMMARY	1
1. INTRODUCTION	2
A. ENVIRONMENTAL RADIATION	3
B. LABORATORY METHODS.....	4
2. ENVIRONMENTAL MONITORING AND SAMPLING.....	6
A. PILGRIM NUCLEAR POWER STATION	6
B. SEABROOK NUCLEAR POWER STATION	9
3. 2018 ENVIRONMENTAL MONITORING RESULTS	11
A. PILGRIM NUCLEAR POWER STATION	11
B. SEABROOK NUCLEAR POWER STATION	12
D. REFERENCES	14
E. RESULTS TABLES.....	15

EXECUTIVE SUMMARY

The Massachusetts Department of Public Health (MDPH) Bureau of Environmental Health's (MDPH/BEH) Environmental Toxicology and Radiation Control Programs collaborate to conduct routine environmental monitoring in nuclear power station Emergency Planning Zones (EPZs) within the Commonwealth. This monitoring is part of the Department's regulatory responsibility, and provides a system of watchfulness over environmental radiation in Massachusetts communities surrounding nuclear power plants. These EPZs include communities located within a 10-mile radius of Pilgrim Nuclear Power Station (Pilgrim) in Plymouth, MA and Seabrook Nuclear Power Station (Seabrook) in Seabrook, NH. The Vermont Yankee Nuclear Power Station (VY), in Vernon, VT, ceased operations on December 29, 2014 and is undergoing decommissioning. Routine monitoring within the VY 10-mile EPZ was discontinued in 2017 after the US Nuclear Regulatory Commission (NRC) reduced the EPZ to the perimeter of the plant. This report summarizes the 2018 monitoring activities and results for the Pilgrim and Seabrook nuclear plant EPZs¹.

Report Highlights

- Overall, no radiation indicators or radionuclides were detected at a level of health concern.
- Radiation monitoring results in 2018 for areas surrounding the two nuclear power stations affecting Massachusetts have been either non-detect, naturally occurring, at levels expected to be present in the environment from background fallout from historic bomb testing and past nuclear accidents, or attributable to a known source.
- Iodine-131 was detected in Irish moss samples both within and outside the Pilgrim EPZ and at a background location outside the Seabrook EPZ, all at levels not of health concern.
 - At Pilgrim, Iodine-131 was detected at just above the laboratory reporting levels. Iodine-131 has previously been detected at the same locations, similarly at levels just above the laboratory reporting levels. Iodine-131 was not detected in any of the Pilgrim air or surface water samples, and MDPH determined the source is not likely to be the nuclear power plant.
 - Iodine-131 has also been detected previously at the background location outside the Seabrook EPZ. Based on the location of this sample, the Iodine-131 is most likely from either a water treatment plant or medical facility located nearby, where Iodine-131 may be used for treating certain thyroid disorders.

¹Pilgrim permanently shut down the nuclear power reactors on May 31, 2019. The site entered Phase I of plant decommissioning on June 11, 2019 when the nuclear fuel from the reactor was safely transferred into the spent fuel pool, and the site was certified as having permanent cessation of operations and permanent removal of fuel by Federal regulators. In August of 2019 Pilgrim was sold by the Entergy Corporation to Holtec International for completion of the remaining decommissioning steps.

1. INTRODUCTION

The MDPH/BEH radiation environmental monitoring program is designed to monitor radiation levels and to protect residents in the Commonwealth from exposure to radiation. Samples of environmental media collected within and just outside the EPZs surrounding nuclear power plants by MDPH/BEH, or provided by the utilities that operate the nuclear power plants, are analyzed for radiation by the MDPH/BEH Massachusetts Environmental Radiation Laboratory (MERL). Environmental media analyzed in 2018 include: air, surface water, milk, fish, shellfish, sediment, vegetation and food crops. In addition to the samples analyzed for radiation by MERL, MDPH/BEH has a network of stationary monitors surrounding Pilgrim that measures gamma radiation in real-time, which is monitored online by MDPH/BEH staff. The C-IO Research & Education Foundation, Inc., a non-profit organization under contract to MDPH/BEH, conducts direct radiation monitoring in Massachusetts communities within the Seabrook EPZ and provides summary reports to MDPH/BEH.

The radiation environmental monitoring of Pilgrim and Seabrook EPZs has been in place since the 1980s. A focused investigation of tritium in groundwater on the Pilgrim Nuclear Power Plant property is ongoing and not part of this report. Updates on this monitoring effort are posted on the MDPH website: [Tritium investigation update reports](#). The environmental monitoring program for Massachusetts communities within the VY EPZ began in 2011, continuing through 2017. The VY reactor shut down permanently on December 29, 2014. Routine monitoring was discontinued in 2017 after the U.S. Nuclear Regulatory Commission (NRC) reduced the EPZ to the perimeter of the plant.

The NRC requires specific environmental monitoring and annual reporting by operating nuclear power plants. The NRC reports summarizing Seabrook's and Pilgrim's environmental monitoring can be found on its website: [Pilgrim's 2018 Environmental Radiological Monitoring Report](#) and [Seabrook's 2018 Environmental Radiological Monitoring Report](#).

This report contains background information regarding environmental radiation and laboratory methods used to analyze samples for radiation; sample location and analyses for the Pilgrim and Seabrook EPZs; and a summary of the monitoring results.

A. ENVIRONMENTAL RADIATION

Background radiation in the environment comes from three general sources: naturally occurring radiation, radioactive fallout from past weapons testing or nuclear accidents, and man-made sources.

Naturally occurring radionuclides, such as Potassium-40 and Beryllium-7, are present in most environmental media. Potassium-40 is a radioactive form of potassium, which is an essential nutrient. Beryllium-7 is produced when cosmic energy collides with nitrogen and oxygen in the atmosphere (Delaygue et al., 2015). Additional natural sources of radiation, including cosmic radiation, radon, and carbon-14, contribute to an annual background radiation dose of approximately 310 mrems/year (US NRC, 2017a). Man-made sources include medical procedures (e.g., diagnostic x-rays) and various consumer products (e.g., certain construction material, combustible fuels, televisions, smoke detectors) (US NRC 2017b). Background and man-made sources contribute to the estimated 620 mrem annual dose of environmental radiation for average U.S. residents (US NRC, 2017c).

Table 1. Background Radiation Dose for Average U.S. Resident

Source	Millirems/year
Natural background radiation	310
Man-made sources	310
Total of all sources	620

Source: US NRC, 2017c

Background radiation includes fallout radiation from historical weapons testing, which occurred primarily in the 1950s and 1960s, and from nuclear power plant accidents such as Chernobyl and Fukushima. This fallout includes radioisotopes such as Cesium-137 (Cs-137) and Strontium-90 (Sr-90), which persist in the environment due to their 28-30 year half-lives.

During active operation, nuclear power plants emit direct gamma radiation from nuclear reactor systems; noble gases, tritium, Iodine-131, Carbon-14, and particulates from the station's air stack; and discharge water containing tritium as well as other radionuclides that emit alpha, beta and gamma radiation (Luykx and Fraser, 1983; UNSCEAR, 2008). Noble gases are chemically inert, have short half-lives, disperse quickly in the environment, and do not bioconcentrate or easily incorporate into biological tissue. Tritium is created when water passes through the reactor core; the hydrogen atoms in the water molecules and other trace elements like boron absorb neutrons from the fission of the reactor fuel. Tritium is lighter and more mobile in water than other radionuclides and is a sentinel indicator of radionuclides in water bodies. Both Iodine-131 and particulates (notably Cesium-137, Cobalt-60, Iron-59, Magnesium-54, Strontium-90 and Zinc-65) have environmental and public health significance: their half-lives range from weeks to years, they are readily incorporated into biological tissue, and they will bioconcentrate. Iodine-131 is usually the first radioactive element detected in the event of an accidental release of power plant radiation (ATSDR, 2002). Carbon-14 is a naturally occurring radionuclide, which can also be released in relatively small amounts from nuclear power plants, primarily due to its formation in the coolant system (Yim and Caron, 2006).

Exposure to radiation from nuclear power plants may occur from permitted air or liquid discharges or from unmonitored releases or leaks. MDPH/BEH evaluates possible routes of exposure for radionuclides, particularly those that accumulate in the food chain, and samples environmental media along these routes to measure potential exposure to radiation.

B. LABORATORY METHODS

The MDPH/BEH Radiation Control Program's Massachusetts Environmental Radiation Laboratory (MERL) analyzes samples for a suite of more than 30 radioactive isotopes (e.g., radioisotopes, or radionuclides). Gamma spectroscopy is used to identify and detect environmentally significant and naturally-occurring radioisotopes; gas proportion counters measure gross beta and alpha radiation; and liquid scintillation counters measure tritium. Environmental media sample results are compared to typical background levels. In the event that gamma emitters are present above typical background, the MERL protocol calls for additional testing at an outside laboratory for alpha emitters, such as transuranic (high atomic number) elements, and beta emitters, such as Strontium-90. MERL maintains its standard of excellence in analytical capability through participation with several federal agencies in inter-laboratory quality assurance measures.

Analysis methods by media are summarized below:

AIR

Air filters are collected weekly and analyzed for gross alpha and gross beta radioactivity using a gas proportion counter. Gross alpha and beta analysis is a screening-level tool that does not identify individual radionuclides; therefore, air filters are also analyzed quarterly for gamma emitting radionuclides using gamma spectroscopy. Results are compared to results from a background monitor located in Boston.

Air cartridges are analyzed weekly for iodine-131 using gamma spectroscopy.

Direct gamma radiation in air is measured with thermoluminescent dosimeters (TLDs) and analyzed using gamma spectroscopy.

WATER

Surface water samples are tested for total alpha and beta radioactivity with a gas proportional counter, and for gamma-emitting radionuclides with a gamma spectrometer. Water samples are also tested for tritium with a liquid scintillation counter.

MILK

Milk is a good indicator media for radioactive elements, particularly iodine-131, which can be detected in milk soon after cows graze on contaminated pastures or feed. Hence, cow's milk is tested for gamma radionuclides, including iodine-131, using gamma spectroscopy.

SEAFOOD, SEDIMENT, VEGETATION, AND FOOD CROPS

Seafood, sediment, vegetation and food crops were chosen to represent various stages of the food chain where radionuclides may be identified. Mollusks (such as clams and mussels) filter-feed sediment and sand where heavy and sediment-bound radionuclides may accumulate; lobsters eat clams, mussels and small fish; and radionuclides biomagnify from smaller to larger surface-dwelling fish.

Analyses of vegetation and crop samples aim to identify radionuclides that may settle on surfaces or be absorbed through the roots. Samples are tested for gamma-emitting radionuclides using a gamma spectrometer.

QUALITY ASSURANCE

Laboratory sample detection levels are affected by sample size, time between collection and analysis, and equipment processing and counting time. Where detection levels fall outside our analytic sensitivity guidelines they are noted in the tables as “NR” (Result is not reported for quality control reason).

2. ENVIRONMENTAL MONITORING AND SAMPLING

This section describes the two nuclear power station EPZs in Massachusetts and summarizes the environmental samples collected and analyzed in 2018.

A. PILGRIM NUCLEAR POWER STATION

The Pilgrim Nuclear Power Station (Pilgrim) is located in Plymouth, MA. Five Massachusetts communities are included in the 10-mile EPZ of Pilgrim: Carver, Duxbury, Kingston, Marshfield, and Plymouth, all shown below in Figure 1.




Figure 1. The MDPH/BEH Radiation -monitoring network at Pilgrim

MDPH/BEH's radiation monitoring conducted within and outside the Pilgrim EPZ is a combination of independent direct monitoring of airborne radiation; air, milk and cranberry sampling; and analysis of split samples provided by Entergy, the utility that owned Pilgrim in 2018, of water, fish, shellfish, sediment, Irish moss, and crops. Figure 1 shows locations of the air monitors for Pilgrim. Sample locations for water, fish and shellfish, sediment, Irish moss and crops are shown in Figure 2.




Figure 2. Pilgrim EPZ and sampling locations

Air/Direct Radiation

MDPH/BEH's direct radiation monitoring at Pilgrim is comprised of three systems operating on real-time, weekly, and quarterly bases. The redundant systems are designed to independently monitor the land areas within the 10-mile EPZ and to verify the utility's radiation monitoring.

MDPH/BEH maintains a network of 15 stationary radiation monitoring stations that detects gamma radiation in real-time and transmits data to a computer which is remotely accessed by staff. Emergency alerts are sent to MDPH and Massachusetts Emergency Management Agency (MEMA) officials if radiation is detected at levels greater than three times the typical background level. In 2016, MDPH/BEH completely replaced the older system with new monitors and servers, and installed an internet-based communication system.

MDPH/BEH analyzes samples collected from an air particulate filter and a charcoal air cartridge located just outside the Pilgrim utility's fence. Filters are analyzed for gross beta and gross alpha radioactivity and cartridges are analyzed for iodine-131. A filter composite sample is also analyzed quarterly for additional gamma-emitting radionuclides. The same analyses are done for an air particulate filter and charcoal cartridge collected from a background location in Boston.

MDPH/BEH also has a network of 39 TLDs placed throughout the Pilgrim EPZ and surrounding communities, which measure total gamma radiation in milliroentgen (mR). The majority of the TLDs are located in the inner region of the EPZ, and three are near the plant border. These TLDs are collected and analyzed quarterly, and the results are compared to those of a background location in Boston.

Surface Water

Entergy collects seawater on a monthly basis from the Pilgrim discharge canal and the Powder Point Bridge in Duxbury and provides split samples to MDPH/BEH for analysis of gamma-emitting radionuclides. MERL also analyzes monthly composites of weekly surface water samples from both locations for tritium.

Milk

MDPH/BEH collects samples of cow's milk monthly from a farm in Duxbury. The milk is analyzed for gamma-emitting radionuclides, including Iodine-131. Although this farm is located just outside the EPZ (11 miles from Pilgrim), it is the closest dairy farm to Pilgrim with available milk samples.

Fish and Shellfish

Entergy provides annual split samples of fish and lobster collected from the Pilgrim discharge canal to MERL for analysis. Entergy also provides MERL with split samples of background fish and lobster from Cape Cod Bay.

Entergy collects *Mytilus* (i.e., blue or common mussels) semiannually from Green Harbor in Marshfield and collects soft shell clams semiannually from Duxbury Bay and Plymouth Harbor. These three locations are reported to be background locations by Entergy for federal reporting requirements, but are considered to be "indicator" locations by MDPH/BEH because they fall within the 10-mile EPZ. MERL analyzes the split samples for gamma-emitting radionuclides. Entergy also collects mussels from the Pilgrim discharge canal.

Sediment

In 2018 Entergy collected semi-annual sediment samples from the Pilgrim discharge canal and Green Harbor in Marshfield; MERL analyzed the split samples.

Irish moss

Irish moss (i.e., *Chondrus*) readily absorbs iodine and is a good reference indicator of iodine-131 in the environment. Entergy collects samples of Irish moss from the Pilgrim discharge canal and a background location at Brant Rock in Marshfield semiannually; split samples are analyzed by MERL.

Crops

MDPH/BEH collects and analyzes background cranberry samples from a bog in East Taunton annually.

In 2018 MDPH collected samples of green beans, green squash and tomatoes during the growing season, from a farm in Kingston

B. SEABROOK NUCLEAR POWER STATION

The Seabrook Nuclear Power Station (Seabrook) is located in Seabrook, New Hampshire, approximately two miles north of the Massachusetts border. Six Massachusetts communities are included in the 10-mile EPZ of Seabrook: Amesbury, Merrimac, Newbury, Newburyport, Salisbury, and West Newbury as shown in Figure 3.




Figure 3. Seabrook EPZ and sampling locations within Massachusetts

Radiation monitoring conducted within and outside the Seabrook EPZ includes the following environmental media: air, surface water, fish, shellfish, sediment, Irish moss, crops, and milk. MDPH/BEH receives split samples from Nextera, the utility that owns Seabrook, for all media except milk and air. Sampling locations and activities within Massachusetts are described below. Sampling locations are shown in Figure 3.

Air/Direct Radiation

MDPH/BEH collects air particulate filters and charcoal cartridges weekly at the Salisbury Fire Station. Filters are analyzed for gross beta and alpha radioactivity, and cartridges for iodine-131. Additionally, a

filter composite is analyzed quarterly for gamma-emitting radionuclides. The same analyses are done for air particulate filters and charcoal cartridges collected at the background location in Boston.

MDPH/BEH measures total ambient gamma radiation using a network of 34 TLDs placed at locations throughout the Seabrook EPZ in Massachusetts. These are collected and analyzed quarterly and results are compared to those of a background location in Boston.

MDPH/BEH contracts with the C-10 Research & Education Foundation, Inc. to conduct radiation monitoring in Massachusetts communities located in the Seabrook EPZ. The C-10 system consists of a network of 14 real-time radiation sensors and weather probes located in Massachusetts within a 10-mile radius of Seabrook station. Beta, gamma, and weather data are collected and uploaded every 15 minutes to a secure web-based central repository. C-10 compiles and graphs the data monthly and sends reports to MDPH/BEH. The 14 monitoring sites within the Seabrook 10-mile EPZ are located at private homes, schools, and businesses. MDPH and MEMA officials receive text alerts from C-10 if levels are greater than three times the typical background readings.

Surface Water

Seawater samples are typically collected monthly by Nextera from a background location in Ipswich Bay. MERL analyzes split samples for gamma-emitting radionuclides. MERL also analyzes surface water samples for tritium.

Milk

MDPH/BEH collects samples of cow's milk monthly from a farm located in Rowley and MERL analyzes the samples for gamma-emitting radionuclides, including iodine-131.

Fish and Shellfish

Nextera semi-annually collects samples of fish and shellfish, including lobster, *Modiolus* (i.e., Atlantic ribbed mussels) and *Mytilus* (i.e., blue or common mussels), from Ipswich Bay, which is considered a background location; MERL analyzes the split samples for gamma-emitting radionuclides.

Sediment

Nextera semi-annually collects sediment samples from Ipswich Bay and the tidal flats on Plum Island, both background locations; MERL analyzes the split samples for gamma-emitting radionuclides.

Irish moss

As noted earlier, Irish moss readily absorbs iodine and is a sentinel indicator of environmental iodine-131. Nextera collects samples of Irish moss semiannually from a background location in Ipswich Bay, and split samples are analyzed by MERL for gamma-emitting radionuclides.

Crops

In 2018 MDPH collected green beans, strawberries and tomatoes from a farm located within the Seabrook EPZ in Salisbury, and collected strawberries, tomatoes, and zucchini from a farm in Ipswich, which is outside the Seabrook EPZ.

3. 2018 ENVIRONMENTAL MONITORING RESULTS

Results of environmental monitoring conducted by MDPH/BEH in the Massachusetts communities in the vicinity of the Pilgrim and Seabrook nuclear power stations are discussed below and presented in Tables 1-7. The tables are organized by nuclear power station and by sample media.

A. PILGRIM NUCLEAR POWER STATION

Sampling results for Pilgrim are provided in Tables 2, 3 and 4. Only three radionuclides – Potassium-40, Beryllium-7, and Iodine-131, were detected in samples collected either within or outside the Pilgrim EPZ.

Naturally occurring Potassium-40 was detected in all samples of environmental media analyzed for Potassium-40 from both within and outside of the Pilgrim EPZ. Beryllium-7, which is also naturally occurring, was detected in all four of the quarterly composite air samples; in clams from both Plymouth and Duxbury Harbors; in mussels from the discharge canal; and in Irish moss samples from both the Pilgrim discharge canal and the background location at Brant Rock. For both Potassium-40 and Beryllium-7, levels detected in samples collected within the EPZ are generally consistent with levels detected in background samples.

Although not detected at levels of human health concern (Delacroix et al., 2002) Iodine-131 was detected in the Irish moss sample collected in June from the discharge canal (at 45 pCi/kg) and the sample collected in May from the background location at Brant Rock, in Marshfield (at 50 pCi/kg), at levels just above the respective laboratory reporting levels. Iodine-131 is monitored carefully as it may be released from nuclear power plants. Iodine-131 is also commonly used for treating individuals with overactive thyroids and thyroid cancer, which can result in discharge of Iodine-131 into sanitary sewers (Rose et al., 2012). Iodine-131 was not detected in any of the surface water samples collected for Pilgrim, nor was it detected in any of the air samples collected from the charcoal cartridges just outside the Pilgrim utility fence. In 2015, Iodine-131 was similarly detected at levels just above the laboratory reporting levels in Irish moss samples collected at both the discharge canal and Brant Rock. As in 2015, the Iodine-131 detected in the Irish moss samples does not appear to be connected to PNPS operations.

In addition to naturally occurring Beryllium-7 and Potassium-40, which were detected at levels consistent with background levels, air filter and cartridge analyses indicated low levels of gross alpha and gross beta radiation. Levels of gross alpha and beta radiation measured in the Pilgrim EPZ ranged from 0.002 – 0.018 and 0.015 – 0.068 pCi/m³, respectively. These levels are consistent with those measured at the background location in Boston, of 0.003 – 0.011 and 0.015 – 0.049 pCi/m³, respectively. No gamma-emitting radionuclides of concern were detected in quarterly composite air samples.

Real-time monitoring did not detect radiation greater than typical background levels of approximately 0.007 - 0.009 mRoentgen/hour with the exception of brief increases up to 0.02 mRoentgen/hour. Brief increases are expected due to rainfall washout from naturally occurring radionuclides such as airborne radon daughters and cosmic radiation events. No alerts at three times background were recorded. TLD total gamma exposure results ranged from 11.7 to 31.1 mRoentgen/quarter (i.e., 0.005 – 0.014 mRoentgen/hour) with an average of 16.1 mRoentgen/quarter (0.007 mRoentgen/hour). This value is compared to an average value of 13.5 mRoentgen/quarter measured at a background location in Boston, and corresponds to an average gamma exposure of 2.6 mRoentgen/quarter above background.

B. SEABROOK NUCLEAR POWER STATION

Seabrook sampling results are provided in Tables 5, 6 and 7. As with Pilgrim, the only radionuclides detected either within or outside the Seabrook EPZ were Potassium-40, Beryllium-7 and Iodine-131.

Naturally occurring Potassium-40 was detected in all samples of environmental media from both within and outside the Seabrook EPZ. Naturally occurring Beryllium-7 was detected in Irish moss (i.e., chondrus) and mussels (i.e., modiolus and mytilus) collected from the background sampling location in Ipswich Bay, in strawberries collected from the farm in Salisbury, and in the four quarterly air samples collected at the Salisbury Fire Station. As with Pilgrim, levels of Potassium-40 and Beryllium-7 detected within the Seabrook EPZ are consistent with background levels.

In addition to naturally occurring Beryllium-7 and Potassium-40, detected at levels consistent with background, analyses of air filter and cartridge samples found low levels of gross alpha ($0.004 - 0.011$ pCi/m³) and gross beta ($0.012 - 0.032$ pCi/m³) radiation, consistent with levels measured at the background location in Boston ($0.003 - 0.011$ pCi/m³ and $0.015 - 0.049$ pCi/m³, for gross alpha and gross beta radiation, respectively). No gamma radionuclides of concern were detected in quarterly composite air samples.

Although not detected at levels of human health concern (Delacroix et al., 2002) Iodine-131 was detected at a level of 194 pCi/kg in the Irish moss (i.e., chondrus) sample collected in May from the background location in Ipswich Bay, approximately 20 miles from the Seabrook EPZ. Radioactive iodine was found in Irish moss from the same location in previous years (2011, 2012, 2014 2015, and 2017). Iodine-131 was not detected in any of the surface water samples collected from Ipswich Bay, or in any of the air samples collected from the charcoal cartridge at the Salisbury Fire Station. Thus, sewage discharge and a nearby medical center are the most likely sources of the Iodine-131 detected in the Irish moss.

In 2018, real-time monitoring for the Seabrook EPZ did not show gamma radiation levels above typical background levels at most stations (approximately 0.010 mRoentgen/hour) with the exception of brief increases (typically up to approximately 0.02 mRoentgen/hour). Brief increases are expected due to rainfall washout from naturally occurring radionuclides such as airborne radon daughters and cosmic radiation events. Beta readings ranged from approximately 40 to 50 counts per minute with the exception of brief increases similar to the gamma results.

TLD results for total gamma exposure ranged from 10.6 to 21.3 mRoentgen/quarter ($0.005 - 0.010$ mRoentgen/hour) with an average exposure of 17.2 mRoentgen/quarter (0.008 mRoentgen/hour), compared to an average of 13.8 mRoentgen/quarter at the background location in Boston. The result is an average gamma exposure level of 3.4 mRoentgen/quarter over background.

C. SUMMARY

Radiation monitoring results in 2018 for Massachusetts have been either non-detect or naturally occurring (i.e., Potassium-40, Beryllium-7). No detectible radionuclides were at levels of health concern or were indicative of an unintentional release of radiation at Pilgrim or Seabrook. Iodine-131 was detected at levels just above the laboratory reporting levels in samples of Irish moss collected both outside the Pilgrim EPZ and from the Pilgrim discharge canal. Iodine-131 was also detected at higher levels in samples of Irish moss collected from a background location outside the Seabrook EPZ.

C. REFERENCES

- Amund et al., 1996. *Contaminants in Fish and Sediments in the North Atlantic Ocean*. TemaNord.
- U.S. Agency for Toxic Substances and Disease Registry (ATSDR). 2002. Case Studies in Environmental Medicine: Radiation Exposure from Iodine 131.
- U.S. Agency for Toxic Substances and Disease Registry (ATSDR). 2004.. *Toxicological Profile for Cesium*.
- Burger et al., 2007. "Radionuclides in Marine Fishes and Birds from Amchitka and Kiska Islands in the Aleutians: Establishing a baseline." *Health Physics*. March 2007, Vol. 92, No.3.
- Delacroix et al., 2002. "Radionuclide and Radiation Protection Data Handbook 2002." *Radiation Protection Dosimetry* 98(1)
- Delaygue et al., 2015. "Modelling the stratospheric budget of beryllium isotopes." *Chemical and Physical Meteorology*. 67:1
- Luykx, F and G. Fraser. 1983. "Radioactive effluents from nuclear power stations and nuclear fuel reprocessing plants in the European Community, Discharge Data 1976 – 1980, Radiological Aspects." Report prepared for Commission of the European Communities, Directorate – General Employment, Social Affairs and Education, Health and Safety Directorate.
- Rose et al., 2012. "Medically-derived I-131 in municipal sewage effluent". *Water Research* 46 (2012):5663-5671.
- United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR). 2008. Volume I: Sources and Effects of Ionizing Radiation; Annex B: Exposures of the Public and Workers from Various Sources of Radiation. [Accessed at: https://www.unscear.org/unscear/publications/2008_1.html](https://www.unscear.org/unscear/publications/2008_1.html).
- US United States Nuclear Regulatory Commission (NRC). 2017a. Natural Background Sources. <https://www.nrc.gov/about-nrc/radiation/around-us/sources/nat-bg-sources.html>.
- US United States Nuclear Regulatory Commission(NRC). 2017b. Man-Made Sources. <https://www.nrc.gov/about-nrc/radiation/around-us/sources/man-made-sources.html>.
- US United States Nuclear Regulatory Commission(NRC). 2017c. Doses in Our Daily Lives. <http://www.nrc.gov/about-nrc/radiation/around-us/doses-daily-lives.html>.
- Yim, MS and F Caron (2006). Life cycle and management of carbon-14 from nuclear power generation. *Prog Nucl Energ*. 48:2-36.

D. RESULTS TABLES

Table 2. Pilgrim Nuclear Power Station 2018 Environmental Monitoring Data - Air Samples

Sample Type	Location	Date	I-131* (pCi/m ³)	Be-7* (pCi/m ³)	K-40* (pCi/m ³)	Mn-54* (pCi/m ³)	Fe-59* (pCi/m ³)	Co-60* (pCi/m ³)	Zn-65* (pCi/m ³)	Cs-137* (pCi/m ³)	Gross Alpha (pCi/m ³)	Gross Beta (pCi/m ³)
Air	Pilgrim Station	1/8/2018	<0.0101	-	-	-	-	-	-	-	0.00641	0.0228
Air	Pilgrim Station	1/12/2018	<0.0277	-	-	-	-	-	-	-	0.01640	0.0527
Air	Pilgrim Station	1/19/2018	<0.0162	-	-	-	-	-	-	-	0.00735	0.0239
Air	Pilgrim Station	1/23/2018	<0.0342	-	-	-	-	-	-	-	0.01770	0.0684
Air	Pilgrim Station	2/1/2018	<0.0107	-	-	-	-	-	-	-	0.00477	0.0197
Air	Pilgrim Station	2/15/2018	<0.0066	-	-	-	-	-	-	-	0.00438	0.0226
Air	Pilgrim Station Quarterly composite	2/15/2018	-	0.096	0.262	<0.001	<0.003	<0.001	<0.002	<0.0001	-	-
Air	Pilgrim Station	2/23/2018	<0.0146	-	-	-	-	-	-	-	0.00486	0.0290
Air	Pilgrim Station	2/28/2018	<0.0247	-	-	-	-	-	-	-	0.00638	0.0292
Air	Pilgrim Station	3/9/2018	<0.0131	-	-	-	-	-	-	-	0.00261	0.0160
Air	Pilgrim Station	3/16/2018	<0.0147	-	-	-	-	-	-	-	0.00380	0.0169
Air	Pilgrim Station	3/23/2018	<0.0186	-	-	-	-	-	-	-	0.00332	0.0297
Air	Pilgrim Station	3/29/2018	<0.0203	-	-	-	-	-	-	-	0.00307	0.0251
Air	Pilgrim Station	4/5/2018	<0.0152	-	-	-	-	-	-	-	0.00518	0.0240
Air	Pilgrim Station	4/12/2018	<0.0188	-	-	-	-	-	-	-	0.00331	0.0206
Air	Pilgrim Station	4/20/2018	<0.0110	-	-	-	-	-	-	-	0.00517	0.0184
Air	Pilgrim Station	4/26/2018	<0.0211	-	-	-	-	-	-	-	0.00820	0.0258
Air	Pilgrim Station	5/2/2018	<0.0146	-	-	-	-	-	-	-	0.00528	0.0193
Air	Pilgrim Station	5/10/2018	<0.0143	-	-	-	-	-	-	-	0.00652	0.0246
Air	Pilgrim Station Quarterly composite	5/15/2018	-	0.126	0.145	<0.001	<0.004	<0.001	<0.002	<0.001	-	-
Air	Pilgrim Station	5/18/2018	<0.0149	-	-	-	-	-	-	-	0.00463	0.0176
Air	Pilgrim Station	5/24/2018	<0.0196	-	-	-	-	-	-	-	0.00617	0.0218
Air	Pilgrim Station	5/31/2018	<0.0140	-	-	-	-	-	-	-	0.00649	0.0235
Air	Pilgrim Station	6/7/2018	<0.0115	-	-	-	-	-	-	-	0.00489	0.0159
Air	Pilgrim Station	6/15/2018	<0.0104	-	-	-	-	-	-	-	0.00429	0.0210
Air	Pilgrim Station	6/20/2018	<0.0216	-	-	-	-	-	-	-	0.00360	0.0267
Air	Pilgrim Station	6/29/2018	<0.0086	-	-	-	-	-	-	-	0.00241	0.0191
Air	Pilgrim Station	7/6/2018	<0.0159	-	-	-	-	-	-	-	0.00911	0.0302
Air	Pilgrim Station	7/12/2018	<0.0235	-	-	-	-	-	-	-	0.00513	0.0229
Air	Pilgrim Station	7/25/2018	<0.00745	-	-	-	-	-	-	-	0.00548	0.0153
Air	Pilgrim Station	8/3/2018	<0.0118	-	-	-	-	-	-	-	0.00664	0.0209

- " = Not analyzed

* I-131 = iodine 131; Ba-140 = Barium 140; Be-7 = Beryllium 7; Co-60 = Cobalt 60; Cs-134 = Cesium 134; Cs-137=Cesium 137; Fe-59=Iron 59; H-3* = Tritium; K-40 = Potassium-40; Mn-54=Manganese-54; Zn-65=Zinc 65

"<" = value is less than the listed MDA (Minimum Detectable Activity) value

"NR"= result not reported for quality control reasons.

Sample Type	Location	Date	I-131* (pCi/m ³)	Be-7* (pCi/m ³)	K-40* (pCi/m ³)	Mn-54* (pCi/m ³)	Fe-59* (pCi/m ³)	Co-60* (pCi/m ³)	Zn-65* (pCi/m ³)	Cs-137* (pCi/m ³)	Gross Alpha (pCi/m ³)	Gross Beta (pCi/m ³)
Air	Pilgrim Station	8/9/2018	<0.0222	-	-	-	-	-	-	-	0.01500	0.0375
Air	Pilgrim Station Quarterly composite	8/15/2018	-	0.105	<0.016	<0.001	<0.003	<0.001	<0.001	<0.001	-	-
Air	Pilgrim Station	8/16/2018	<0.0178	-	-	-	-	-	-	-	0.01050	0.0328
Air	Pilgrim Station	8/23/2018	<0.0170	-	-	-	-	-	-	-	0.00884	0.0264
Air	Pilgrim Station	8/30/2018	<0.0137	-	-	-	-	-	-	-	0.01210	0.0448
Air	Pilgrim Station	9/6/2018	<0.0130	-	-	-	-	-	-	-	0.00521	0.0238
Air	Pilgrim Station	9/13/2018	<0.0152	-	-	-	-	-	-	-	0.00508	0.0180
Air	Pilgrim Station	9/21/2018	<0.0117	-	-	-	-	-	-	-	0.00548	0.0226
Air	Pilgrim Station	9/27/2018	<0.0150	-	-	-	-	-	-	-	0.00622	0.0203
Air	Pilgrim Station	10/3/2018	<0.0143	-	-	-	-	-	-	-	0.00931	0.0266
Air	Pilgrim Station	10/10/2018	<0.0161	-	-	-	-	-	-	-	0.00836	0.0251
Air	Pilgrim Station	10/17/2018	<0.0166	-	-	-	-	-	-	-	0.00874	0.0250
Air	Pilgrim Station	10/26/2018	<0.0106	-	-	-	-	-	-	-	0.00583	0.0217
Air	Pilgrim Station	10/31/2018	<0.0204	-	-	-	-	-	-	-	0.00568	0.0233
Air	Pilgrim Station	11/8/2018	<0.0122	-	-	-	-	-	-	-	0.00781	0.0248
Air	Pilgrim Station	11/15/2018	<0.0122	-	-	-	-	-	-	-	0.00800	0.0247
Air	Pilgrim Station Quarterly composite	11/15/2018	-	0.078	0.272	<0.001	<0.003	<0.001	<0.002	<0.001	-	-
Air	Pilgrim Station	11/23/2018	<0.0154	-	-	-	-	-	-	-	0.00560	0.0243
Air	Pilgrim Station	11/30/2018	<0.0165	-	-	-	-	-	-	-	0.00613	0.0217
Air	Pilgrim Station	12/6/2018	<0.0199	-	-	-	-	-	-	-	0.00735	0.0215
Air	Pilgrim Station	12/14/2018	<0.0115	-	-	-	-	-	-	-	0.00492	0.0331
Air	Pilgrim Station	12/20/2018	<0.0185	-	-	-	-	-	-	-	0.00511	0.0320
Air	Pilgrim Station	12/27/2018	<0.0182	-	-	-	-	-	-	-	0.00329	0.0278
Air	Background	1/2/2018	<0.0185	-	-	-	-	-	-	-	0.00877	0.0326
Air	Background	1/9/2018	<0.0199	-	-	-	-	-	-	-	0.00894	0.0311
Air	Background	1/16/2018	<0.0156	-	-	-	-	-	-	-	0.01010	0.0353
Air	Background	1/22/2018	<0.0219	-	-	-	-	-	-	-	0.01100	0.0489
Air	Background	1/29/2018	<0.0184	-	-	-	-	-	-	-	0.01140	0.0313
Air	Background	2/5/2018	<0.0175	-	-	-	-	-	-	-	0.00949	0.0270
Air	Background	2/12/2018	<0.0187	-	-	-	-	-	-	-	0.00999	0.0300
Air	Background quarterly composite	2/15/2018	-	0.099	0.280	<0.001	0.003	<0.001	<0.002	<0.001	-	-

- " = Not analyzed

* I-131 = iodine 131; Ba-140 = Barium 140; Be-7 = Beryllium 7; Co-60 = Cobalt 60; Cs-134 = Cesium 134; Cs-137=Cesium 137; Fe-59=Iron 59; H-3* = Tritium; K-40 = Potassium-40; Mn-54=Manganese-54; Zn-65=Zinc 65

"<" = value is less than the listed MDA (Minimum Detectable Activity) value

"NR"= result not reported for quality control reasons.

Sample Type	Location	Date	I-131* (pCi/m ³)	Be-7* (pCi/m ³)	K-40* (pCi/m ³)	Mn-54* (pCi/m ³)	Fe-59* (pCi/m ³)	Co-60* (pCi/m ³)	Zn-65* (pCi/m ³)	Cs-137* (pCi/m ³)	Gross Alpha (pCi/m ³)	Gross Beta (pCi/m ³)
Air	Background	2/20/2018	<0.0139	-	-	-	-	-	-	-	0.00701	0.0345
Air	Background	2/26/2018	<0.0152	-	-	-	-	-	-	-	0.00605	0.0226
Air	Background	3/5/2018	<0.0120	-	-	-	-	-	-	-	0.00490	0.0259
Air	Background	3/12/2018	<0.0172	-	-	-	-	-	-	-	0.00407	0.0152
Air	Background	3/19/2018	<0.0172	-	-	-	-	-	-	-	0.00625	0.0257
Air	Background	3/26/2018	<0.0184	-	-	-	-	-	-	-	0.00430	0.0199
Air	Background	4/2/2018	<0.0178	-	-	-	-	-	-	-	0.00691	0.0253
Air	Background	4/9/2018	<0.0174	-	-	-	-	-	-	-	0.00530	0.0264
Air	Background	4/17/2018	<0.0160	-	-	-	-	-	-	-	0.00570	0.0252
Air	Background	4/23/2018	<0.0202	-	-	-	-	-	-	-	0.00383	0.0204
Air	Background	4/30/2018	<0.0188	-	-	-	-	-	-	-	0.00436	0.0203
Air	Background	5/7/2018	<0.0173	-	-	-	-	-	-	-	0.00591	0.0292
Air	Background	5/14/2018	<0.0180	-	-	-	-	-	-	-	0.00401	0.0187
Air	Background quarterly composite	5/15/2018	-	0.124	<0.025	<0.001	<0.004	<0.001	<0.002	<0.001	-	-
Air	Background	5/21/2018	<0.0164	-	-	-	-	-	-	-	0.00404	0.0176
Air	Background	5/29/2018	<0.0111	-	-	-	-	-	-	-	0.00335	0.0234
Air	Background	6/4/2018	<0.0190	-	-	-	-	-	-	-	0.00445	0.0197
Air	Background	6/11/2018	<0.0134	-	-	-	-	-	-	-	0.00334	0.0189
Air	Background	6/18/2018	<0.0165	-	-	-	-	-	-	-	0.00263	0.0151
Air	Background	6/25/2018	<0.0173	-	-	-	-	-	-	-	0.00477	0.0254
Air	Background	7/2/2018	<0.0172	-	-	-	-	-	-	-	0.00700	0.0250
Air	Background	7/9/2018	<0.0182	-	-	-	-	-	-	-	0.00810	0.0276
Air	Background	7/16/2018	<0.0178	-	-	-	-	-	-	-	0.00602	0.0257
Air	Background	7/23/2018	<0.0179	-	-	-	-	-	-	-	0.00618	0.0224
Air	Background	7/30/2018	<0.0167	-	-	-	-	-	-	-	0.00455	0.0198
Air	Background	8/6/2018	<0.0183	-	-	-	-	-	-	-	0.00660	0.0234
Air	Background	8/13/2018	<0.0153	-	-	-	-	-	-	-	0.00747	0.0408
Air	Background quarterly composite	8/15/2018	-	0.109	<0.025	<0.001	<0.004	<0.001	<0.002	<0.001	-	-
Air	Background	8/20/2018	<0.0186	-	-	-	-	-	-	-	0.00719	0.0290
Air	Background	8/27/2018	<0.0180	-	-	-	-	-	-	-	0.00547	0.0263
Air	Background	9/4/2018	<0.0165	-	-	-	-	-	-	-	0.00658	0.0364
Air	Background	9/10/2018	<0.0222	-	-	-	-	-	-	-	0.00481	0.0278

- " = Not analyzed

* I-131 = iodine 131; Ba-140 = Barium 140; Be-7 = Beryllium 7; Co-60 = Cobalt 60; Cs-134 = Cesium 134; Cs-137=Cesium 137; Fe-59=Iron 59; H-3* = Tritium; K-40 = Potassium-40; Mn-54=Manganese-54; Zn-65=Zinc 65

"<" = value is less than the listed MDA (Minimum Detectable Activity) value

"NR"= result not reported for quality control reasons.

Sample Type	Location	Date	I-131* (pCi/m ³)	Be-7* (pCi/m ³)	K-40* (pCi/m ³)	Mn-54* (pCi/m ³)	Fe-59* (pCi/m ³)	Co-60* (pCi/m ³)	Zn-65* (pCi/m ³)	Cs-137* (pCi/m ³)	Gross Alpha (pCi/m ³)	Gross Beta (pCi/m ³)
Air	Background	9/17/2018	<0.0178	-	-	-	-	-	-	-	0.00381	0.0222
Air	Background	9/24/2018	<0.0176	-	-	-	-	-	-	-	0.00294	0.0203
Air	Background	10/1/2018	<0.0135	-	-	-	-	-	-	-	0.00607	0.0221
Air	Background	10/9/2018	<0.0126	-	-	-	-	-	-	-	0.00764	0.0225
Air	Background	10/15/2018	<0.0223	-	-	-	-	-	-	-	0.00954	0.0277
Air	Background	10/22/2018	<0.0153	-	-	-	-	-	-	-	0.00990	0.0279
Air	Background	10/29/2018	<0.0138	-	-	-	-	-	-	-	0.00804	0.0230
Air	Background	11/5/2018	<0.0177	-	-	-	-	-	-	-	0.00866	0.0290
Air	Background	11/13/2018	<0.0146	-	-	-	-	-	-	-	0.00643	0.0251
Air	Background quarterly composite	11/15/2018	-	0.089	0.147	<0.001	<0.003	<0.001	<0.002	<0.001	-	-
Air	Background	11/19/2018	<0.0194	-	-	-	-	-	-	-	0.00945	0.0268
Air	Background	11/26/2018	<0.0199	-	-	-	-	-	-	-	0.00732	0.0262
Air	Background	12/3/2018	<0.0176	-	-	-	-	-	-	-	0.00881	0.0179
Air	Background	12/10/2018	<0.0190	-	-	-	-	-	-	-	0.00755	0.0303
Air	Background	12/17/2018	<0.0179	-	-	-	-	-	-	-	0.00350	0.0333
Air	Background	12/24/2018	<0.0153	-	-	-	-	-	-	-	0.00329	0.0265
Air	Background	12/31/2018	<0.0140	-	-	-	-	-	-	-	0.00339	0.0245

- = Not analyzed

* I-131 = iodine 131; Ba-140 = Barium 140; Be-7 = Beryllium 7; Co-60 = Cobalt 60; Cs-134 = Cesium 134; Cs-137=Cesium 137; Fe-59=Iron 59; H-3* = Tritium; K-40 = Potassium-40; Mn-54=Manganese-54; Zn-65=Zinc 65

"<" = value is less than the listed MDA (Minimum Detectable Activity) value

"NR"= result not reported for quality control reasons.

Table 3. Pilgrim Nuclear Power Station 2018 Environmental Monitoring Data – Liquid Matrices

Sample Type	Location	Date	K-40* (pCi/L)	Mn-54* (pCi/L)	Fe-59* (pCi/L)	Co-60* (pCi/L)	Zn-65* (pCi/L)	I-131* (pCi/L)	Cs-134* (pCi/L)	Cs-137* (pCi/L)	Ba-140* (pCi/L)	H-3* (pCi/L)
Surface water	Discharge Canal	1/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Discharge Canal	1/31/2018	301	<2.5	<5.9	<2.4	<5.0	<15.8	-	<2.6	-	-
Surface water	Discharge Canal	2/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Discharge Canal	2/27/2018	351	<3.1	<7.3	<2.8	<6.3	<13.1	-	<2.9	-	-
Surface water	Discharge Canal	3/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Discharge Canal	4/3/2018	296	<3.0	<7.1	<2.7	<6.1	<13.2	-	<3.0	-	-
Surface water	Discharge Canal	4/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Discharge Canal	5/1/2018	1440	<2.9	<6.5	<2.5	<6.1	<11.1	-	<2.8	-	-
Surface water	Discharge Canal	5/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Discharge Canal	5/29/2018	291	<2.9	<6.7	<2.9	<6.4	<10.0	-	<2.8	-	-
Surface water	Discharge Canal	6/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Discharge Canal	7/3/2018	323	<3.0	<6.0	<2.8	<6.4	<6.0	-	<3.0	-	-
Surface water	Discharge Canal	7/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Discharge Canal	7/31/2018	878	<2.8	<6.2	<3.2	<5.7	<5.1	-	<3.2	-	-
Surface water	Discharge Canal	8/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Discharge Canal	8/28/2018	916	<3.2	<6.4	<3.3	<7.1	<3.9	-	<3.3	-	-
Surface water	Discharge Canal	9/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Discharge Canal	10/2/2018	342	<3.1	<6.4	<2.9	<6.6	<10.5	-	<2.7	-	-
Surface water	Discharge Canal	10/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Discharge Canal	10/30/2018	286	<3.0	<6.7	<2.8	<6.4	<11.0	-	<3.0	-	-
Surface water	Discharge Canal	11/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Discharge Canal	11/27/2018	325	<2.9	<6.4	<2.8	<6.4	<5.5	-	<2.9	-	-
Surface water	Discharge Canal	12/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Discharge Canal ¹	1/2/2019	297	<3.1	<7.6	<2.9	<6.8	<17.7	-	<2.9	-	-
Surface water	Powder Point Bridge ²	1/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Powder Point Bridge ²	1/31/2018	301	<2.5	<5.8	<2.6	<5.0	<12.7	-	<2.6	-	-
Surface water	Powder Point Bridge ²	2/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Powder Point Bridge ²	2/27/2018	271	<2.5	<5.6	<2.6	<5.1	<8.4	-	<2.7	-	-
Surface water	Powder Point Bridge ²	3/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Powder Point Bridge ²	4/3/2018	291	<2.5	<5.7	<2.6	<5.8	<9.1	-	<2.6	-	-
Surface water	Powder Point Bridge ²	4/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Powder Point Bridge ²	5/1/2018	275	<2.4	<5.7	<2.4	<5.2	<8.3	-	<2.5	-	-
Surface water	Powder Point Bridge ²	5/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Powder Point Bridge ²	5/29/2018	315	<2.3	<5.0	<2.5	<5.3	<6.9	-	<2.5	-	-

- = Not analyzed

* I-131 = iodine 131; Ba-140 = Barium 140; Be-7 = Beryllium 7; Co-60 = Cobalt 60; Cs-134 = Cesium 134; Cs-137=Cesium 137; Fe-59=Iron 59; H-3* = Tritium; K-40 = Potassium-40; Mn-54=Manganese-54; Zn-65=Zinc 65

< = value is less than the listed MDA (Minimum Detectable Activity) value

“NR”= result not reported for quality control reasons.

Sample Type	Location	Date	K-40* (pCi/L)	Mn-54* (pCi/L)	Fe-59* (pCi/L)	Co-60* (pCi/L)	Zn-65* (pCi/L)	I-131* (pCi/L)	Cs-134* (pCi/L)	Cs-137* (pCi/L)	Ba-140* (pCi/L)	H-3* (pCi/L)
Surface water	Powder Point Bridge ²	6/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Powder Point Bridge ²	7/3/2018	663	<2.4	<4.7	<2.6	<5.3	<4.0	-	<2.7	-	-
Surface water	Powder Point Bridge ²	7/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Powder Point Bridge ²	7/31/2018	299	<2.5	<5.4	<2.2	<5.3	<4.3	-	<2.8	-	-
Surface water	Powder Point Bridge ²	8/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Powder Point Bridge ²	8/28/2018	338	<2.7	<4.9	<2.8	<5.4	<3.2	-	<2.6	-	-
Surface water	Powder Point Bridge ²	9/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Powder Point Bridge ²	10/2/2018	730	<2.5	<5.9	<2.6	<5.2	<7.8	-	<2.5	-	-
Surface water	Powder Point Bridge ²	10/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Powder Point Bridge ²	10/30/2018	298	<2.6	<5.2	<2.8	<5.7	<7.9	-	<2.7	-	-
Surface water	Powder Point Bridge ²	11/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Powder Point Bridge ²	11/27/2018	668	<2.5	<5.0	<2.7	<5.3	<4.0	-	<2.6	-	-
Surface water	Powder Point Bridge ²	12/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Powder Point Bridge ^{1,2}	1/2/2019	319	<3.1	<7.4	<2.9	<6.5	<10.9	-	<2.9	-	-
Milk	Duxbury	1/12/2018	1360	-	-	-	-	<3.9	<2.7	<3.2	<13	-
Milk	Duxbury	2/15/2018	1330	-	-	-	-	<3.1	<2.7	<3.1	<10.5	-
Milk	Duxbury	3/16/2018	1280	-	-	-	-	<3.1	<2.5	<3.0	<10.6	-
Milk	Duxbury	4/12/2018	1230	-	-	-	-	<3.0	<2.6	<3.0	<10.2	-
Milk	Duxbury	5/10/2018	1240	-	-	-	-	<2.3	<2.3	<2.9	<9.2	-
Milk	Duxbury	6/7/2018	1270	-	-	-	-	<3.1	<2.6	<2.8	<10.3	-
Milk	Duxbury	7/12/2018	1270	-	-	-	-	<3.3	<2.6	<3.0	<10.3	-
Milk	Duxbury	8/9/2018	1310	-	-	-	-	<3.5	<2.7	<3.4	<11.2	-
Milk	Duxbury	9/27/2018	1130	-	-	-	-	<3.2	<2.5	<3.0	<10.5	-
Milk	Duxbury	10/10/2018	1220	-	-	-	-	<3.6	<2.6	<3.1	<11.3	-
Milk	Duxbury	11/8/2018	1210	-	-	-	-	<3.2	<2.6	<3.1	<10.6	-
Milk	Duxbury	12/6/2018	1390	-	-	-	-	<3.3	<2.7	<3.2	<11.3	-

(1) Sample is composite for December 2018

(2) Sample considered "background" for the purpose of NRC regulations, but considered "indicator" by MDPH because it falls within the 10-mile EPZ.

- " = Not analyzed

* I-131 = iodine 131; Ba-140 = Barium 140; Be-7 = Beryllium 7; Co-60 = Cobalt 60; Cs-134 = Cesium 134; Cs-137=Cesium 137; Fe-59=Iron 59; H-3* = Tritium; K-40 = Potassium-40; Mn-54=Manganese-54; Zn-65=Zinc 65

"<" = value is less than the listed MDA (Minimum Detectable Activity) value

"NR"= result not reported for quality control reasons.

Table 4. Pilgrim Nuclear Power Station 2018 Environmental Monitoring Data – Solid matrices

Sample	Location	Date	Be-7* (pCi/kg)	K-40* (pCi/kg)	Mn-54* (pCi/kg)	Fe-59* (pCi/kg)	Co-60* (pCi/kg)	Zn-65* (pCi/kg)	Cs-137* (pCi/kg)	I-131* (pCi/kg)
Atlantic Menhaden	PNPS Discharge Canal	7/20/2018	<45.3	4180	<5.4	<12.9	<5.1	<11.9	<4.8	-
Striped Bass	PNPS Discharge Canal	7/20/2018	<43.6	3850	<4.7	<12.2	<4.8	<11.0	<4.8	-
Winter Flounder	Cape Cod Bay (background)	5/2/2018	<45.3	4200	<4.8	<13.3	<4.8	<12.0	<4.8	-
Winter Flounder	PNPS - Discharge Canal	5/1/2018	<36.9	3910	<4.3	<12.6	<4.9	<9.8	<4.3	-
Lobster	Cape Cod Bay (background)	7/20/2018	<35	2640	<3.9	<9.8	<4.4	<9.5	<4.0	-
Lobster	PNPS Discharge Canal	7/21/2018	<238	2480	NR	NR	NR	<54.0	NR	-
Mytilus ¹	Green Harbor, Marshfield (background)	5/17/2018	<43.1	1510	<4.2	<8.5	<3.7	<8.4	<3.8	-
Mytilus ¹	Green Harbor, Marshfield (background)	10/8/2018	NR	1050	<7.0	NR	<4.7	<18	<3.9	-
Mytilus ¹	PNPS Discharge Canal	6/14/2018	35.1	1480	<3.3	<8.7	<3.8	<7.6	<3.5	-
Mytilus ¹	PNPS Discharge Canal	12/6/2018	NR	1430	<6.3	NR	<4.1	<15.8	<4.0	-
Softshell Clams	Duxbury	5/17/2018	69.8	1640	<3.6	<7.3	<3.6	<8.2	<3.7	-
Softshell Clams	Duxbury	10/8/2018	NR	1120	<6.0	NR	<4.2	<15.3	<3.6	-
Softshell Clams	Plymouth Harbor	5/18/2018	59.3	1610	<3.42	<7.3	<3.7	<7.3	<3.5	-
Softshell Clams	Plymouth Harbor	10/8/2018	NR	1220	<6.9	<371	<4.9	<17.4	<4.1	-
Sediment	Green Harbor, Marshfield	5/17/2018	-	10900	-	-	<22.5	-	<24.1	-
Sediment	Green Harbor, Marshfield	10/8/2018	-	10900	-	-	<29.1	-	<29.2	-
Sediment	PNPS Discharge Canal	6/14/2018	-	30800	-	-	<32.9	-	<35.3	-
Sediment	PNPS Discharge Canal	12/6/2018	-	20200	-	-	<26.7	-	<26.0	-
Irish Moss	Brant Rock (background)	5/12/2018	371	18400	<13.7	<37.4	<14.3	<35.4	<13.9	50.0
Irish Moss	Brant Rock (background)	10/9/2018	NR	25500	<25.2	NR	<15.7	<65.6	<14.6	-
Irish Moss	PNPS Discharge Canal	6/14/2018	657	33600	<16.5	<47.3	<18.8	<45.5	<15.9	45.2
Irish Moss	PNPS Discharge Canal	12/6/2018	1880	28200	NR	NR	<25.7	NR	<20.3	-
Cranberries	E. Taunton (background)	10/22/2018	<43.1	728	<4.4	<8.7	<5.1	<9.8	<4.9	-
Green Beans	Cretinon's Farm, Kingston	7/25/2018	<55.1	2540	<6.6	<11.8	<7.0	<13.5	<6.7	-
Green Squash	Cretinon's Farm, Kingston	7/25/2018	<38.3	2040	<4.9	<10.0	<4.4	<11.0	<4.9	-
Tomatoes	Cretinon's Farm, Kingston	8/23/2018	<32.8	2220	<4	<8.2	<4.5	<9.9	<3.9	-

¹Blue, or common mussel

- = Not analyzed

* I-131 = iodine 131; Ba-140 = Barium 140; Be-7 = Beryllium 7; Co-60 = Cobalt 60; Cs-134 = Cesium 134; Cs-137=Cesium 137; Fe-59=Iron 59; H-3* = Tritium; K-40 = Potassium-40; Mn-54=Manganese-54; Zn-65=Zinc 65

< = value is less than the listed MDA (Minimum Detectable Activity) value

"NR"= result not reported for quality control reasons.

Table 5. Seabrook Nuclear Power Station 2018 Environmental Monitoring Data - Air Samples

Sample Type	Location	Date	I-131* (pCi/m ³)	Be-7* (pCi/m ³)	K-40* (pCi/m ³)	Mn-54* (pCi/m ³)	Fe-59* (pCi/m ³)	Co-60* (pCi/m ³)	Zn-65* (pCi/m ³)	Cs-137* (pCi/m ³)	Gross Alpha (pCi/m ³)	Gross Beta (pCi/m ³)
Air	Salisbury Fire Station	1/3/2018	<0.0195	-	-	-	-	-	-	-	0.0096	0.0273
Air	Salisbury Fire Station	1/10/2018	<0.0237	-	-	-	-	-	-	-	0.0102	0.0275
Air	Salisbury Fire Station	1/17/2018	<0.0249	-	-	-	-	-	-	-	0.0097	0.0250
Air	Salisbury Fire Station	1/25/2018	<0.0147	-	-	-	-	-	-	-	0.0099	0.0318
Air	Salisbury Fire Station	1/31/2018	<0.0237	-	-	-	-	-	-	-	0.0112	0.0271
Air	Salisbury Fire Station	2/7/2018	<0.0166	-	-	-	-	-	-	-	0.0071	0.0234
Air	Salisbury Fire Station	2/14/2018	<0.0208	-	-	-	-	-	-	-	0.0089	0.0300
Air	Salisbury Fire Station quarterly composite	2/15/2018	-	0.076	0.097	<0.0006	<0.002	<0.0006	<0.001	<0.001	-	-
Air	Salisbury Fire Station	2/21/2018	<0.0190	-	-	-	-	-	-	-	0.0065	0.0305
Air	Salisbury Fire Station	3/2/2018	<0.0179	-	-	-	-	-	-	-	0.0050	0.0238
Air	Salisbury Fire Station	3/7/2018	<0.0222	-	-	-	-	-	-	-	0.0075	0.0214
Air	Salisbury Fire Station	3/12/2018	<0.0284	-	-	-	-	-	-	-	0.0048	0.0117
Air	Salisbury Fire Station	3/21/2018	<0.0151	-	-	-	-	-	-	-	0.0042	0.0144
Air	Salisbury Fire Station	3/27/2018	<0.0198	-	-	-	-	-	-	-	0.0043	0.0193
Air	Salisbury Fire Station	4/4/2018	<0.0196	-	-	-	-	-	-	-	0.0062	0.0238
Air	Salisbury Fire Station	4/11/2018	<0.0133	-	-	-	-	-	-	-	0.0073	0.0213
Air	Salisbury Fire Station	4/18/2018	<0.0193	-	-	-	-	-	-	-	0.0070	0.0232
Air	Salisbury Fire Station	4/24/2018	<0.0185	-	-	-	-	-	-	-	0.0067	0.0184
Air	Salisbury Fire Station	5/2/2018	<0.0114	-	-	-	-	-	-	-	0.0056	0.0132
Air	Salisbury Fire Station	5/9/2018	<0.0175	-	-	-	-	-	-	-	0.0080	0.0302
Air	Salisbury Fire Station quarterly composite	5/15/2018	-	0.097	0.099	<0.0007	<0.004	<0.0007	<0.002	<0.0006	-	-
Air	Salisbury Fire Station	5/16/2018	<0.0159	-	-	-	-	-	-	-	0.0055	0.0157
Air	Salisbury Fire Station	5/23/2018	<0.0169	-	-	-	-	-	-	-	0.0064	0.0163
Air	Salisbury Fire Station	5/30/2018	<0.0187	-	-	-	-	-	-	-	0.0072	0.0208
Air	Salisbury Fire Station	6/5/2018	<0.0159	-	-	-	-	-	-	-	0.0065	0.0162
Air	Salisbury Fire Station	6/12/2018	<0.0192	-	-	-	-	-	-	-	0.0060	0.0220
Air	Salisbury Fire Station	6/19/2018	<0.0121	-	-	-	-	-	-	-	0.0039	0.0156
Air	Salisbury Fire Station	6/27/2018	<0.0208	-	-	-	-	-	-	-	0.0055	0.0189
Air	Salisbury Fire Station	7/5/2018	<0.0143	-	-	-	-	-	-	-	0.0075	0.0239
Air	Salisbury Fire Station	7/11/2018	<0.0151	-	-	-	-	-	-	-	0.0066	0.0203
Air	Salisbury Fire Station	7/17/2018	<0.0187	-	-	-	-	-	-	-	0.0069	0.0161

- = Not analyzed

* I-131 = iodine 131; Ba-140 = Barium 140; Be-7 = Beryllium 7; Co-60 = Cobalt 60; Cs-134 = Cesium 134; Cs-137=Cesium 137; Fe-59=Iron 59; H-3* = Tritium; K-40 = Potassium-40; Mn-54=Manganese-54; Zn-65=Zinc 65

"<" = value is less than the listed MDA (Minimum Detectable Activity) value

"NR"= result not reported for quality control reasons.

Sample Type	Location	Date	I-131* (pCi/m ³)	Be-7* (pCi/m ³)	K-40* (pCi/m ³)	Mn-54* (pCi/m ³)	Fe-59* (pCi/m ³)	Co-60* (pCi/m ³)	Zn-65* (pCi/m ³)	Cs-137* (pCi/m ³)	Gross Alpha (pCi/m ³)	Gross Beta (pCi/m ³)
Air	Salisbury Fire Station	7/24/2018	<0.0189	-	-	-	-	-	-	-	0.0072	0.0148
Air	Salisbury Fire Station	8/1/2018	<0.0147	-	-	-	-	-	-	-	0.0072	0.0185
Air	Salisbury Fire Station	8/7/2018	<0.0194	-	-	-	-	-	-	-	0.0114	0.0305
Air	Salisbury Fire Station	8/14/2018	<0.0200	-	-	-	-	-	-	-	0.0105	0.0291
Air	Salisbury Fire Station quarterly composite	8/15/2018	-	0.124	0.269	<0.0008	<0.004	<0.0007	<0.002	<0.0007	-	-
Air	Salisbury Fire Station	8/22/2018	<0.0202	-	-	-	-	-	-	-	0.0083	0.0229
Air	Salisbury Fire Station	8/28/2018	<0.0208	-	-	-	-	-	-	-	0.0110	0.0295
Air	Salisbury Fire Station	9/4/2018	<0.0153	-	-	-	-	-	-	-	0.0086	0.0303
Air	Salisbury Fire Station	9/12/2018	<0.0168	-	-	-	-	-	-	-	0.0066	0.0196
Air	Salisbury Fire Station	9/19/2018	<0.0208	-	-	-	-	-	-	-	0.0076	0.0213
Air	Salisbury Fire Station	9/25/2018	<0.0185	-	-	-	-	-	-	-	0.0074	0.0214
Air	Salisbury Fire Station	10/3/2018	<0.0150	-	-	-	-	-	-	-	0.0069	0.0187
Air	Salisbury Fire Station	10/10/2018	<0.0245	-	-	-	-	-	-	-	0.0090	0.0235
Air	Salisbury Fire Station	10/16/2018	<0.0252	-	-	-	-	-	-	-	0.0082	0.0238
Air	Salisbury Fire Station	10/24/2018	<0.0177	-	-	-	-	-	-	-	0.0070	0.0186
Air	Salisbury Fire Station	10/31/2018	<0.0218	-	-	-	-	-	-	-	0.0091	0.0173
Air	Salisbury Fire Station	11/7/2018	<0.0278	-	-	-	-	-	-	-	0.0060	0.0202
Air	Salisbury Fire Station	11/13/2018	<0.0200	-	-	-	-	-	-	-	0.0084	0.0231
Air	Salisbury Fire Station quarterly composite	11/15/2018	-	0.082	0.010	<0.0007	<0.002	<0.0006	<0.002	<0.0006	-	-
Air	Salisbury Fire Station	11/21/2018	<0.0143	-	-	-	-	-	-	-	0.0058	0.0209
Air	Salisbury Fire Station	11/28/2018	<0.0191	-	-	-	-	-	-	-	0.0078	0.0240
Air	Salisbury Fire Station	12/6/2018	<0.0238	-	-	-	-	-	-	-	0.0063	0.0169
Air	Salisbury Fire Station	12/14/2018	<0.0182	-	-	-	-	-	-	-	0.0063	0.0293
Air	Salisbury Fire Station	12/19/2018	<0.0282	-	-	-	-	-	-	-	0.0102	0.0364
Air	Salisbury Fire Station	12/26/2018	<0.0187	-	-	-	-	-	-	-	0.0050	0.0290
Air	Background	1/2/2018	<0.0185	-	-	-	-	-	-	-	0.0088	0.0326
Air	Background	1/9/2018	<0.0199	-	-	-	-	-	-	-	0.0089	0.0311
Air	Background	1/16/2018	<0.0156	-	-	-	-	-	-	-	0.0101	0.0353
Air	Background	1/22/2018	<0.0219	-	-	-	-	-	-	-	0.0110	0.0489
Air	Background	1/29/2018	<0.0184	-	-	-	-	-	-	-	0.0114	0.0313
Air	Background	2/5/2018	<0.0175	-	-	-	-	-	-	-	0.0095	0.0270
Air	Background	2/12/2018	<0.0187	-	-	-	-	-	-	-	0.0100	0.0300

- " = Not analyzed

* I-131 = iodine 131; Ba-140 = Barium 140; Be-7 = Beryllium 7; Co-60 = Cobalt 60; Cs-134 = Cesium 134; Cs-137=Cesium 137; Fe-59=Iron 59; H-3* = Tritium; K-40 = Potassium-40; Mn-54=Manganese-54; Zn-65=Zinc 65

"<" = value is less than the listed MDA (Minimum Detectable Activity) value

"NR"= result not reported for quality control reasons.

Sample Type	Location	Date	I-131* (pCi/m ³)	Be-7* (pCi/m ³)	K-40* (pCi/m ³)	Mn-54* (pCi/m ³)	Fe-59* (pCi/m ³)	Co-60* (pCi/m ³)	Zn-65* (pCi/m ³)	Cs-137* (pCi/m ³)	Gross Alpha (pCi/m ³)	Gross Beta (pCi/m ³)
Air	Background quarterly composite	2/15/2018	-	0.099	0.280	<0.0008	<0.003	<0.0006	<0.002	<0.0006	-	-
Air	Background	2/20/2018	<0.0139	-	-	-	-	-	-	-	0.0070	0.0345
Air	Background	2/26/2018	<0.0152	-	-	-	-	-	-	-	0.0061	0.0226
Air	Background	3/5/2018	<0.0120	-	-	-	-	-	-	-	0.0049	0.0259
Air	Background	3/12/2018	<0.0172	-	-	-	-	-	-	-	0.0041	0.0152
Air	Background	3/19/2018	<0.0172	-	-	-	-	-	-	-	0.0063	0.0257
Air	Background	3/26/2018	<0.0184	-	-	-	-	-	-	-	0.0043	0.0199
Air	Background	4/2/2018	<0.0178	-	-	-	-	-	-	-	0.0069	0.0253
Air	Background	4/9/2018	<0.0174	-	-	-	-	-	-	-	0.0053	0.0264
Air	Background	4/17/2018	<0.0160	-	-	-	-	-	-	-	0.0057	0.0252
Air	Background	4/23/2018	<0.0202	-	-	-	-	-	-	-	0.0038	0.0204
Air	Background	4/30/2018	<0.0188	-	-	-	-	-	-	-	0.0044	0.0203
Air	Background	5/7/2018	<0.0173	-	-	-	-	-	-	-	0.0059	0.0292
Air	Background	5/14/2018	<0.0180	-	-	-	-	-	-	-	0.0040	0.0187
Air	Background quarterly composite	5/15/2018	-	0.124	<0.025	<0.0008	<0.004	<0.0007	<0.002	<0.0007	-	-
Air	Background	5/21/2018	<0.0164	-	-	-	-	-	-	-	0.0040	0.0176
Air	Background	5/29/2018	<0.0111	-	-	-	-	-	-	-	0.0034	0.0234
Air	Background	6/4/2018	<0.0190	-	-	-	-	-	-	-	0.0045	0.0197
Air	Background	6/11/2018	<0.0134	-	-	-	-	-	-	-	0.0033	0.0189
Air	Background	6/18/2018	<0.0165	-	-	-	-	-	-	-	0.0026	0.0151
Air	Background	6/25/2018	<0.0173	-	-	-	-	-	-	-	0.0048	0.0254
Air	Background	7/2/2018	<0.0172	-	-	-	-	-	-	-	0.0070	0.0250
Air	Background	7/9/2018	<0.0182	-	-	-	-	-	-	-	0.0081	0.0276
Air	Background	7/16/2018	<0.0178	-	-	-	-	-	-	-	0.0060	0.0257
Air	Background	7/23/2018	<0.0179	-	-	-	-	-	-	-	0.0062	0.0224
Air	Background	7/30/2018	<0.0167	-	-	-	-	-	-	-	0.0046	0.0198
Air	Background	8/6/2018	<0.0183	-	-	-	-	-	-	-	0.0066	0.0234
Air	Background	8/13/2018	<0.0153	-	-	-	-	-	-	-	0.0075	0.0408
Air	Background quarterly composite	8/15/2018	-	0.109	<0.024	<0.001	<0.003	<0.001	<0.002	<0.0005	-	-
Air	Background	8/20/2018	<0.0186	-	-	-	-	-	-	-	0.0072	0.0290
Air	Background	8/27/2018	<0.0180	-	-	-	-	-	-	-	0.0055	0.0263

- " = Not analyzed

* I-131 = iodine 131; Ba-140 = Barium 140; Be-7 = Beryllium 7; Co-60 = Cobalt 60; Cs-134 = Cesium 134; Cs-137=Cesium 137; Fe-59=Iron 59; H-3* = Tritium; K-40 = Potassium-40; Mn-54=Manganese-54; Zn-65=Zinc 65

"<" = value is less than the listed MDA (Minimum Detectable Activity) value

"NR"= result not reported for quality control reasons.

Sample Type	Location	Date	I-131* (pCi/m ³)	Be-7* (pCi/m ³)	K-40* (pCi/m ³)	Mn-54* (pCi/m ³)	Fe-59* (pCi/m ³)	Co-60* (pCi/m ³)	Zn-65* (pCi/m ³)	Cs-137* (pCi/m ³)	Gross Alpha (pCi/m ³)	Gross Beta (pCi/m ³)
Air	Background	9/4/2018	<0.0165	-	-	-	-	-	-	-	0.0066	0.0364
Air	Background	9/10/2018	<0.0222	-	-	-	-	-	-	-	0.0048	0.0278
Air	Background	9/17/2018	<0.0178	-	-	-	-	-	-	-	0.0038	0.0222
Air	Background	9/24/2018	<0.0176	-	-	-	-	-	-	-	0.0029	0.0203
Air	Background	10/1/2018	<0.0135	-	-	-	-	-	-	-	0.0061	0.0221
Air	Background	10/9/2018	<0.0126	-	-	-	-	-	-	-	0.0076	0.0225
Air	Background	10/15/2018	<0.0223	-	-	-	-	-	-	-	0.0095	0.0277
Air	Background	10/22/2018	<0.0153	-	-	-	-	-	-	-	0.0099	0.0279
Air	Background	10/29/2018	<0.0138	-	-	-	-	-	-	-	0.0080	0.0230
Air	Background	11/5/2018	<0.0177	-	-	-	-	-	-	-	0.0087	0.0290
Air	Background	11/13/2018	<0.0146	-	-	-	-	-	-	-	0.0064	0.0251
Air	Background quarterly composite	11/15/2018	-	0.089	0.147	<0.0008	<0.003	<0.0008	<0.002	<0.0007	-	-
Air	Background	11/19/2018	<0.0194	-	-	-	-	-	-	-	0.0095	0.0268
Air	Background	11/26/2018	<0.0199	-	-	-	-	-	-	-	0.0073	0.0262
Air	Background	12/3/2018	<0.0176	-	-	-	-	-	-	-	0.0088	0.0179
Air	Background	12/10/2018	<0.0190	-	-	-	-	-	-	-	0.0076	0.0303
Air	Background	12/17/2018	<0.0179	-	-	-	-	-	-	-	0.0035	0.0333
Air	Background	12/24/2018	<0.0153	-	-	-	-	-	-	-	0.0033	0.0265
Air	Background	12/31/2018	<0.0140	-	-	-	-	-	-	-	0.0034	0.0245

- " = Not analyzed

* I-131 = iodine 131; Ba-140 = Barium 140; Be-7 = Beryllium 7; Co-60 = Cobalt 60; Cs-134 = Cesium 134; Cs-137=Cesium 137; Fe-59=Iron 59; H-3* = Tritium; K-40 = Potassium-40; Mn-54=Manganese-54; Zn-65=Zinc 65

"<" = value is less than the listed MDA (Minimum Detectable Activity) value

"NR"= result not reported for quality control reasons.

Table 6. Seabrook Nuclear Power Station 2018 Environmental Monitoring Data – Liquid Matrices

Sample Type	Location	Date	K-40* (pCi/L)	Mn-54* (pCi/L)	Fe-59* (pCi/L)	Co-60* (pCi/L)	Zn-65* (pCi/L)	I-131* (pCi/L)	Cs-134* (pCi/L)	Cs-137* (pCi/L)	Ba-140* (pCi/L)	H-3* (pCi/L)
Surface water	Ipswich bay ¹	1/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Ipswich bay ¹	1/19/2018	220	<3.0	<9.3	<2.7	<6.7	NR	-	<2.9	-	-
Surface water	Ipswich bay ¹	2/15/2018	345	<3.3	<8.2	<2.8	<6.5	<27.3	-	<2.8	-	<300
Surface water	Ipswich bay ¹	3/15/2018	280	<3.3	<8.7	<2.7	<6.6	NR	-	<2.9	-	<300
Surface water	Ipswich bay ¹	4/11/2018	274	<2.5	<5.2	<2.3	<5.0	<4.0	-	<2.6	-	<300
Surface water	Ipswich bay ¹	5/14/2018	278	<2.4	<5.6	<2.6	<5.2	<10.0	-	<2.6	-	<300
Surface water	Ipswich bay ¹	6/12/2018	326	<2.4	<4.7	<2.6	<5.3	<5.1	-	<2.5	-	<300
Surface water	Ipswich bay ¹	7/10/2018	919	<3.4	<8.1	<3.3	<7.1	<18.3	-	<3.2	-	-
Surface water	Ipswich bay ¹	7/15/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Ipswich bay ¹	8/14/2018	263	<3.4	<7.5	<3.1	<6.7	<14.7	-	<2.9	-	<300
Surface water	Ipswich bay ¹	9/12/2018	313	<2.7	<7.9	<2.6	<5.8	NR	-	<2.6	-	-
Surface water	Ipswich bay ¹	9/14/2018	-	-	-	-	-	-	-	-	-	<300
Surface water	Ipswich bay ¹	10/17/2018	357	<3.2	<8.0	<2.8	<6.8	<23.5	-	<3.2	-	<300
Surface water	Ipswich bay ¹	11/12/2018	266	<3.2	<7.9	<2.9	<6.6	<21.9	-	<2.9	-	<300
Surface water	Ipswich bay ¹	12/4/2018	268	<3.3	<10.7	<2.8	<7.3	NR	-	<3.0	-	<300
Milk	Rowley	1/3/2018	1310	-	-	-	-	<2.7	<2.6	<3.2	<11.1	
Milk	Rowley	2/7/2018	1380	-	-	-	-	<3.5	<2.8	<3.2	<11.3	
Milk	Rowley	3/7/2018	1390	-	-	-	-	<2.6	<2.5	<3.0	<10.0	
Milk	Rowley	4/11/2018	1360	-	-	-	-	<3.2	<2.7	<3.0	<10.1	
Milk	Rowley	5/2/2018	2550	-	-	-	-	<3.3	<2.7	<2.9	<10.3	
Milk	Rowley	6/5/2018	1420	-	-	-	-	<3.3	<2.7	<3.1	<10.3	
Milk	Rowley	7/11/2018	1450	-	-	-	-	<2.5	<2.3	<3.1	<9.5	
Milk	Rowley	8/1/2018	1870	-	-	-	-	<2.5	<2.5	<3.1	<9.6	
Milk	Rowley	9/4/2018	948	-	-	-	-	<2.9	<3.0	<3.5	<11.5	
Milk	Rowley	10/3/2018	1470	-	-	-	-	<3.4	<2.6	<2.9	<10.5	
Milk	Rowley	11/14/2018	1400	-	-	-	-	<3.3	<2.6	<3.1	<10.5	
Milk	Rowley	12/19/2018	1420	-	-	-	-	<3.2	<2.5	<3.0	<10.2	

(1) Background sample

-" = Not analyzed

* I-131 = iodine 131; Ba-140 = Barium 140; Be-7 = Beryllium 7; Co-60 = Cobalt 60; Cs-134 = Cesium 134; Cs-137=Cesium 137; Fe-59=Iron 59; H-3* = Tritium; K-40 = Potassium-40; Mn-54=Manganese-54; Zn-65=Zinc 65

"<" = value is less than the listed MDA (Minimum Detectable Activity) value

"NR"= result not reported for quality control reasons.

Table 7. Seabrook Nuclear Power Station 2018 Environmental Monitoring Data –Solid Matrices

Sample	Location	Date	Be-7* (pCi/kg)	K-40* (pCi/kg)	Mn-54* (pCi/kg)	Fe-59* (pCi/kg)	Co-60* (pCi/kg)	Zn-65* (pCi/kg)	Cs-137* (pCi/kg)	I-131* (pCi/kg)
Eel Pout	Ipswich Bay (background)	5/18/2018	<298	3160	<21.8	<72.4	<23.1	<49.3	<21.7	-
Flounder	Ipswich Bay (background)	8/14/2018	<913	3410	<31.2	<290	<25.8	<71.9	<24.2	-
Lobster	Ipswich Bay (background)	5/21/2018	<50.2	2480	<4.6	<14.0	<4.1	<10.2	<4.0	-
Lobster	Ipswich Bay (background)	11/18/2018	<46.4	2510	<4.9	<14.5	<4.7	<10.9	<4.4	-
Modiolus ¹	Ipswich Bay (background)	5/14/2018	31	1360	<3.5	<9.5	<4.0	<7.9	<3.6	-
Modiolus ¹	Ipswich Bay (background)	11/12/2018	14	1070	<2.8	<6.6	<3.0	<6.0	<3.1	-
Mytilus ¹	Ipswich Bay (background)	5/17/2018	51	1340	<4.5	<10.3	<4.5	<9.2	<3.9	-
Mytilus ¹	Ipswich Bay (background)	11/13/2018	136	1390	<3.7	<8.8	<4.2	<8.7	<3.8	-
Sediment	Ipswich Bay - beach (background)	5/17/2018	-	16,400	-	-	<23.2	-	<21.1	-
Sediment	Ipswich Bay - beach (background)	11/14/2018	-	17,500	-	-	<22.8	-	<22.7	-
Sediment	Ipswich Bay - subtidal (background)	5/14/2018	-	12,400	-	-	<25.6	-	<26.8	-
Sediment	Ipswich Bay - subtidal (background)	11/12/2018	-	13,400	-	-	<24.2	-	<26.7	-
Irish Moss	Ipswich Bay (background)	5/14/2018	1,050	31500	<15.5	<40.5	<15.3	<40.7	<14.5	194
Irish Moss	Ipswich Bay (background)	11/12/2018	3910	18300	<13.2	<32.7	<12.1	<30.9	<12.7	<53.4
Green Beans	Bartlett Farm, Salisbury	7/17/2018	<41.8	2560	<5.2	<9.97	<6.3	<11.6	<5.4	-
Strawberries	Bartlett Farm, Salisbury	6/19/2018	37	1590	<3.8	<7.9	<4.7	<9.0	<3.9	-
Strawberries	Russell Orchards, Ipswich (background)	6/19/2018	<36.8	1310	<5.3	<11.1	<5.1	<11.2	<4.9	-
Tomatoes	Russell Orchards, Ipswich (background)	8/14/2018	<29.7	2040	<3.6	<7.37	<4.1	<8.3	<3.7	-
Tomatoes	Bartlett Farm, Salisbury	8/14/2018	<32.1	1900	<4.0	<8.77	<4.2	<9.6	<3.9	-
Zucchini	Russell Orchards, Ipswich (background)	7/17/2018	<40.0	1670	<4.7	<8.85	<4.6	<10.7	<4.4	-

¹Mytilus (i.e., blue or common mussel) samples collected on Plum Island; Modiolus (i.e., Atlantic ribbed mussel) samples collected offshore.

- = Not analyzed

* I-131 = iodine 131; Ba-140 = Barium 140; Be-7 = Beryllium 7; Co-60 = Cobalt 60; Cs-134 = Cesium 134; Cs-137=Cesium 137; Fe-59=Iron 59; H-3* = Tritium; K-40 = Potassium-40; Mn-54=Manganese-54; Zn-65=Zinc 65

"<" = value is less than the listed MDA (Minimum Detectable Activity) value

"NR"= result not reported for quality control reasons.