




Massachusetts 2020 Air Quality Report

June 2021



Department of Environmental Protection
Bureau of Air and Waste
Division of Air and Climate Programs
Air Assessment Branch
Wall Experiment Station
37 Shattuck Street
Lawrence, Massachusetts 01843

ACKNOWLEDGEMENTS

This 2020 Air Quality Report was prepared by the Massachusetts Department of Environmental Protection (MassDEP), Air Assessment Branch (AAB), which collects representative samples of ambient air for a number of pollutants at monitoring stations located across the Commonwealth. All samples are collected in a precise and scientifically sound manner in order to properly characterize the quality of the air in the Commonwealth.

The photo on the cover is a view of the Worcester Airport monitoring station in Worcester, MA.

This report is available on MassDEP's web site at

www.mass.gov/eea/agencies/massdep/air/quality/air-monitoring-reports-and-studies.html

Questions about this report may be directed to:

Sean Dunn
Air Assessment Branch
Wall Experiment Station
Lawrence, MA 01843-1343
(978) 242-1335
email: sean.m.dunn@mass.gov

Table of Contents

Section 1 – Ambient Air Monitoring Program	5
1.1 - Program Overview	5
1.2 - National Ambient Air Quality Standards (NAAQS)	8
1.3 - Pollutant Health Effects and Sources	9
1.4 - Monitoring Network Description	11
Section 2 – Attainment of Air Quality Standards.....	12
2.1 - Attainment Status Summary	12
2.2 - 2020 Ozone Season	13
2.3 - Daily Ozone and PM Forecasts	16
Section 3 – Massachusetts Air Quality Data Summaries	17
3.1 - Ozone Summary	17
3.2 - Sulfur Dioxide (SO ₂) Summary	21
3.3 - Nitrogen Dioxide (NO ₂) Summary.....	24
3.4 - Carbon Monoxide (CO) Summary.....	27
3.5 - Particulate Matter 10 Microns (PM ₁₀) Summary.....	30
3.6 - Particulate Matter 2.5 Microns (PM _{2.5}) Summary	32
3.7 - Speciation	38
3.8 - Interagency Monitoring of Protected Visual Environments (IMPROVE).....	38
3.9 - Quality Assurance and Quality Control (QA/QC).....	38
Section 4 – PAMS/Air Toxics Monitoring.....	39
4.1 - Photochemical Assessment Monitoring Stations (PAMS) Monitoring.....	39
4.2 - Air Toxics Monitoring.....	39

List of Abbreviations

AAB.....	Air Assessment Branch
AQS.....	Air Quality System
AQI.....	Air Quality Index
BC	Black Carbon
BP	Barometric Pressure
CAA.....	Clean Air Act
CFR	Code of Federal Regulations
CO.....	Carbon Monoxide
CO ₂	Carbon Dioxide
FEM	Federal Equivalent Method
FRM	Federal Reference Method
EPA	United States Environmental Protection Agency
IMPROVE	Interagency Monitoring of Protected Visual Environments
MassDEP.....	Massachusetts Department of Environmental Protection
NAAQS.....	National Ambient Air Quality Standards
NATTS.....	National Air Toxics Trends Station
NCore	National Core Monitoring Network
NO	Nitric Oxide
NO _x	Nitrogen Oxides
NO _y	Total Reactive Oxidized Nitrogen
NO ₂	Nitrogen Dioxide
NO ₃	Nitrate
O ₃	Ozone
PAH.....	Polycyclic Aromatic Hydrocarbon
PAMS.....	Photochemical Assessment Monitoring Stations
Pb	Lead
ppb	parts per billion by volume
ppm	parts per million by volume
PM _{2.5}	Particulate matter ≤ 2.5 microns aerodynamic diameter
PM ₁₀	Particulate matter ≤ 10 microns aerodynamic diameter
PRECIP	Precipitation
QA/QC	Quality Assurance and Quality Control
REL.....	Relative Humidity
SIP.....	State Implementation Plan
SO ₂	Sulfur Dioxide
SO ₄	Sulfate
SUN.....	Solar Radiation
TEMP	Temperature
TOTAL UV	Ultraviolet radiation
TSA	Technical Systems Audit
µg/m ³	micrograms per cubic meter
VOCs	Volatile Organic Compounds
WS/WD.....	Wind Speed/Wind Direction
WSv/WDv	Wind Speed Vector/Wind Direction Vector

Section 1 – Ambient Air Monitoring Program

1.1 - Program Overview

Introduction

The Massachusetts Department of Environmental Protection (MassDEP) is responsible for monitoring outdoor air quality in Massachusetts, and for developing plans and regulatory programs to reduce emissions of pollutants that adversely affect public health, welfare, and the environment.

MassDEP's Air Assessment Branch (AAB) operates a network of air monitoring stations throughout the Commonwealth. During 2020, MassDEP operated 22 monitoring stations located in 18 cities and towns. MassDEP also received data from the Wampanoag Tribe of Gay Head (Aquinnah), which operates an air monitoring station on Martha's Vineyard.

MassDEP submits ambient air quality data to the national Air Quality System (AQS) database that is administered by the U.S. Environmental Protection Agency (EPA). Continuous monitoring data is sent to the AirNow website, <https://www.airnow.gov/>, which reports data from all U.S. monitoring stations. MassDEP's MassAir Online website provides air quality information and allows users to click on a map of the state to find current air quality data from the MassDEP continuous air monitoring network. MassAir Online is found at www.mass.gov/eea/agencies/massdep/air/quality/. The EPA also makes historical AQS data for all U.S. monitoring stations available at <https://www.epa.gov/outdoor-air-quality-data>.

Why is Air Quality Data Collected?

Ambient air quality data is used for a number of purposes, including to:

- Provide information about air quality to the public;
- Provide short-term and long-term information regarding air pollution and public health;
- Verify compliance with National Ambient Air Quality Standards (NAAQS);
- Assess the effectiveness of current air pollution control regulations and programs;
- Support development of policies and regulations aimed at reducing air pollution;
- Support long-term trend analysis and special research; and
- Fulfill requirements to report ambient air quality data to EPA.

What is Monitored?

MassDEP monitors parameters in the following categories:

Criteria pollutants for which EPA has established NAAQS. The criteria pollutants monitored are:

- sulfur dioxide (SO₂)
- ozone (O₃)
- carbon monoxide (CO)
- nitrogen dioxide (NO₂)
- lead (Pb)

- particulate matter ≤ 10 microns (PM_{10})
- particulate matter ≤ 2.5 microns ($PM_{2.5}$)

Non-criteria pollutants do not have NAAQS but can contribute to the formation of ozone and particulate matter and/or be toxic. The non-criteria pollutants monitored include:

- nitric oxide (NO)
- total nitrogen oxides (NO_x)
- total reactive oxidized nitrogen (NO_y)
- volatile organic compounds (VOCs) – ozone precursors and reaction product chemicals
- black carbon (i.e., soot)
- toxics – health-relevant VOCs, semi-volatile organic compounds (SVOCs), carbonyls and metals

Meteorological parameters monitored include:

- wind speed/wind direction (WS/WD) and vector (WSv/WDv)
- relative humidity (REL)
- temperature (TEMP)
- barometric pressure (BP)
- solar radiation (SUN)
- precipitation (PRECIP)
- ultraviolet radiation (TOTAL UV)


Monitoring Station Locations

Monitoring stations are sited to provide data for various purposes. Some are located where maximum pollutant concentrations are expected, while others are positioned in areas that will provide data that is representative of larger geographical areas. Local topography and pollutant source areas are factors that determine how well a particular monitor's location will represent a region.

Networks of monitors are located throughout the state. These networks are designed to reflect pollutant concentrations for all of Massachusetts. Section III of this report contains data summaries for each pollutant measured and maps showing the monitor locations for each network. Appendix A contains a list of monitoring stations.

The map on page 3 shows Massachusetts cities and towns where air monitors were located during 2020.

2020 Air Monitoring Network



1.2 - National Ambient Air Quality Standards (NAAQS)

Below are the current NAAQS for criteria pollutants set by EPA. Primary Standards are designed to protect public health, including the health of sensitive populations such as asthmatics, children, and the elderly. Secondary Standards are designed to protect public welfare, including protection against decreased visibility, damage to crops, vegetation, and buildings.

National Ambient Air Quality Standards					
Pollutant		Primary/ Secondary	Averaging Time	Level	Form
Carbon Monoxide		primary	8-hour	9 ppm	Not to be exceeded more than once per year
			1-hour	35 ppm	
Lead		primary and secondary	Rolling 3-month average	0.15 $\mu\text{g}/\text{m}^3$	Not to be exceeded
Nitrogen Dioxide		primary	1-hour	100 ppb	98 th percentile of 1-hr daily maximum concentrations, averaged over 3 years
		primary and secondary	Annual	53 ppb	Annual Mean
Ozone		primary and secondary	8-hour	0.070 ppm	Annual fourth-highest daily maximum 8-hr concentration, averaged over 3 years
Particle Pollution	PM _{2.5}	primary	Annual	12 $\mu\text{g}/\text{m}^3$	annual mean, averaged over 3 years
		secondary	Annual	15 $\mu\text{g}/\text{m}^3$	annual mean, averaged over 3 years
		primary and secondary	24-hour	35 $\mu\text{g}/\text{m}^3$	98 th percentile, averaged over 3 years
	PM ₁₀	primary and secondary	24-hour	150 $\mu\text{g}/\text{m}^3$	Not to be exceeded more than once per year on average over 3 years
Sulfur Dioxide		primary	1-hour	75 ppb	99 th percentile of 1-hr daily maximum concentrations, averaged over 3 years
		secondary	3-hour	0.5 ppm	Not to be exceeded more than once per year

$\mu\text{g}/\text{m}^3$ = micrograms per cubic meter

ppm = parts per million

ppb = parts per billion

1.3 - Pollutant Health Effects and Sources

Ozone (O₃)

- Tropospheric O₃ (ground-level) and Stratospheric O₃ (upper atmosphere) are the same chemical compound, just found at different places in the atmosphere. Stratospheric O₃, found at greater than 30,000 feet above the surface of the earth, is beneficial to all life because it filters out the sun's harmful UV radiation before it reaches the earth's surface. Ground-Level O₃ on the other hand is a health and environmental problem. This report pertains exclusively to ground-level O₃.
- O₃ is a respiratory irritant that can reduce lung function and resistance to infection. It can cause asthma attacks, nasal congestion, and throat irritation. It can inflame and damage (possibly permanently) cells that line the lungs and aggravate chronic lung diseases. In addition, several studies have found a strong link between increases in ground-level O₃ and increased risk of premature death.
- O₃ is toxic to vegetation, inhibiting growth and causing leaf damage.
- O₃ deteriorates materials such as rubber and fabrics.
- Ground-level O₃ is unique in that it is formed by the reactions that occur between certain precursor pollutants in the presence of intense, high-energy sunlight during the hot summer months. The complexity of the process and the amount of time needed to complete these reactions can result in the buildup of ground-level ozone concentrations far downwind from the original source of the precursors.
- Sources of ground-level O₃ precursors, i.e., nitrogen oxides and hydrocarbons, include motor vehicles, lawn and garden equipment, power plants and other industrial sources.

Carbon Monoxide (CO)

- CO binds with hemoglobin in the blood, reducing the amount of oxygen carried to organs and tissues.
- Symptoms of high CO exposure include shortness of breath, chest pain, headaches, confusion, and loss of coordination. The health threat is most severe for those with cardiovascular disease.
- Motor vehicle emissions are the largest source of CO, which is produced from incomplete combustion of carbon in fuels.
- Industrial processes and non-transportation fuel combustion (e.g., boilers, lawn and garden equipment) also are sources of CO.

Sulfur Dioxide (SO₂)

- SO₂ combines with water vapor to form acidic aerosols harmful to the respiratory tract, aggravating symptoms associated with lung diseases such as asthma and bronchitis.
- SO₂ is a primary contributor to acid deposition. Impacts of acid deposition include: acidification of lakes and streams, damage to vegetation, damage to materials, and diminution of visibility.
- SO₂ is a product of fuel combustion (e.g., the burning of coal and oil that contains sulfur). Sources include power plants and business and residential sources burning heating oil.

Nitrogen Dioxide (NO₂)

- NO₂ lowers resistance to respiratory infections and aggravates symptoms associated with asthma and bronchitis.
- NO₂ contributes to acid deposition. Impacts of acid deposition include acidification of lakes and streams, damage to vegetation, damage to materials, and diminution of visibility.
- NO₂ and nitric oxide (NO) contribute to the formation of ozone.
- NO₂ is formed from the oxidation of NO. Major sources of NO are fuel combustion, space heating, power plants and motor vehicles.

Particulate Matter (PM₁₀ and PM_{2.5})

- Particulate matter is tiny airborne particles or aerosols, which include dust, dirt, soot, smoke, and liquid droplets. Fine particulate matter (mostly below 2.5 microns in size) are not only the result of direct emissions but can be formed in the atmosphere by chemical reactions involving gaseous pollutants.
- The numbers 2.5 and 10 refer to the particle size (equal to or less than that size), measured in microns, which are collected by the monitors. Several thousand PM_{2.5} particles could fit on the period at the end of this sentence.
- The small size of these particles allows easy entry into the human respiratory system. Long-term exposure causes the particles to accumulate in the lungs and affects breathing and produces respiratory symptoms. The small particles can migrate through the lungs and into the circulatory system and potentially produce cardio-vascular symptoms, as well as impacts from toxic components contained in the particulate matter.
- Particulate matter causes soiling and corrosion of materials.
- Particulate matter contributes to atmospheric haze that degrades visibility.
- Sources of particulates include industrial process emissions, motor vehicles, incinerators, power plants, and other fuel combustion sources.

Lead (Pb)

- Lead is an elemental metal that is found in nature.
- Exposure to lead can occur by inhalation or ingestion with food, water, soil or dust particles.
- Children, infants, and fetuses are the most susceptible to the effects of lead exposure.
- Lead causes intellectual disability, brain damage, and liver disease. It may be a factor in high blood pressure and damages the nervous system.
- Lead enters the atmosphere from the incineration of lead containing materials and from the manufacture and processing of lead containing products or materials like storage batteries, smelting and removal of lead-containing paint.

1.4 - Monitoring Network Description

The following describes the ambient air monitoring network in 2020.

Network Size	<ul style="list-style-type: none"> ▪ 23 Monitoring Stations (22 operated by MassDEP and 1 by the Wampanoag Tribe) ▪ 19 cities and towns with monitoring stations
Number of Continuous Monitors	<p>Continuous monitors measure air quality 24 hours per day. The data are reported as hourly averages.</p> <ul style="list-style-type: none"> ▪ Criteria pollutant monitors measure pollutants for which NAAQS have been set. <ul style="list-style-type: none"> ❑ 3 – Trace-level CO monitors ❑ 10 – NO₂, NO and NO_x monitors ❑ 17 – O₃ monitors (1 Tribal) ❑ 6 – Trace-level SO₂ monitors ❑ 17 – Continuous PM_{2.5} (1 collocated) ▪ Meteorological monitors track weather conditions. <ul style="list-style-type: none"> ❑ 13 – Barometric pressure ❑ 13 – Relative humidity ❑ 13 – Solar radiation ❑ 13 – Temperature ❑ 13 – Wind speed/wind direction ❑ 1 – Wind speed vector/wind direction vector ❑ 2 – Precipitation ❑ 1 – Ultraviolet radiation ▪ Other Monitors <ul style="list-style-type: none"> ❑ 3 – Total Reactive Oxidized Nitrogen (NO/NO_y) ❑ 1 – Photochemical assessment monitoring station (PAMS). PAMS monitors measure VOCs using an automated gas chromatograph (auto-GC) on an hourly basis during the summer. ❑ 7 – Black Carbon
Number of Intermittent Monitors	<p>Intermittent monitors collect discrete samples for a specific time period. The samples are collected every day, every third day, or every sixth day. The data are averaged in 3-hour or 24-hour intervals.</p> <ul style="list-style-type: none"> ▪ Criteria pollutant monitors measure pollutants that have NAAQS. <ul style="list-style-type: none"> ❑ 4 – PM₁₀ monitors (1 collocated) ❑ 11 – PM_{2.5} Federal Reference Method monitors (1 collocated) ▪ Non-criteria pollutant monitors measure pollutants that do not have NAAQS. <ul style="list-style-type: none"> ❑ 3 – VOCs and carbonyls ❑ 2 – Speciation. These monitors measure for PM_{2.5}, nitrates, and organics ❑ 1 – PM₁₀ for metals analysis

Section 2 – Attainment of Air Quality Standards

2.1 - Attainment Status Summary

The federal Clean Air Act (CAA) contains timeframes and milestones for states to meet and maintain NAAQS for criteria pollutants, which include carbon monoxide, lead, nitrogen dioxide, ozone, particulate matter, and sulfur dioxide. EPA sets NAAQS at levels to protect public health and the environment. The EPA must review each NAAQS every five years and may update the standards based on new scientific information as well as establish new monitoring requirements. Each state is required to monitor the ambient air to determine whether it meets each standard. If monitoring shows that the air quality does not meet a standard, the state must develop and implement pollution control strategies to attain that standard. Once air quality meets a standard, a state must develop a plan to maintain that standard while accounting for future economic and emissions growth. Taken together, these plans and control strategies constitute the State Implementation Plan (SIP).

Carbon Monoxide

Monitored levels of CO in Massachusetts meet the CO standards. Prior to the mid-1980s, Massachusetts was in nonattainment of the CO standards. However, with the adoption of numerous control programs, CO emissions significantly decreased and monitored levels of CO met the standards beginning in 1987. Massachusetts is designated as Unclassifiable/Attainment for the CO standards. Based on EPA's most recent review of the CO standards, in August 2011 EPA retained the existing primary CO standards of 9 ppm measured over 8 hours, and 35 ppm measured over 1 hour. Since EPA did not change the standards, no new designation process was triggered.

Lead

Monitored levels of lead in Massachusetts meet the lead standards. In October 2009, EPA lowered the lead standards (primary and secondary) from 1.5 $\mu\text{g}/\text{m}^3$ to 0.15 $\mu\text{g}/\text{m}^3$ averaged over a rolling 3-month period. In November 2011, EPA designated all of Massachusetts as Unclassifiable/Attainment for the 2009 standard. Based on EPA's most recent review of the lead standards, in September 2016 EPA retained the existing lead standards. Since EPA did not change the standards, no new designation process was triggered.

Nitrogen Dioxide

Monitored levels of NO₂ in Massachusetts meet the NO₂ standards. Based on EPA's most recent review of the NO₂ standards, in January 2010 EPA established a new 1-hour NO₂ standard of 100 ppb and new near-road monitoring requirements. In January 2012, EPA designated all of Massachusetts as Unclassifiable/Attainment for the 2010 standard.

Sulfur Dioxide

Monitored levels of SO₂ in Massachusetts meet the SO₂ standards. Based on EPA's most recent review of the SO₂ standards, in June 2010 EPA established a new 1-hour SO₂ standard of 75 ppb. In December 2018, EPA designated all of Massachusetts as Unclassifiable/Attainment for the 2010 standard.

Particulate Matter

There are standards for two types of particulate matter: PM₁₀ and PM_{2.5}. Monitored levels of PM₁₀ and PM_{2.5} in Massachusetts meet the respective standards.

Based on EPA's most recent review of the PM_{2.5} standards, in December 2012 EPA lowered the primary annual PM_{2.5} standard to 12 µg/m³. In December 2014, EPA designated all of Massachusetts as Unclassifiable/Attainment for the 2012 standard.

Ozone

In 1979, EPA established an ozone standard (0.12 ppm) based on the maximum 1-hour ozone concentration that occurred each day during the ozone monitoring season. Massachusetts was designated as Nonattainment with this standard.

In 1997, EPA established new 8-hour ozone standards (0.08 ppm) that were designed to be more representative of exposure over time, rather than just the maximum concentration (the 1-hour standard was revoked in 2005). Massachusetts was designated as Nonattainment for these standards at that time. Through a combination of state and regional controls, Massachusetts' air quality attained the 1997 standards by the 2009 attainment deadline.

In 2009, EPA lowered the 8-hour ozone standards to 0.075 ppm. In April 2012, EPA designated Dukes County as Nonattainment for the 2009 ozone standards and designated the remainder of Massachusetts as Unclassifiable/Attainment. Dukes County attained the 2009 ozone standard by the 2015 attainment deadline.

Based on EPA's most recent review of the ozone standards, in October 2015, EPA lowered the 8-hour ozone standards to 0.070 ppm. EPA has designated all of Massachusetts as Unclassifiable/Attainment for the 2015 standard.

2.2 - 2020 Ozone Season

In 2020, there were three days when the 8-hour ozone standard of 0.070 ppm was exceeded in Massachusetts. Based on the most recent three years of data (2018–2020), no monitoring locations violated the 0.070 ppm standard.

While ozone concentrations have trended downward over the past several decades due to air pollution control programs, ozone concentrations vary each year due to varying weather patterns. In general, the chemical reactions that produce elevated ozone concentrations occur when high energy sunlight (present on hot Summer days) facilitates the react of ozone "precursor" pollutants – VOCs and NO_x, which results in ozone formation. Typically, Massachusetts ozone exceedances occur when a high-pressure area well south of New England creates a broad southwesterly airflow which contains precursors from the upwind coastal urban corridor to New England. This is where reactions in the atmosphere result in elevated levels of ozone. This typical pattern also moves slowly, promoting heat wave conditions that can last several days, allowing pollutants to build up.

Difference Between Ozone Exceedances and Violations

An ozone exceedance occurs when monitored ozone concentrations exceed the ozone NAAQS. Ozone monitoring data is collected as an hourly average of continuous data which is then used to determine the highest 8-hour average value for the day. An exceedance of the 8-hour standard is an 8-hour averaged value that is greater than 0.070 ppm. An ozone exceedance occurs when a monitor records ambient levels of ozone above the standard. Monitoring an ozone exceedance does not mean that a violation of the ozone standard has occurred, because a violation of an ozone standard (as opposed to an exceedance) is based on three-year averages of data at each monitor.

An ozone violation of the 8-hour standard is determined using the annual 4th-highest daily maximum eight-hour ozone value at each monitor. A violation requires a three-year average of the annual 4th-highest daily maximum eight-hour value that is greater than 0.070 ppm. In other words, the eight-hour values for each day during a year for a specific monitor are ranked from highest to lowest. Then, the 4th-highest value for three consecutive years is averaged. If the three-year average is greater than 0.070 ppm, a violation of the 8-hour standard has occurred at that monitoring site.

Exceedance Days and Total Exceedance Trends

Figure 1 shows the number of 8-hour exceedance days and the total number of exceedances for the past ten years. Note that years 2011-2015 show what exceedances would have been had the 0.070 ppm 8-hour standards been in effect. Figure 2 shows the most recent ozone design values (i.e., the 4th highest 8-hour O₃ value averaged over three years) relative to the 2015 ozone NAAQS. Monitoring sites with less than three years of data, and therefore no three-year averages, are not included in Figure 2.

Figure 1
8-hour Ozone Exceedance Days and Total Exceedances 2011-2020
 Exceedances Based on 0.070 ppm 8-hour Standard





Figure 2
Ozone Design Values (2018-2020)



2.3 - Daily Ozone and PM Forecasts

MassDEP provides the public with daily air quality forecasts for ozone from April through September and for fine particles all year-round using weather maps and meteorological factors to predict whether or not conditions will result in elevated pollution levels. The daily air quality forecasts are available from www.mass.gov/eea/agencies/massdep/air/quality/. EPA web sites that contain regional and national pollution forecasts using data that is provided by participating states are located at www.epa.gov/region01/airquality/forecast.html and <http://airnow.gov/>. The table below describes the ratings used in the daily air quality forecasts.

Air Quality Index (AQI): Ozone		
Index Values	Levels of Health Concern	Cautionary Statements
0-50	Good	None
51-100	Moderate	Unusually sensitive people should consider reducing prolonged or heavy exertion outdoors.
101-150	Unhealthy for Sensitive Groups	Active children and adults, and people with lung disease, such as asthma, should reduce prolonged or heavy exertion outdoors.
151-200	Unhealthy	Active children and adults, and people with lung disease, such as asthma, should avoid prolonged or heavy exertion outdoors. Everyone else, especially children, should reduce prolonged or heavy exertion outdoors.
201-300	Very Unhealthy	Active children and adults, and people with lung disease, such as asthma, should avoid all outdoor exertion. Everyone else, especially children, should avoid prolonged or heavy exertion outdoors.

Air Quality Index (AQI): Particle Pollution		
Index Values	Levels of Health Concern	Cautionary Statements
0-50	Good	None
51-100	Moderate	Unusually sensitive people should consider reducing prolonged or heavy exertion.
101-150	Unhealthy for Sensitive Groups	People with heart or lung disease, older adults, and children should reduce prolonged or heavy exertion.
151-200	Unhealthy	People with heart or lung disease, older adults, and children should avoid prolonged or heavy exertion. Everyone else should reduce prolonged or heavy exertion.
201-300	Very Unhealthy	People with heart or lung disease, older adults, and children should avoid all physical activity outdoors. Everyone else should avoid prolonged or heavy exertion.

Section 3 – Massachusetts Air Quality Data Summaries

3.1 - Ozone Summary

2020 Ozone Data Summary

A summary of the data collected during the 2020 ozone season (March 1 – September 30) is shown below (in parts per million). MassDEP operated 16 ozone monitors during 2020. The Wampanoag Tribe operated one ozone monitor in Aquinnah on Martha’s Vineyard.


O ₃ 2020			1 st	2 nd	3 rd	4 th	8-HR
CITY	COUNTY	ADDRESS	MAX 8-HR	MAX 8-HR	MAX 8-HR	MAX 8-HR	MAX>0.070 STD
Aquinnah (Tribal)	Dukes	Herring Creek Drive	0.078	0.065	0.061	0.060	1
Boston	Suffolk	Harrison Avenue	0.065	0.061	0.057	0.057	0
Brockton	Plymouth	Clinton Street	0.064	0.062	0.061	0.060	0
Chelmsford EPA	Middlesex	Technology Drive	0.068	0.067	0.062	0.059	0
Chelmsford NR	Middlesex	Manning Road	0.056	0.052	0.051	0.051	0
Chicopee	Hampden	Anderson Road	0.063	0.062	0.061	0.061	0
Fairhaven	Bristol	School Street	0.076	0.072	0.065	0.064	2
Fall River	Bristol	Globe Street	0.073	0.071	0.067	0.066	2
Greenfield	Franklin	Barr Avenue	0.060	0.055	0.052	0.052	0
Haverhill	Essex	Washington Street	0.069	0.063	0.060	0.056	0
Lynn	Essex	Parkland Avenue	0.068	0.061	0.061	0.059	0
Milton	Norfolk	Canton Avenue	0.057	0.056	0.054	0.053	0
Pittsfield	Berkshire	Silver Lake Blvd	0.058	0.058	0.056	0.054	0
Truro	Barnstable	Collins Road	0.073	0.071	0.063	0.063	2
Uxbridge	Worcester	E. Hartford Ave	0.065	0.063	0.063	0.059	0
Ware	Hampshire	Skyline Drive	0.065	0.062	0.062	0.061	0
Worcester	Worcester	Airport Drive	0.067	0.066	0.065	0.063	0

Standard: 8-hour = 0.070 ppm

1st, 2nd, 3rd, 4th MAX 8-HR = Maximum 8-hour Value for the 1st, 2nd, 3rd and 4th Highest Day

8-HR MAX > 0.070 STD = Number of Measured Daily 8-hour Maximum Values Greater Than the 0.070 ppm 8-hour Standard

Ozone Monitor Locations



8-hour Ozone Exceedance Trends

Figure 3 shows the trend for each monitor for the past ten years based on the 0.070 ppm 8-hour standard.

Figure 3
8-hour Ozone Exceedance Trends 2011 – 2020
 Exceedances Based on 0.070 ppm 8-hour Standard






Figure 3 cont.
8-hour Ozone Exceedance Trends 2011 – 2020
 Exceedances Based on 0.070 ppm 8-hour Standard



3.2 - Sulfur Dioxide (SO₂) Summary

2020 SO₂ Data Summary

A summary of the 2020 SO₂ data is shown below (in parts per billion). MassDEP operated six SO₂ monitors during 2020. All values are well below applicable NAAQS.

SO ₂ 2020			1 st	2 nd	99 th		1 st	2 nd
CITY	COUNTY	ADDRESS	MAX	MAX	PCTL	ARITH	MAX	MAX
			1-HR	1-HR	1-HR	MEAN	24-HR	24-HR
Boston	Suffolk	Kenmore Square	2.8	2.1	1.8	0.30	1.2	1.0
Boston	Suffolk	Harrison Avenue	2.5	2.1	2.0	0.48	1.1	0.9
Fall River	Bristol	Globe Street	6.0	4.5	3.3	0.27	1.3	1.3
Springfield	Hampden	Liberty Street	4.8	2.9	2.8	0.17	3.2	1.6
Ware	Hampshire	Skyline Drive	1.9	1.1	1.0	0.15	0.7	0.7
Worcester	Worcester	Summer Street	3.1	2.5	2.3	0.17	0.9	0.7

Standards: 1-hour = 75 ppb 3-hour = 0.5 ppm (500 ppb)


1st, 2nd MAX 1-HR = First and Second Highest 1-hour Value

99th PCTL 1-HR = 99th Percentile of the 1-hour Maximum Value

ARITH MEAN = Annual Arithmetic Mean

1st, 2nd MAX 24-HR = First and Second Highest 24-hour Value


Sulfur Dioxide Monitor Locations



SO₂ Trends

Figure 4 shows the trend of the 1-hour 99th percentile for each SO₂ monitor for the past 10 years relative to the 1-hour standard of 75 ppb.

Figure 4
Sulfur Dioxide Trends 2011 – 2020
1-hour 99th Percentile Annual Average



3.3 - Nitrogen Dioxide (NO₂) Summary

2020 NO₂ Data Summary

A summary of the 2020 NO₂ data is shown below (in parts per billion). MassDEP operated 10 NO₂ monitors during 2020. All values are well below applicable NAAQS.


NO ₂ 2020			1 st MAX 1-HR	2 nd MAX 1-HR	98 th PERCENTILE VALUE	ARITH MEAN
CITY	COUNTY	ADDRESS				
Boston	Suffolk	Kenmore Square	47	46	42	10.32
Boston	Suffolk	Harrison Avenue	54	48	42	9.34
Boston	Suffolk	Von Hillern Street	50	49	46	12.48
Chelmsford	Middlesex	Manning Road	46	43	39	10.99
Chicopee	Hampden	Anderson Road	43	37	33	4.30
Lynn	Essex	Parkland Avenue	41	41	34	5.05
Milton	Norfolk	Canton Avenue	38	32	25	3.46
Springfield	Hampden	Liberty Street	50	48	44	9.68
Ware	Hampshire	Skyline Drive	43	42	23	2.20
Worcester	Worcester	Summer Street	53	51	45	10.11

Standards: 1-hour = 100 ppb Annual Arithmetic Mean = 53 ppb

1st, 2nd MAX 1-HR = First and Second Highest 1-hour Value

ARITH MEAN = Annual Arithmetic Mean


Nitrogen Dioxide Monitor Locations



NO₂ Trends

Figure 5 shows the trend of the 1-hour 98th percentile annual average for each NO₂ monitor over the past 10 years relative to the 1-hour standard of 100 ppb.

Figure 5
Nitrogen Dioxide Trends 2011 – 2020
 1-hour 98th Percentile Annual Average



3.4 - Carbon Monoxide (CO) Summary


2020 CO Data Summary

A summary of the 2020 CO data is shown below (in parts per million). MassDEP operated three CO monitors during 2020. All values are well below applicable NAAQS.

CO 2020			1 st	2 nd	1 st	2 nd
			MAX	MAX	MAX	MAX
CITY	COUNTY	ADDRESS	1-HR	1-HR	8-HR	8-HR
Boston	Suffolk	Harrison Avenue	1.618	1.573	1.2	1.1
Boston	Suffolk	Von Hillern Street	1.048	1.042	0.7	0.6
Worcester	Worcester	Summer Street	2.281	2.119	1.2	1.1

Standards: 1-hour = 35 ppm 8-hour = 9 ppm
 1st, 2nd MAX 1-HR = First and Second Highest 1-hour Value
 1st, 2nd MAX 8-HR = First and Second Highest 8-hour Value


Carbon Monoxide Monitor Locations



CO Trends

Figure 6 shows the trend of the 2nd maximum 8-hour average for each CO monitor over the past 10 years relative to the 8-hour standard of 9 ppm.

Figure 6
Carbon Monoxide Trends 2011 – 2020
2nd Maximum 8-hour Values



3.5 - Particulate Matter 10 Microns (PM₁₀) Summary

2020 PM₁₀ Data Summary

A summary of the 2020 PM₁₀ data is shown below (in µg/m³). MassDEP operated four PM₁₀ monitors in 2020. All values are well below applicable NAAQS.

PM ₁₀ 2020			1 st	2 nd	3 rd	4 th	DAYS	
CITY	COUNTY	ADDRESS	MAX	MAX	MAX	MAX	MAX	ARITH
			24-HR	24-HR	24-HR	24-HR	>STD	MEAN
Boston	Suffolk	Harrison Avenue	26	25	23	23	0	12.5
Boston	Suffolk	Harrison Avenue*	27	23	22	21	0	12.2
Ware	Hampshire	Skyline Drive	20	16	16	14	0	8.1
Worcester	Worcester	Summer St	36	33	23	22	0	12.7


Standard: 24-hour = 150 µg/m³

* = Collocated monitors

1st, 2nd, 3rd, 4th 24-HR MAX = First, Second, Third and Fourth Highest 24-hour Values for the Year

ARITH MEAN = Annual Arithmetic Mean


PM₁₀ Monitor Locations



PM₁₀ Trends

Figure 7 shows the 2020 calendar year 24-hour maximum concentration for each PM₁₀ monitor relative to the 24-hour standard of 150 µg/m³.

Figure 7
PM₁₀ Trends 2011 – 2020
24-Hour Calendar Year Maximum



3.6 - Particulate Matter 2.5 Microns (PM_{2.5}) Summary

During 2020, MassDEP operated 11 Federal Reference Method (FRM) filter-based PM_{2.5} monitors and 17 Federal Equivalence Method (FEM) continuous PM_{2.5} monitors.

FRM monitors require the manual set-up and collection of filters that measure 24-hour samples every three or six days. The filters are weighed prior to placement in the field and then weighed again after the sample is collected to determine the amount of PM_{2.5} collected on the filter.

FEM monitors measure PM_{2.5} on an hourly basis. MassDEP operates FRM and FEM monitors side-by-side at some locations for comparison. In general, FEM monitors provide slightly higher values than FRM monitors, which may be due to measurement of fresh hourly PM_{2.5} samples with FEM versus slightly aged 24-hour samples with the FRM filter method.

2020 PM_{2.5} FRM Data Summary

A summary of the 2020 PM_{2.5} FRM data is shown below (in µg/m³).

PM _{2.5} 2020			1 st	2 nd	3 rd	4 th	98 th	
CITY	COUNTY	ADDRESS	MAX 24-HR	MAX 24-HR	MAX 24-HR	MAX 24-HR	PERCENTILE 24-HR	ARITH MEAN
Boston	Suffolk	Kenmore Square	17.0	14.5	12.9	10.9	12.9	6.28
Boston	Suffolk	Harrison Avenue	18.2	14.3	12.7	11.9	12.7	6.37
Boston	Suffolk	Von Hillern Street	14.9	12.2	11.9	10.9	12.2	7.38
Brockton	Plymouth	Clinton Street	13.8	11.5	11.3	10.8	11.5	6.59
Chicopee	Hampden	Anderson Road	31.4	15.9	14.0	13.6	15.9	6.85
Chicopee	Hampden	Anderson Road*	30.4	15.4	11.0	10.3	15.4	6.31
Greenfield	Franklin	Barr Avenue	33.3	22.1	18.4	16.0	18.4	7.72
Haverhill	Essex	Washington Street	16.8	13.7	12.4	12.3	13.7	7.21
Pittsfield	Berkshire	Silver Lake Blvd	32.4	30.6	22.7	16.4	22.7	6.70
Springfield	Hampden	Liberty Street	18.3	16.9	14.5	13.6	16.9	6.91
Worcester	Worcester	Summer Street	15.4	13.3	12.5	10.5	13.3	7.00

Standards: Annual Mean = 12.0 µg/m³ (primary) 24-hour (98th percentile) = 35 µg/m³


* = Collocated monitors

1st, 2nd, 3rd, 4th 24-HR MAX = First, Second, Third, and Fourth Highest 24-hour Values for the Year

98th PERCENTILE 24-HR = 98th Percentile Value for the Year

ARITH MEAN = Annual Arithmetic Mean



PM_{2.5} Monitor Locations



PM_{2.5} FRM Trends

Figure 8 shows trends of the annual arithmetic mean for each PM_{2.5} FRM monitor over the past 10 years relative to the standard of 12 µg/m³.

Figure 8
PM_{2.5} Trends 2011 – 2020
 FRM Annual Arithmetic Mean



2020 PM_{2.5} FEM Data Summary

A summary of the 2020 PM_{2.5} FEM data is shown below (in µg/m³).

FEM 2020			1 st	2 nd	3 rd	4 th	98 th	
CITY	COUNTY	ADDRESS	MAX 24-HR	MAX 24-HR	MAX 24-HR	MAX 24-HR	PERCENTILE 24-HR	ARITH MEAN
Boston	Suffolk	Kenmore Sq	25.0	19.7	18.3	18.1	18.1	6.92
Boston	Suffolk	Harrison Ave	25.6	21.7	21.6	20.2	14.3	5.75
Boston	Suffolk	Von Hillern St	26.0	20.7	20.2	19.6	16.2	8.27
Boston	Suffolk	Von Hillern St*	23.5	19.5	17.5	17.1	15.0	6.93
Brockton	Plymouth	Clinton Street	22.6	21.3	20.0	19.6	17.7	7.52
Chelmsford	Middlesex	Manning Road	24.0	22.9	21.0	20.6	13.9	6.70
Chicopee	Hampden	Anderson Road	26.4	25.0	23.7	20.3	18.1	6.58
Fall River	Bristol	Globe Street	21.4	20.5	19.5	17.4	16.8	6.22
Greenfield	Franklin	Barr Avenue	28.9	28.5	28.4	27.7	20.2	8.21
Haverhill	Essex	Washington St	22.4	20.0	19.9	14.5	13.5	5.23
Lynn	Essex	Parkland Ave	19.3	18.3	17.0	15.2	13.4	5.51
North Adams	Berkshire	Holden Street	20.0	18.4	16.1	15.8	14.5	5.39
Pittsfield	Berkshire	Silver Lake Blvd	27.7	26.0	21.8	21.5	17.3	7.58
Springfield	Hampden	Liberty Street	35.0	33.2	29.5	28.0	22.1	8.67
Ware	Hampshire	Skyline Drive	17.2	15.2	15.0	14.7	14.2	6.80
Weymouth	Norfolk	Bridge St	21.5	20.1	20.0	18.8	15.7	5.22
Worcester	Worcester	Summer Street	18.0	17.5	17.4	17.4	16.3	8.04

Standards: Annual Mean = 12.0 µg/m³ (primary) 24-hour (98th percentile) = 35 µg/m³

* = Collocated monitors

1st, 2nd, 3rd, 4th 24-HR MAX = First, Second, Third, and Fourth Highest 24-hour Values for the Year



98th PERCENTILE 24-HR = 98th Percentile Value for the Year

ARITH MEAN = Annual Arithmetic Mean

PM_{2.5} FEM Trends

Figure 9 shows trends of the annual arithmetic mean for each PM_{2.5} FEM monitor over the past 10 years relative to the standard of 12 µg/m³.

Figure 9
PM_{2.5} Trends 2011– 2020
FEM Annual Arithmetic Mean



3.7 - Speciation

MassDEP collects PM_{2.5} samples for speciation in Boston (Harrison Avenue) and Chicopee. Speciation involves analysis of particulate matter to determine its chemical composition and to identify air pollution sources that affect the area around the monitoring station. Pollutants analyzed include elements (e.g., metals), sulfates, nitrates, and carbon (total and organic).

3.8 - Interagency Monitoring of Protected Visual Environments (IMPROVE)

IMPROVE is a nationwide program designed to assess air quality at rural locations where air pollution may affect visibility over long distances (e.g., mountain ranges or scenic vistas). During 2020, the National Park Service operated an IMPROVE sampler at the Truro monitoring site, and the Wampanoag Tribe operated an IMPROVE sampler at its Martha's Vineyard monitoring site. These samplers acquire PM_{2.5} filter samples for speciation analysis to determine effects on visibility. Data can be viewed at the IMPROVE web site at <http://vista.cira.colostate.edu/improve/Data/data.htm>.

3.9 - Quality Assurance and Quality Control (QA/QC)

In order to ensure that all air quality data is of acceptable and consistent quality, MassDEP has developed standard operating procedures (SOPs) based on federal requirements that include quality control and quality assurance protocols that systematically assess the entire sample collection and data handling system on an ongoing basis. Ambient air monitoring quality assurance requirements are contained in the Federal Regulations at 40 CFR Part 58, Appendix A – E. Each year MassDEP certifies that it is in compliance with the federal requirements. A few of the considerations that affect sample collection data quality are:

- Site Placement
- Intake Probe Material
- Intake Probe Height
- Spacing from roadways and trees

For data processing there are quantitative statistics and qualitative descriptors used to interpret the degree of acceptability and utility of data. Examples of these data quality indicators are:

- Representativeness
- Precision
- Bias
- Detectability
- Completeness
- Comparability

MassDEP's Air Assessment Branch maintains a Quality Assurance Group that ensures samples are collected correctly and conducts performance audits throughout the air monitoring network to verify data validity. There also is a Quality Control Group that reviews daily monitored and historical data for validity, tracks precision results, finalizes monthly values, and submits air quality data to EPA's database in a timely manner. Computer software tools, report queries, and "eyes on" data reviews all are used to validate data before it is submitted to EPA. EPA also conducts its own performance audits on MassDEP samplers and every three years conducts a thorough Technical Systems Audit (TSA). The latest TSA was conducted in 2019.

Section 4 – PAMS/Air Toxics Monitoring

4.1 - Photochemical Assessment Monitoring Station (PAMS) Monitoring

Ground-level ozone is a secondary pollutant and is not emitted directly to the atmosphere from a stack or tailpipe, but forms in the atmosphere from the photochemical reactions of other pollutants such as VOCs and NO_x. The PAMS program was established to gather information on the ozone formation process to help agencies identify effective ozone pollution control strategies. In 2020, MassDEP operated one PAMS station in Lynn.

Instruments at the Lynn site measure pollutants and meteorological parameters that are specific to the photochemical processes by which ozone is created in the atmosphere at ground level. In addition to the relevant NAAQS pollutants (e.g., ozone, NO₂), the Lynn site monitors VOCs every six days throughout the year and on an hourly basis during the hottest part of the summer (June, July, and August). Meteorology is a critical component of ozone formation and the Lynn PAMS site has a full complement of meteorological sensors including wind speed, wind direction, temperature, relative humidity, barometric pressure, solar radiation, and precipitation.

4.2 - Air Toxics Monitoring

Toxic air pollutants are known or suspected to cause cancer or other serious health effects. Air toxics include certain VOCs and toxic metals (e.g., arsenic, cadmium). The Boston (Harrison Avenue) monitoring site is designated as a National Air Toxics Trends Station (NATTS) and monitors a number of toxic air pollutants, including VOCs, carbonyls, metals, black carbon and polycyclic aromatic hydrocarbons (PAHs). Data from this site is compared with data from a network of similar sites across the country to identify transport, trends, and site-specific characteristics of these pollutants. The table below summarizes the 2020 metals data.

2020 Metal	Boston (Harrison Ave)	
	Max Value µg/m ³	Mean µg/m ³
Antimony	0.00390	0.00143
Arsenic	0.00205	0.00045
Beryllium	0.00002	0.00000
Cadmium	0.00018	0.00006
Chromium	0.00743	0.00419
Cobalt	0.00029	0.00010
Lead	0.00720	0.00233
Manganese	0.01270	0.00496
Mercury	0.00002	0.00000
Nickel	0.00351	0.00094
Selenium	0.00061	0.00022

MassDEP monitors VOCs in Lynn as part of the PAMS program, some of which are classified as air toxics. In February 2020, MassDEP began monitoring VOCs in Weymouth. The table below summarizes 24-hour concentrations of target VOCs measured at the Boston (Harrison Avenue), Lynn, and Weymouth (Bridge Street) sites for 2020.

2020 Compounds	Boston (Harrison Ave)		Lynn		Weymouth (Bridge St)	
	Max Value ppb	Mean ppb	Max Value ppb	Mean ppb	Max Value ppb	Mean ppb
benzene	0.37	0.13	0.28	0.12	0.23	0.11
toluene	0.54	0.25	0.53	0.17	0.76	0.19
ethylbenzene	0.09	0.03	0.05	0.02	0.11	0.04
xylene	0.31	0.13	0.23	0.09	0.71	0.15
acrolein	0.40	0.09	0.17	0.09	0.26	0.10
acrylonitrile	<0.01	<0.01	0.01	<0.01	0.02	<0.01
1,3-butadiene	0.05	0.01	0.03	0.01	0.04	0.01
carbon tetrachloride	0.09	0.08	0.09	0.08	0.10	0.08
chloroform	0.03	0.02	0.04	0.02	0.03	0.02
1,2-dichloroethane	0.02	0.02	0.02	0.02	0.02	0.02
ethylene oxide	0.15	0.06	0.10	0.06	0.20	0.07
1,1,1-trichloroethane	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
tetrachloroethylene	0.10	0.01	0.05	0.00	0.02	0.01
trichloroethylene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
vinyl chloride	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
acetaldehyde	1.15	0.60	1.60	0.86	1.19	0.66
formaldehyde	4.50	1.79	5.90	2.85	5.18	1.67
naphthalene	0.02	0.01	0.03	0.01	0.09	0.01

Notes:

1. Weymouth (Bridge St) was a temporary site that did not meet EPA siting criteria. MassDEP established a permanent site in April 2021 on Monatiquot Street in Weymouth that meets EPA siting criteria.
2. Table does not include values affected by sampling canister contamination.

Appendix A 2020 Monitoring Stations

City/Town	Address	Parameters Monitored
Aquinnah*	Herring Creek Road	O ₃ , PM _{2.5} speciated (IMPROVE)
Boston	Kenmore Square	NO ₂ , NO, NO _x , SO ₂ , PM _{2.5}
Boston	1159 Harrison Ave	O ₃ , NO ₂ , NO, NO _x , NO _y , SO ₂ , CO, PM _{2.5} , PM ₁₀ , PM _{Coarse} , PM _{2.5} speciated, Black Carbon, VOCs, Carbonyls, metals, PAHs, WS/WD, WSv/WDv, TEMP, SUN, REL, BP
Boston	19 Von Hillern Street	NO ₂ , NO, NO _x , CO, PM _{2.5} , Black Carbon, WS/WD, TEMP, SUN, REL, BP
Brockton	170 Clinton Street	O ₃ , PM _{2.5}
Chelmsford	11 Technology Drive	O ₃
Chelmsford	5 Manning Road	O ₃ , NO ₂ , NO, NO _x , PM _{2.5} , Black Carbon
Chicopee	Anderson Road	O ₃ , NO ₂ , NO, NO _x , PM _{2.5} , PM _{2.5} speciated, WS/WD, TEMP, SUN, REL, BP
Fairhaven	30 School Street	O ₃ , WS/WD, TEMP, SUN, REL, BP
Fall River	659 Globe Street	O ₃ , SO ₂ , PM _{2.5}
Greenfield	16 Barr Avenue	O ₃ , PM _{2.5} , Black Carbon, WS/WD, TEMP, SUN, REL, BP
Haverhill	685 Washington Street	O ₃ , PM _{2.5} , WS/WD, TEMP, SUN, REL, BP
Lynn	390 Parkland Avenue	O ₃ , NO ₂ , NO _y , PM _{2.5} , VOCs, Carbonyls, WS/WD, TEMP, SUN, REL, BP, PRECIP, TOTAL UV
Milton	1904 Canton Avenue	O ₃ , NO ₂ , NO, NO _x , WS/WD, TEMP, SUN, REL, BP
North Adams	86 Holden Street	PM _{2.5} , Black Carbon
Pittsfield	25 Silver Lake Drive	O ₃ , PM _{2.5} , Black Carbon, WS/WD, TEMP, SUN, REL, BP
Springfield	600 Liberty Street	NO ₂ , NO, NO _x , SO ₂ , PM _{2.5} , Black Carbon
Truro	6 Collins Road	O ₃ , WS/WD, TEMP, SUN, REL, BP, PM _{2.5} speciated (IMPROVE)
Uxbridge	366 E. Hartford Ave	O ₃ , WS/WD, TEMP, SUN, REL, BP
Ware	36 Skyline Drive	O ₃ , NO ₂ , NO, NO _x , NO _y , SO ₂ , PM _{2.5} , PM ₁₀ , WS/WD, TEMP, SUN, REL, BP, PRECIP
Weymouth	6 Bridge Street	PM _{2.5} , VOCs, Carbonyls
Worcester	375 Airport Drive	O ₃ , WS/WD, TEMP, SUN, REL, BP
Worcester	Summer Street	NO ₂ , NO, NO _x , SO ₂ , CO, PM _{2.5} , PM ₁₀

*Tribal Site