



\*Ceftriaxone 500mg IM x1 dose  
 Doxycycline 100mg orally twice a day for 7 days  
 Metronidazole 500mg orally twice a day for 7 days<sup>1</sup>  
 Recommend: Hepatitis B and HPV vaccines, if not previously vaccinated.  
 \*See CDC guidelines for special populations and for patients weighing ≥150 kg

**STI testing**  
 Urine NAAT: GC/Ch/Trich  
 Blood: HIV<sup>2</sup>, Syphilis serology, Hep C Ab, Hep B sAg, sAb, cAb

Additional baseline labs for HIV nPEP<sup>3</sup>: Serum creatinine, ALT/AST  
 If STI testing was declined, readdress HIV, Hep B sAg, sAb, cAb testing (see note below.)<sup>3</sup>  
 For medications, see CDC nPEP guidelines or consult ID

Site specific testing (oral, anal, vaginal): generally **NOT** recommended during acute timeframe.  
 Patients with genital s/s of infection: management per ED provider

1. Prophylaxis and testing for trichomoniasis is only recommended for patients with a vagina.
2. HIV – ½ antigen/antibody combo immunoassay (4th generation) testing preferred. Ab= antibody. sAg= Surface Antigen, sAb= Surface Antibody, cAb= Core Antibody
3. If a patient declines serology, decisions regarding nPEP administration should be made by the medical provider, in consultation with the patient, based on the patient’s medical history and assault-related risk factors.