Massachusetts Department of Conservation and Recreation Division of Water Supply Protection, Office of Watershed Management Forest Management Project Proposal Summary for Public Comment

Location, goals, and summary of proposed forest management.

Item Information/Description
NS-25-02
2025
Quabbin
Shutesbury
Richard G MacLean
77
New Salem
02
Located Between Camel Brook and Cooleyville Rd in Shutesbury
NS-16-02, PE-19-15-01
This proposal seeks to advance overall watershed goals of increasing age and species
diversity through harvest. Species diversity will be improved by harvesting areas with
low species diversity while retaining underrepresented species. Age diversity goals will
be pursued by opening the forest canopy over existing regeneration and providing open patches with high sunlight to create opportunities for new regeneration to
establish. This proposal contains a prominent hill, vulnerable to high wind disturbance,
that has a stand of tall even aged white pine on it. Introducing openings at the top of
the hill, with thinning in between openings, will provide an opportunity for new
regeneration to grow and increase the resiliency of that area to large scale wind
disturbance. In the pine hemlock stand, small openings cooccurring with hemlock
regeneration will help encourage hemlock regeneration and provide a refugia for
hemlock which is currently struggling with multiple insect and climate stressors.

Forest Cover Types and Acreages

Overstory Forest Types	Acres
White pine - hemlock	51
White pine - oak	14
Mixed oak	8
White pine	4

Understory Cover Types and Relative Importance

Understory Cover Type	Relative area covered (Dominant, Secondary, Minor, None)
Tree seedlings and saplings	Primary
Mountain laurel	Secondary
Mesic site - witch hazel, highbush	Minor
blueberry	
Dry site -Huckleberry, blueberry	Minor

Understory Cover Type	Relative area covered (Dominant, Secondary, Minor, None)
Mesic site - cinnamon fern, mixed	None
hardwood	
Hayscented fern	None
Invasive shrubs/vines	Minor
Other	

Vegetation Topic	Description
General Description, Forest Composition, Stand History, and Harvest History	The proposed harvest area covers a hilltop and the surrounding slopes between the West Branch of the Swift and streams which feed into Camel Brook. A skid road/old cart road wraps the base of the hill leading from Cornwell Rd east below the peak of the hill and the heading north towards the DWSP boundary with private abutters. The area can be divided into four general stands.
	A majority of the site is a white pine – hemlock stand wrapping around the hill from the west south around to the east. It varies in relative proportion of the two dominant species from heavily pine in the west to more heavily hemlock on the south and eastern sides. There is a secondary component of northern red oak scattered through the southern and eastern portions of the stand. Canopy dominant hemlock mortality from hemlock wooly adelgid is present but is scattered to individual snags. A few areas with a group of canopy hemlock mortality near camel brook, west of Cornwell Road have diverse regeneration now small pole sized. Just north of the intersection of the old skid trail and Cornwell Rd is a 1.4 acre patch opened up in a 1987 has regenerated to pure white pine.
	The northern slope of the hill leading up to the top is a fourteen acre white pine - oak stand. This stand contains over 100 ' tall emergent white pine on a southwest to northwest aspect with some prominence and exposure to open private land to the north west. This position increases vulnerability to large scale hurricane winds or sever winter prevailing winds.
	The top of the eastern and south facing slopes is dominated by a mixed oak cover with a minor white pine component. There are a few acres on the southwest corner that are unmanageable and a few more on the eastern face that would only be manageable with a cable skidder.
	Finally, there is a four acre stand of overstory white pine and remnant dying red pine. Under this canopy is a midstory of pole sized northern red oak, hemlock, white pine, black birch, and red maple. This stand is located at the corner of Cornwell Rd and Cooleyville Rd and the tall white pine may represent a hazard to those roads in the event of a large wind disturbance.
	Past management has occurred throughout the proposed area in multiple entries. Nearly all of the area was harvested with a shelterwood prep cut in 1963 in the seventh commercial harvest at Quabbin. The area west of Cornwell Rd and the top and east/northeastern slope of the hill were harvested in 1984 and 1985 likely related to the record spongy moth outbreak in 1981. Finally, the eastern lowest slopes were treated in 1987 by the Quabbin internal crew.

Vegetation Topic	Description
Advance Regeneration description	The proposed area was sampled at roughly 1 plot per acre. Approximately one third of the proposed area is in monoculture regeneration, almost entirely hemlock, with a small percentage in black birch and white pine. This monoculture hemlock regeneration is distributed across the white pine – hemlock stand. A fifth of the area has little to no regeneration with sparse hemlock, scattered mountain laurel or witch hazel, or no woody stems at all. In the next fifth, regeneration is prevented by interfering dense mountain laurel, and to a smaller degree tall dense witch hazel. All the sampled plots categorized as interfered are in the white pine – oak and mixed oak stands at the top and northern slope of hill. Another tenth was classified as marginally regenerated, but even in these cases where there was some marginal diversity regeneration was still dominated by hemlock. Only one twentieth of the area was found to have diverse regeneration, and again, in these plots hemlock was the dominant species.
Terrestrial Invasive Plants description	Terrestrial invasive species were only observed in small populations of barberry on Cornwell Rd, as well as adjacent to the landing south of the Camel Brook bridge in the former gravel pit.

Description of Wetland Resources Present

rescription of weetiana resources i resent		
Resource Type	Description of resources present	
Wetlands	There are small seasonal wetlands associated with the intermittent stream west of	
	Cornwell Rd. These will not be harvested but may need to be crossed to access areas	
	of higher ground. All attempts will be made to avoid crossings. There is a small	
	amount of wetland immediately adjacent to the West Branch of the Swift River that	
	will also be avoided.	
Streams	The proposed harvest area is bounded by the West Branch of the Swift River to the	
	east, intermittent streams feeding Camel Brook to the west and Camel Brook to the	
	southwest. No streams crossings will be needed.	
Vernal pools	None known.	
Seeps	None known.	

Description of Soils by Hydric Class

Soil Hydric Classes	% of area	Soil series and any further comments
Excessively Drained	39	Hinckley sandy loam
Well-drained Thin	11	Chatfield-Hollis complex
Well-drained thick	44	Two thirds Montauk fine sandy loam, one third Canton fine sandy loam
		and possibly a small amount of Newfields fine sandy loam.
Moderately well-drained	5	Sudbury sandy loam with a small amount of Walpole fine sandy loam.
Poorly to very poorly drained	1	Walpole fine sandy loam

Proposed Silvicultural Activities

Topic	Description
Site Selection and	The portion of the proposed area made prominent above the surrounding area by the small
Silvicultural	hill leads to vulnerability of large scale wind disturbance for the even aged stand. By
Objectives	harvesting here and increasing age diversity through regeneration of a new age class, overall
	forest resiliency can be increased. In the surrounding lower areas currently dominated by
	scattered hemlock regeneration increasing light to the understory will encourage a greater
	diversity of species recruitment and increase the species diversity of the future forest.
Silviculture	With a focus on this site of recruiting new age classes, silviculture will primarily rely on group
Prescription	selection openings with species diverse edge retention trees. Size will vary by cover, and
	prominence. In the lowest areas of the white pine – hemlock stand opening sizes may be as
	small as half an acre and focus on release of existing hemlock regeneration in order to treat
	those low-lying areas as somewhat of a refuge for a hemlock dominant cover type. Steeper
	slopes will similarly be treated with intermediate thinning to small openings to reduce soil
	erosion hazards.
	In areas with more exposed prominence, larger openings will target the tallest most
	vulnerable white pine, as well as targeting the areas of dense mountain laurel. This will
	provide a large amount of light as well as high mechanical disturbance of the mountain laurel
	in order to provide an opportunity for fast growing shade intolerant tree species to establish
	before the mountain laurel can recover and impede regeneration. Larger openings on
	prominent hilltops have also been shown in research to have a higher benefit to migrating
	bird and insect populations than a similar sized opening on flat or low-lying ground.
	Across Cornwell Rd the current cover type is similar but contains evidence that it was in red
	and white pine plantation after the original taking. Remnant red pine will be removed to
	reduce hazardous conditions created by red pine scale related mortality. Larger openings will
	be implemented to release the few areas of diverse hardwood regeneration that are already
	present there.
	present mere.

Climate Change Considerations: DWSP has determined that the decision to implement this project is consistent with EEA climate goals and guidelines and agency land management objectives. Carbon and climate change considerations specific to the activities proposed for this project are discussed below.

Proposed Activity	Alignment of Activity with Climate Oriented Strategies and Recommendations	
Full overstory removal, partial stand, patch regeneration cut. (see page 4, Silvicultural Prescription, group selection openings)	Patch cutting is a regeneration technique that straddles the boundary between classic even-aged and uneven-aged forest management systems. Foresters select appropriate areas ('patches' or 'groups') covering a portion of the stand to harvest rather than removing the entire stand and then return periodically to repeat the process in other portions of the stand. In using patch cutting there is no final regeneration cut. Patch size and shape are determined by many different factors including overstory condition, desired species composition in the regeneration layer, other desirable herbaceous and woody vegetation, location, stand re-entry period, etc. Harvesting in patches aligns with many climate-smart forestry practices:	
	 Increasing structural diversity improves resiliency by reducing the impact of age/size related disturbances. Extending regeneration periods minimizes short term impacts to groundwater and nutrient cycling. Partial stand overstory removals more closely align with natural disturbance patterns. More carbon is left on the landscape for longer periods, and withinpatch live tree, snag, and coarse debris retention allow for development of old forest characteristics. Can also be used as opportunities to increase the stocking of future climate adapted species, current climate imperiled species, or other types of desirable vegetation. 	
Diffuse overstory removal, partial	Partial cutting via single trees or small groups in a mature stand can advance a	
cut, late rotation regeneration	variety of management objectives as well as climate-smart practices . Single	
related.	tree or very small group removals, if used exclusively and repeatedly, will	
(see page 4, Silvicultural Prescription, intermediate thinning on steeper slopes)	perpetuate an uneven-aged stand condition with a species mix shifted towards higher shade tolerance. However, this type of harvest can also serve within an even-aged system to establish regeneration of species of lower shade tolerance under a partial canopy for subsequent release using larger group or patch cuts (irregular shelterwood) or complete-stand overstory	
	removals. Advantages of partial overstory removals include, but not limited to:	
	 Partial cutting retains carbon on the landscape for extended periods while regeneration develops. Reducing competition for resources improves growth and carbon sequestration rates on residual trees. Promotion of a diversity of age classes enhances overall forest resiliency. Maintenance of continuous forest corridors provides for wildlife habitat. As part of a regeneration system this method can be used to help guide species diversity towards more future-adapted mixes. 	

Proposed Activity	Alignment of Activity with Climate Oriented Strategies and Recommendations
General/other Climate Change	With higher severity storms a concern with changing climate emergent white
Considerations	pine will be a focus of this harvest. Similarly, when retained, care will be taken to retain tall white pine in clusters, and away from areas of public use to reduce hazards.

Equipment and Access Constraints and Considerations

4-1	onstraints and considerations
Constraint Topic	Description and Considerations
Proposed Equipment	Forwarder will be required to transport wood to landings across DCR internal roads.
requirements	
Proposed wetland or	This lot will take advantage of the new bridge across Camel Brook when completed. No
stream crossings	temporary stream crossings will be needed, a temporary wetland crossing may be necessary to
	access areas on the west side of Cornwell Rd.
Further wetland	
comments	
Vernal Pools	None known.
Access improvements	Completion of the Camel Brook bridge is needed. Improvement of Cornwell Rd to allow trucks
needed	to access from 202 and reach the intersection of Cornwell Rd and Cooleyville Rd would be very
	helpful.
Other EQ issues	None known.
In-kind Services	None planned.
Other Access	An old skid/cart road is used as an unauthorized trail by abutters and will be protected during
Concerns (parking,	operations/required to be restored upon completion.
trails, etc.)	

Subwatershed Analysis

Sub-Watershed number/name	Total DCR- owned acres in this sub- watershed	Acres regenerated on DCR land in the last 10 years in this sub- watershed	Total DCR-owned acres remaining for regenerating up to the 25% per 10 year limit for this subwatershed	Acres in this sub-watershed that are part of this proposed lot
59: Camel Brook	351	11.6	76.2	37
90: West Branch of Swift	620	5	150	29
Mainstem				

Additional comments on Subwatershed analysis: No comments.

Wildlife and Habitat Observations and Considerations

Wildlife/Habitat	Observations and Considerations
Natural Heritage Priority Habitats?	Yes. There are NHESP priority polygons along the northern section and along the eastern side of this lot.
State Listed species present:	NHESP has determined that certain state-listed sensitive species or habitats may exist within the lot proposal area. To protect them from unnecessary disturbance, detailed information regarding affected species and their locations is not included in this report. DWSP will coordinate with NHESP and follow recommendations to protect these species during the proposed activity.
Rare Natural Communities:	None known.
General Wildlife Comments	Current browse evidence is well distributed across the proposed area with browsed down hemlock, missing browse preferred regeneration species, and even non preferred species browse present throughout. The areas of extensive mountain laurel may be the result of past higher browse levels which prevented any tree regeneration from establishing and selected for the unpalatable mountain laurel monocultures present today.


Cultural Resources Description and proposed protection measures

Cultural Resource	Description and proposed protection measures
Historical features present; comments regarding protection	
Description of site characteristics in relation to Ancient sites modeling or other verified evidence	Surface stone is a relatively minor component outside of the steeper slopes and ledgey areas. Microtopography is somewhat uneven west of Cornwell road and the southern slopes leading down towards Camel Brook. The low lying areas adjacent to the West Branch of the Swift River have the least microtopography.
	Slope is variable but the areas of steepest slopes are the ledge areas on the southwest side of the hilltop, and the southeast to east face. If applicable, DWSP will follow the recommendations of DCR's Archeologist regarding protection of sensitive sites.

Division of Water Supply Protection Office of Watershed Management



NS-25-02 -- Locus Map






Division of Water Supply Protection Office of Watershed Management



NS-25-02 -- Stand Map






Division of Water Supply Protection Office of Watershed Management



NS-25-02 -- Soil Drainage Class






Division of Water Supply Protection Office of Watershed Management



NS-25-02 -- Wetlands and Wildlife Resources





Division of Water Supply Protection Office of Watershed Management





NS-25-02 -- Cultural Resources and Landscape Characteristics

