# Massachusetts Department of Conservation and Recreation Division of Water Supply Protection, Office of Watershed Management Forest Management Project Proposal Summary for Public Comment

Location, goals, and summary of proposed forest management.

	of proposed forest management.	
Proposal Summary Item	Item Information/Description	
Lot Proposal ID	PT-26-09	
Fiscal Year	2026	
Watershed	Quabbin	
Town(s)	Petersham	
Forester	Helen Johnson	
Estimated Acres by	21 to 29 total acres in regeneration openings and 50 to 64 acres of thinning.	
Treatment Type	Regeneration openings will range in size from 1/5 to 2 acres each, with an average size	
	of around 1.25 acres.	
Total Proposal Acres	88.5	
Block	Petersham	
Compartment and/or	9	
Working Unit		
Location and Boundary	This proposal is east of the <b>East Branch of Fever Brook</b> in Petersham, MA. It's bordered	
Description	to the south by a DCR access road (Dugway Road), to the west and north by Fever	
	Brook, and to the east by the height of land.	
Previous Proposal?	No	
Project Goals and	This site was selected due to its severely inadequate regeneration and lack of vertical	
Summary Description	structure, particularly near East Fever Brook where white pine and hemlock tower over	
	a largely empty understory.	
	The overstory has been weakened by forest health concerns, and white pine is highly vulnerable to breakage in high winds even when healthy. If a severe windstorm were to cause rapid loss of those tall trees, the watershed forest filter would be severely degraded, leaving the Brook and nearby Quabbin Reservoir vulnerable.	
	The primary goal of this proposal is to strengthen the watershed forest filter and increase its resilience by initiating a new cohort of young trees, which will be too small to be blown over by wind. This will be accomplished by creating openings of up to 2 acres where there are clusters of trees that are diseased, declining, or have poor stem structure, thereby increasing age class diversity while improving overall forest health and vigor.	

#### **Forest Cover Types and Acreages**

-	,,	
	Overstory Forest Types	Acres
	White Pine-Hemlock	40.86
	Hemlock-Hardwoods	24.25
	White Pine-Oak	11.49
ĺ	White Pine-Hardwoods	11.91

## **Understory Cover Types and Relative Importance**

Understory Cover Type	Relative area covered (Dominant, Secondary, Minor, None)
Tree seedlings and saplings	Secondary
Mountain laurel	Minor
Mesic site - witch hazel, highbush	Minor
blueberry	
Dry site -Huckleberry, blueberry	Minor
Mesic site - cinnamon fern, mixed	None
hardwood	
Hayscented fern	Minor
Invasive shrubs/vines	Minor
Other	Open understory: Dominant

#### **Forest Vegetation Description**

orest Vegetation Description		
Vegetation Topic	Description	
General Description, Forest Composition, Stand History, and Harvest History	White pine and hemlock are the dominant overstory species across the entire proposal, either as a single dominant canopy, or with a lower canopy of hardwoods, including red, white and black oak, red maple and black birch sawlogs. Pole-sized trees of the same species are present but not common.	
	Basal area ranges from 150 to 290 ft²/ac in the White Pine-Hemlock type, to 110 to 230 in Hemlock-Hardwoods and White Pine-Hardwoods, and 90 to 240 in White Pine-Oak. Standing dead trees are common across all stand types, averaging 33 ft²/ac, with a range of 0-90 ft²/ac.	
	The most pervasive forest health concern at this site is hemlock wooly adelgid, which has led to thinning crowns and extensive mortality. Some of the white pines also have thinning crowns, in some cases despite having plenty of growing space, raising the possibility of white pine needle drop fungi. Spongy moth mortality is present but not extensive, and nectria canker was observed on some birches.	
	Past harvests include 53.0 acres of shelterwood prep completed in 1963 (Lot 0005), 7.6 acres of shelterwood prep completed in 1965 (Lot 0019), and 19.8 acres of shelterwood prep completed in 1992 (Lot 0608).	
Advance Regeneration description	The vast majority of the proposal area has inadequate regeneration, ranging from no regeneration and virtually nothing in the understory (51% of the area) to sparse or marginal regeneration (46% of area). Only 2% of the area has adequate regeneration density, but with species limited to black birch, white pine and hemlock. Oak seedlings and saplings are present at extremely low density on only 5% of the proposal area, near Dugway Road. Interfering agents include witch hazel on about 10% of the area, most of it near Dugway Road, and hay-scented fern on 2% of the area.	
Terrestrial Invasive Plants description	A small (~0.05 acre) patch of Japanese knotweed is located in the pullout to the DCR access road just outside the southwest corner of the proposal, and will be avoided in order to prevent its spread. If possible, treatment prior to harvest is recommended.	

#### **Description of Wetland Resources Present**

Resource Type	Description of resources present	
Wetlands	Vegetated wetlands along the East Branch of Fever Brook border the proposal to the	
	north and west, and a small wetland (Verified Vernal Pool #13) borders the proposal	
	to the southeast.	
Streams	A perennial stream bisects the proposal from east to west, becoming braided as it	
	approaches Fever Brook. Outside the proposal are the East Branch of Fever Brook to	
	the west and north, intermittent streams to the northeast and west that flow into	
	Fever Brook, and an intermittent stream to the southeast that flows into Verified	
	Vernal Pool #13.	
Vernal pools	Verified Vernal Pool #13 is 100 feet outside the proposal to the southeast, and will be	
	protected according to the requirements in the 2017 DWSP Land Management Plan.	
Seeps	None known.	

#### **Description of Soils by Hydric Class**

Soil Hydric Classes	% of area	Soil series and any further comments
Excessively Drained	0	
Well-drained Thin	13	Charlton-Chatfield-Hollis association, 15 to 45 percent slopes, very
		rocky
Well-drained thick	87	Montauk-Scituate-Canton association, 3 to 15 percent slopes, extremely stony, Charlton-Chatfield association, 3 to 15 percent slopes, extremely stony, and Montauk-Canton association, 15 to 45 percent slopes, extremely stony
Moderately well-drained	0	
Poorly to very poorly drained	0	

#### **Proposed Silvicultural Activities**

Toposea Silvicultural Activities		
Topic	Description	
Site Selection and	This site was selected due to its severely inadequate regeneration and lack of vertical	
Silvicultural	structure, particularly near East Fever Brook where white pine and hemlock tower over a	
Objectives	largely empty understory.	
	The overstory has been weakened by forest health concerns including hemlock wooly adelgid, white pine needle drop, and spongy moth. Even when healthy, white pine is highly vulnerable to breakage in high winds (see <a href="Hurricane Damage and Forests">Hurricane Of 1938 Battered New England's Woods 75 Years Ago</a> ). If a severe windstorm were to cause rapid loss of those tall trees, the watershed forest filter would be severely degraded, leaving the Brook and nearby Quabbin Reservoir vulnerable.	
	The primary silvicultural objective of this proposal is to strengthen the watershed forest filter	
	and increase its resilience by initiating a new cohort of young trees, which will be too small to be blown over by wind. This will be accomplished by removing clusters of declining trees,	
	thereby increasing age class diversity while improving overall forest health and vigor. Species	
	diversity will be maintained in the overstory, and promoted in the understory by creating	
	, , , , , , , , , , , , , , , , , , , ,	
	openings large enough to regenerate to shade intolerants.	

Topic	Description
Silviculture	Regeneration openings 1/5 to 2 acres in size, totaling 21 to 29 acres and averaging around
Prescription	1.25 acres, will be located where there are clusters of trees that are diseased, declining, or have poor stem structure. Trees on the perimeter of openings will be healthy and vigorous with stable stem structure, and will be either vertical or leaning away from the adjacent opening in order to minimize damage to regeneration if they fall or are cut in the future. Witch hazel and trees ≥5 feet tall within openings will be cut, except for the rare healthy oak saplings, which will be flagged for retention. In 90% of openings over 1/2 acre, 5-10 ft²/acre of basal area will be retained, with healthy oaks and regionally uncommon native species being favored for retention.
	Areas between and around openings will be thinned, with particular attention to improving the vigor and seed bearing potential of oaks and making perimeter trees around openings more windfirm. Declining hemlock and white pine will be targeted for removal to improve forest health and increase biodiversity.

#### **General Climate Change Considerations:**

This silvicultural approach is designed to increase long-term forest and watershed resilience by addressing severe overstory decline and a significant lack of regeneration. Introducing a new age class helps buffer against climate-induced stressors by increasing stand heterogeneity and functional redundancy. These gaps are designed to promote light-demanding, climate-adapted hardwoods—particularly oaks—while retaining some healthy trees for structure, diversity, and future seed input. Removing high-risk stands containing declining hemlock and structurally compromised white pine reduces the potential for catastrophic blowdown and limits future losses from pests and pathogens. As part of this management, I recommend mentioning that select dead trees (snags) will be retained to support wildlife habitat, promote structural diversity, and contribute to long-term carbon storage. As they fall, they add ecological features just as coarse woody debris to help promote old growth characteristics.

DWSP has determined that the decision to implement this project is consistent with EEA climate goals and guidelines and agency land management objectives. Climate change considerations specific to the activities proposed for this project are discussed below.

proposed for this project are discussed below.		
Proposed Activity	Alignment of Activity with Climate Oriented Strategies and Recommendations	
Regeneration patch cuts (full	Patch cutting is a regeneration technique that straddles the boundary between	
overstory removal, partial stand)	classic even-aged and uneven-aged forest management systems. Foresters	
	select appropriate areas ('patches' or 'groups') covering a portion of the stand	
	to harvest rather than removing the entire stand and then return periodically to	
	repeat the process in other portions of the stand. In using patch cutting there is	
	no final regeneration cut. Patch size and shape are determined by many	
	different factors including overstory condition, desired species composition in	
	the regeneration layer, other desirable herbaceous and woody vegetation,	
	location, stand re-entry period, etc. Harvesting in patches aligns with many	
	climate-smart forestry practices:	
	<ul> <li>Increasing structural diversity improves resiliency by reducing the impact of age/size related disturbances.</li> </ul>	
	<ul> <li>Extending regeneration periods minimizes short term impacts to groundwater and nutrient cycling.</li> </ul>	
	Partial stand overstory removals more closely align with natural	
	disturbance patterns.	
	<ul> <li>More carbon is left on the landscape for longer periods, and within-</li> </ul>	
	patch live tree, snag, and coarse debris retention allow for development of old forest characteristics.	
	Can also be used as opportunities to increase the stocking of <b>future climate</b>	
	<b>adapted species, current climate imperiled species</b> , or other types of desirable vegetation.	

#### **Proposed Activity** Alignment of Activity with Climate Oriented Strategies and Recommendations Thinning (diffuse overstory Partial cutting via single trees or small groups in a mature stand can advance a removal, partial cut, regeneration variety of management objectives as well as climate-smart practices. Single tree or very small group removals, if used exclusively and repeatedly, will related) perpetuate an uneven-aged stand condition with a species mix shifted towards higher shade tolerance. However, this type of harvest can also serve within an even-aged system to establish regeneration of species of lower shade tolerance under a partial canopy for subsequent release using larger group or patch cuts (irregular shelterwood) or complete-stand overstory removals. Advantages of partial overstory removals include, but not limited to: Partial cutting **retains carbon on the landscape** for extended periods while regeneration develops. Reducing competition for resources improves growth and carbon **sequestration** rates on residual trees. Promotion of a diversity of age classes enhances overall forest resiliency. Maintenance of continuous forest corridors provides for wildlife habitat. As part of a regeneration system this method can be used to help guide species diversity towards more future-adapted mixes. Protection of soil carbon Routing skid trails to avoid steep slopes and sensitive soils and reinforcement of soft ground with slash (tops and branches from cut trees) prevent soil erosion a. Careful routing of skid trails to avoid steep slopes and sensitive and compaction. soils, and reinforcement of soft Water bars help stabilize skid trails and ensure that excessive erosion is ground with slash. avoided while maintaining the site for future forestry operations. Properly stabilized skid trails will revegetate naturally while being discernable enough to b. Installation of water bars in use in future operations. Beyond compliance with the BMP manual standards, accordance with Massachusetts the size and frequency of water bar installation, and degree of stabilization, Best Management Practices. should be determined by: c. Stabilize the soil at the landing Other uses that may occur between operations, e.g. hiking trails, with conservation mix or snowmobiles trails, use as firebreaks, or unauthorized uses (OHV/ATV) equivalent seed source. The **impacts of future climate conditions**, especially more frequent storms. If the area is already known to be wet, and in the future more frequent storms are expected, more water bars than what may be normally installed are encouraged. **Soil type**. Land managers may consider seeding and mulching water bars on highly erodible soils, steep slopes, or excessively wet areas to ensure longevity and prevent water bar degradation.

Proposed Activity	Alignment of Activity with Climate Oriented Strategies and Recommendations
Temporary stream crossing One stream crossing at a right angle to the stream, located above the braided portion of the stream to avoid the need for additional crossings. A portable bridge will be required if the stream is flowing at the time of the harvest, with the option to use a poled ford instead if the stream is not flowing.	<ul> <li>Temporary stream crossings are occasionally necessary to facilitate forest management activities, though careful project layout can help minimize the number of stream crossings required.</li> <li>Crossing design, installation, maintenance, and removal should be done in accordance with the highest standard BMPs to minimize impacts to sensitive stream and wetland resources.</li> <li>Plan for worst case climate scenarios that generally indicate a greater frequency of higher intensity precipitation events.</li> <li>Changing climatic conditions change will require the continuous evaluation.</li> </ul>
Invasive plant control, including pre- and/or post-harvest and follow up treatments.  Treatment/removal of non-native invasive plants prior to or following a forestry operation in accordance with an approved DWSP Invasive Terrestrial Plant Management Plan.	Strong consensus exists among land managers and climate science experts regarding the threat to future forest health posed by the introduction and spread of invasive plants. Invasive plants can:  • aggressively outcompete native plant species, • dominate understory communities, and even climb, kill, and topple mature trees, • threaten overall biodiversity. • threaten soil health and long-term carbon storage.
	<b>Monitoring and controlling</b> invasive and interfering plant populations prior to and following forestry operations is a critical practice for <b>minimizing the risk of further impacts</b> inadvertently (though not unexpectedly) spread by harvesting-related activities.
Additional Comments	

**Equipment and Access Constraints and Considerations** 

quipment and Access constraints and considerations		
Constraint Topic	Description and Considerations	
Proposed Equipment	Forwarder required to protect the DCR access road if the landing is located on the other side	
requirements	(see Other Access Concerns for more information).	
10401110110	(cos canos recosa concerno con mercano).	
Proposed wetland or	The stream crossing on the perennial stream that bisects the proposal will be located uphill	
stream crossings	(east) of the portion of the stream that's braided. Bridge mats will be required.	
Further wetland	none	
comments		
Vernal Pools  Verified Vernal Pool #13 will be protected by a 200 foot low ground disturbance		
	required in the 2017 DWSP Land Management Plan.	
Access improvements	none	
needed		
Other EQ issues	none	
In-kind Services	Treatment of Japanese knotweed in the road pullout SW of the proposal may be desirable, but	
	would be better started a few years before the harvest.	

Constraint Topic	Description and Considerations
Other Access Concerns (parking, trails, etc.)	The best place for a landing is in the road pullout to the southwest, but that location will not be used in order to avoid transporting the Japanese knotweed that's present at that location. In addition, contractors should take care not to move Japanese knotweed fragments of any size if they park in that area.
	Other possible landing sites include inside the roadside stone wall to the east of the road pullout (portions of the road without a stone wall are too steep), or across the DCR access road at the landing for PT-26-08, which could be operated in conjunction with this proposal.

#### **Subwatershed Analysis**

Sub-Watershed number/name	Total DCR- owned acres in this sub- watershed	Acres regenerated on DCR land in the last 10 years in this subwatershed	Total DCR-owned acres remaining for regenerating up to the 25% per 10 year limit for this subwatershed	Acres in this sub-watershed that are part of this proposed lot
Upper East Fever Brook	1017	32.8	221	61
Lower East Fever Brook	845	0	211	27

Additional comments on Subwatershed analysis: None.

#### Wildlife and Habitat Observations and Considerations

Wildlife/Habitat	Observations and Considerations	
Natural Heritage	There's 0.57 acre of NHESP Priority Habitat in the southwest corner.	
<b>Priority Habitats?</b>		
State Listed species	NHESP has determined that certain state-listed sensitive species or habitats may exist within	
present:	the lot proposal area. To protect them from unnecessary disturbance, detailed information regarding affected species and their locations is not included in this report. DWSP will coordinate with NHESP and follow recommendations to protect these species during the proposed activity.	
Rare Natural	None known.	
Communities:		

Wildlife/Habitat	Observations and Considerations	
General Wildlife Comments	Verified Vernal Pool #13 will be protected by a 200 foot low ground disturbance zone, as required by the 2017 DWSP Land Management Plan; the proposal boundary has been drawn to exclude the 15 foot no cut buffer and the 100 foot shade zone.	
	Dead and dying trees (snags) will be retained and protected whenever possible for wildlife habitat. Large diameter snags and logs, which provide habitat for broad suites of species and are relatively uncommon in the general landscape, will be prioritized for protection.  Wildlife/sign observed on site: bobcat, raven, porcupine, moose and deer.	

**Cultural Resources Description and proposed protection measures** 


<b>Cultural Resource</b>	Description and proposed protection measures
Historical features present; comments regarding protection	A fragmented stone wall borders part of the eastern portion of the DCR access road and may need to be crossed for the landing, as described above. Additional stone walls are located in the northeast corner of the proposal and will be avoided and protected, as will two stone piles in the north third of the proposal.  The Rollin N. Doubleday Sawmill was located on Fever Brook outside the proposal to the southwest.  Any additional cultural features that are located before or during the harvest will be mapped, photographed, flagged, avoided and protected, consistent with the guidelines in the 2017 DWSP Land Management Plan.
Description of site characteristics in relation to Ancient sites modeling or other verified evidence	Slopes are mostly 10-15%, with small areas near Fever Brook and on the hill tops closer to 5%. Aspect is mostly west, but varies from southwest to north. Microtopography is not pronounced.

#### Massachusetts Department of Conservation & Recreation

Division of Water Supply Protection Office of Watershed Management



#### PT-26-09 -- Locus Map






## Massachusetts Department of Conservation & Recreation Division of Water Supply Protection

Office of Watershed Management



### PT-26-09 -- Stand Map






#### Massachusetts Department of Conservation & Recreation

Division of Water Supply Protection Office of Watershed Management



## PT-26-09 -- Soil Drainage Classes






## Massachusetts Department of Conservation & Recreation Division of Water Supply Protection

Office of Watershed Management



## PT-26-09 -- Wetlands and Wildlife Resources





#### Massachusetts Department of Conservation & Recreation

Division of Water Supply Protection Office of Watershed Management





## PT-26-09 -- Cultural Resources and Landscape Characteristics



