# Massachusetts Department of Conservation and Recreation Division of Water Supply Protection, Office of Watershed Management Forest Management Project Proposal Summary for Public Comment

Location, goals, and summary of proposed forest management.

Location, goals, and summary of proposed forest management.					
Proposal Summary Item	Item Information/Description				
Lot Proposal ID	WA-26-85				
Fiscal Year	2026				
Watershed	Wachusett				
Town(s)	West Boylston				
Forester	Greg Buzzell				
Estimated Acres by	12 acres in regeneration patch cuts				
Treatment Type					
<b>Total Proposal Acres</b>	38.2				
Block	n/a				
Compartment and/or	85				
Working Unit					
Location and Boundary	The south and west side are bound by property boundary line; the north side by an				
Description	internal stone wall and partially by property boundary line and the east side by				
	Waushacum Street.				
Previous Proposal?	No				
Project Goals and	This 38-acre forest is part of the original lands that were acquired when the reservoir				
Summary Description	was constructed. Forests provide exceptional water quality protection and yield high-				
	quality water. Active forest management can increase the resistance and resilience of				
	these watershed protection forests to disturbance by deliberately diversifying forest				
	age structure and species composition.				
	The primarily red oak forest in this area is not sufficiently diverse, particularly in age				
	structure with 100% greater than 80 years old. This operation will result in up to 12				
	acres of young forest through the removal of the older overstory in patches. These				
	patches will range in size up to 2 acres and will be located throughout the area taking				
	advantage of where young seedlings and saplings of diverse species are already				
	present.				

#### **Forest Cover Types and Acreages**

Overstory Forest Types	Acres
White Pine - Oak	2.8
Northern red oak	32.7
White pine	1.5

### **Understory Cover Types and Relative Importance**

Understory Cover Type	Relative area covered (Dominant, Secondary, Minor, None)			
Tree seedlings and saplings	Dominant			
Mountain laurel	None			
Mesic site - witch hazel, highbush	Secondary			
blueberry				

Understory Cover Type	Relative area covered (Dominant, Secondary, Minor, None)
Dry site -Huckleberry, blueberry	None
Mesic site - cinnamon fern, mixed	None
hardwood	
Hayscented fern	None
Invasive shrubs/vines	Minor
Other	None

### **Forest Vegetation Description**

orest Vegetation Description  Vegetation Topic Description					
Vegetation Topic  General Description, Forest Composition, Stand History, and Harvest History	At the time of reservoir construction around 1900, this area was described as oak and chestnut sproutland. Prior to that, it was presumably a pasture given the stone walls and hilly character. In the far southwest corner, three small areas were planted, each less than an acre in size. One was planted with white pine in 1918. The second with Norway spruce in 1923 and the third with white pine and Scots pine in 1943. In the far northeast corner on a small flat area at the base of the hill adjacent to John Dee Rd., a tree nursery was established following the closing of the Lamson nursery in Boylston. These nurseries were where over 4.5 million seedlings were grown and planted throughout the properties that were first acquired at the time of reservoir construction. White pines were planted here in 1931 after the nursery was abandoned.  Today, the forest in these 38 acres is almost completely comprised of red oak, black oak and white oak along with far lesser numbers of pignut hickory, red maple, black birch and sugar maple. There are significant numbers of white pine in the northwest corner of this area.  A forest management operation took place on these same 38 acres in 1982. It was a fairly heavy partial removal of the overstory designed to encourage the establishment of regeneration. It may have also been, in part, a salvage of dead oaks resulting from the historic spongy moth infestation of 1981. This was followed in 1989 by a severe thunderstorm that did scattered damage, blowing down and snapping off mature trees. This damage is heavier in the north end with progressively less farther south. The result of these disturbances is an understory and midstory of a variety of primarily hardwood species throughout much of this area. Present are red maple, red oak, black birch, black oak, hickory (both shagbark and pignut), white pine, white oak, sugar maple, sassafras and black cherry. In the south end, there is a lot of spruce regeneration due to the small Norway spruce plantation. In an area particularly heavily c				
Advance Regeneration description	Sampling found that there is adequate regeneration in 44% of the plots and these are well distributed throughout the area. Marginally adequate regeneration was found in 30% of the plots. Present are red maple, red oak, black birch, black oak, hickory (both shagbark and pignut), white pine, white oak, sugar maple, sassafras, black cherry, paper birch, gray birch and spruce.				
Terrestrial Invasive Plants description	Sampling found no invasives however oriental bittersweet was observed in the area of the old nursery.				

### **Description of Wetland Resources Present**

Resource Type	Description of resources present	
Wetlands	None	
Streams	None	
Vernal pools	None	
Seeps	None	

### **Description of Soils by Hydric Class**

Soil Hydric Classes	% of area	Soil series and any further comments
Excessively Drained	7	Merrimac and Hinckley sandy loams
Well-drained Thin	0	
Well-drained thick	93	Canton fine sandy loam
Moderately well-drained	0	
Poorly to very poorly drained	0	

### **Proposed Silvicultural Activities**

Proposed Silvicultural Activities						
Topic	Description					
Site Selection and	This working unit was selected due both to the lack of age diversity in the forests of this					
Silvicultural	subwatershed and in this working unit itself. Most of this area is within subwatershed #4					
Objectives	(Thomas, Quinapoxet and Stillwater Basins). Only 10% of the forest stands within this					
	subwatershed are 20 years old or less. Within the 38 acres of this working unit, none are 20					
	years old or less while 100% of the stands are more than 80 years old with 96% more than 100 years old.					
	The age structure of this working unit is as follows: 0%, 0-20 years old; 0%, 21- 40 years; 0%,					
	41-60 years; 0%, 61-80 years; 4%, 81-100 years; 96%, 100+ years old. The oldest stands date to about 1895 making them 130 years old.					
	Given the lack of young stands in this area and given the presence of good advance					
	regeneration comprised of species well suited to this site, the primary goal will be to increase the proportion of young forest stands in this area.					
Silviculture	With the goal of having a forest with a diverse age structure with at least 3 age-classes					
Prescription	distributed throughout both the subwatershed and within this specific area, the older					
	overstory will be removed in patches. These patches may total as many as 12 acres					
	which would be 1/3rd of the 38.2 acres manageable acres in this area. They will range in size from as small as 1/5th acre up to a maximum of about 2 acres. These will be as well					
	distributed throughout the working unit as possible, taking advantage of where the advance					
	regeneration is present. The Norway spruce, though non-native, provides some unique					
	conifer cover and will be retained.					
	After the operation, the age structure of the forest is estimated to be: 33%, 0-20 years old; 0%, 21-40 years; 0%, 41-60 years; 0%, 61-80 years, 4%, 81-100 years and 63%, 100+ years old.					

Climate Change Considerations: DWSP has determined that the decision to implement this project is consistent with EEA climate goals and guidelines and agency land management objectives. Carbon and climate change considerations specific to the activities proposed for this project are discussed below.

Proposed Activity	Alignment of Activity with Climate Oriented Strategies and Recommendations		
Patch Regeneration Cut (see page 3: Silviculture Prescription)	Patch cutting is a regeneration technique that straddles the boundary between classic even-aged and uneven-aged forest management systems. Foresters select appropriate areas ('patches' or 'groups') covering a portion of the stand to harvest rather than removing the entire stand and then return periodically to repeat the process in other portions of the stand. In using patch cutting there is no final regeneration cut. Patch size and shape are determined by many different factors including overstory condition, desired species composition in the regeneration layer, other desirable herbaceous and woody vegetation, location, stand re-entry period, etc. Harvesting in patches aligns with many climate-smart forestry practices:		
	<ul> <li>Increasing structural diversity improves resiliency by reducing the impact of age/size related disturbances.</li> </ul>		
	<ul> <li>Extending regeneration periods minimizes short term impacts to groundwater and nutrient cycling.</li> </ul>		
	<ul> <li>Partial stand overstory removals more closely align with natural disturbance patterns.</li> </ul>		
	<ul> <li>More carbon is left on the landscape for longer periods, and within- patch live tree, snag, and coarse debris retention allow for development of old forest characteristics.</li> </ul>		
	Can also be used as opportunities to increase the stocking of <b>future climate adapted species, current climate imperiled species</b> , or other types of desirable vegetation.		
General/other Climate Change Considerations	This silvicultural approach focuses on enhancing forest resilience through age-class diversification and regeneration of climate-adapted native species. With 96% of the forest over 100 years old and no stands under 80 years, the management unit is structurally vulnerable to disturbances like pests, storms, and droughts, particularly under a changing climate. The proposed patch cuts mimic the natural disturbance pattern of these past disturbances by selectively removing overstory to take advantage of areas with advance regeneration in a planned, strategic manner. The advance regeneration includes species well-suited to warmer and more variable future climates, such as red oak, hickory, black birch, and white pine. The inclusion of pioneer species like paper and gray birch further improves adaptive capacity and rapid canopy closure, helping to stabilize microclimates and reduce erosion.		

### **Equipment and Access Constraints and Considerations**

Constraint Topic	Description and Considerations				
Proposed Equipment requirements	Mechanized felling and forwarding will be required.				
Proposed wetland or	None needed.				
stream crossings					

Constraint Topic	Description and Considerations
Further wetland	None
comments	
Vernal Pools	None present
Access improvements	None needed
needed	
Other EQ issues	None
In-kind Services	None
Other Access	None
Concerns (parking,	
trails, etc.)	

**Subwatershed Analysis** 

	ub-Watershed number/name	Total DCR- owned acres in this sub- watershed	Acres regenerated on DCR land in the last 10 years in this subwatershed	Total DCR-owned acres remaining for regenerating up to the 25% per 10 year limit for this subwatershed	Acres in this sub-watershed that are part of this proposed lot
4/	Stillwater Basin	599	19	150	38

### Additional comments on Subwatershed analysis:

### Wildlife and Habitat Observations and Considerations

Wildlife/Habitat	Observations and Considerations
Natural Heritage	None
<b>Priority Habitats?</b>	
State Listed species	None known.
present:	
Rare Natural	None known.
Communities:	
<b>General Wildlife</b>	No unusual wildlife sightings were recorded during the site visits. Tree cavities of various sizes
Comments	were observed throughout the lot and will benefit nesting birds or denning mammals. Dead
	snags and live trees with large cavities or loose bark will be retained to the extent possible for
	wildlife value. A few raptor stick nests were observed though none appeared to be active.

**Cultural Resources Description and proposed protection measures** 

<b>Cultural Resource</b>	Description and proposed protection measures
Historical features	There's a stone wall in the north end of this area. It is not anticipated that this wall will need
present; comments	to be crossed.
regarding protection	

Cultural Resource	Description and proposed protection measures
Description of site	Nearly this entire site is greater than 7% sloped and much of it is more than 15% sloped.
characteristics in	There is significant microtopography due to both the stony nature of the soil and numerous
relation to Ancient	pit-and-mounds resulting from the storm of 1989.
sites modeling or	
other verified	
evidence	


## Executive Office of Energy and Environmental Affairs

### Massachusetts Department of Conservation & Recreation

Division of Water Supply Protection Office of Watershed Management



**WA-26-85 -- Locus Map** 






Division of Water Supply Protection Office of Watershed Management



### **WA-26-85** -- Stand Map








Division of Water Supply Protection Office of Watershed Management



**WA-26-85 -- Soil Drainage Classes** 









Division of Water Supply Protection Office of Watershed Management



### WA-26-85 -- Wetlands and Wildlife Resources











## WA-26-85 -- Cultural Resources and Landscape Characteristics

