



DIVISION OF FISHERIES & WILDLIFE

Halfway Pond WMA Habitat Restoration Project Summary SE-HP-TS1

Location

Site: Halfway Pond WMA

Town: Plymouth

District: Southeast District

Project Area

250 Acres

MassWildlife's Approach to Habitat Management

MassWildlife uses habitat restoration and management to conserve both common wildlife and vulnerable species, including rare plants and animals protected by the Massachusetts Endangered Species Act (MESA) and other declining Species of Greatest Conservation Need (SGCN) identified in the Massachusetts State Wildlife Action Plan (SWAP). As part of this effort, biologists and ecologists plan and implement projects to improve, restore, and maintain a variety of healthy habitats to increase biodiversity and climate resiliency across our forests, wetlands, streams, fields, and more.

Biologists plan habitat projects that may include tree cutting, mowing, and mulching to strategically increase open habitats, promote patches of vigorous young forest, restore natural processes, or remove invasive plants. This project has been designed to ensure consistency with recommendations for climate-oriented forest management provided by the Climate Forestry Committee ([Climate Forestry Committee Report, 2024; see below](#)).

Site Significance

Halfway Pond Wildlife Management Area (WMA) is located within a larger collection of conserved lands that is one of the most important landscapes in southern New England for preserving regional biodiversity. Halfway Pond WMA and the surrounding landscape support critically imperiled pine barrens communities. Landscape management is necessary to ensure that the integrity of the pine barrens communities in the Halfway Pond area is strengthened and preserved. Restoration actions will allow these communities to persist and continue to support the specialized wildlife that have evolved to rely upon them.

Pine barrens are considered to be disturbance dependent communities; they require periodic disturbance events to maintain their species composition and open structure. Historically, disturbance in the Halfway Pond area came in the form of regular fire events that occurred over the millennia following the retreat of the last ice sheet. Described in the simplest terms, regularly occurring fire on the landscape will result in: (a) the reduction/removal of many of the generalist plant species that are not capable of tolerating fire; (b) the establishment and persistence of plant communities that tolerate/take

advantage of fire; (c) the removal of accumulated organic fuels; and (d) the continual reset of the structure of these communities that keeps them open and favorable to the species that specialize in them. Conversely, fire exclusion policies have disrupted the natural fire cycle. Now that meaningful fire has not occurred in many areas within the Halfway Pond area in over 50 years, the integrity of pine barrens communities has diminished, the populations of their associated species are in decline, and a major public safety threat from wildfire is building due to the high level of accumulated fuels and closing canopy.

This project will build on past successful habitat work in the greater Southeast Pine Barrens area. Targeted management will add ecological diversity to this already-rich landscape and will directly benefit plants and animals identified as SCGN in the Massachusetts SWAP, along with common wildlife.

Project Activities and Expected Outcomes

Selective tree harvest will be conducted on approximately 250 acres. Tree harvest will occur at varying intensities to stimulate pine barrens shrub layer growth. Individual trees or clusters of trees with high habitat value will be retained to provide food and/or winter cover to wildlife. Pitch pine and tree oak species will be retained. Whole tree removal is the preferred treatment method, but mulching/mowing will be used in some areas where this is not feasible due to dense thickets of pitch pine under 4 inches in diameter. Project design will also establish fire breaks.

Project planning and oversight will be implemented by a team of experienced habitat biologists and restoration ecologists. This project is being done in compliance with MESA Conservation and Management Permit No. 011-135.DFW as amended.

This project will build on past successful habitat work in the area. Planned activities will develop open habitats and diversify the habitats currently available in the area. The health of these pine barrens ecosystems is crucial, as they provide homes for many highly specialized and declining plants and animals, including over 40 species listed under MESA, which are often found only in pine barrens.

Highlights:

- Tree removal will promote the growth of a dense shrub layer across a variety of habitat types that include pitch pine-oak woodland, scrub oak shrubland, sandplain grassland, and heathland. This will provide the high-quality habitat needed for over 40 MESA-listed species and expand nesting, foraging, and cover important for many other wildlife species. This work will also provide high-quality habitat for white-tailed deer, wild turkey, and quail.
- The management practices used in this project will help mitigate the threat of infestation from the invasive southern pine beetle.
- Tree cutting and the use of occasional prescribed fire for long-term habitat maintenance will promote the growth of native barrens understory vegetation, like scrub oak, black huckleberry, and lowbush blueberry, that will provide high-quality habitat for vulnerable and common insects and other types of wildlife.

Climate Considerations

This project was designed to ensure consistency with recommendations for climate-oriented forest management provided by the Climate Forestry Committee and includes:

- Tree thinning to improve this fire influenced landscapes habitat conditions;
- Tree thinning to mitigate against catastrophic wildfire;
- Tree thinning to decrease tree density reducing vulnerability to wildfire, harmful insects, like the southern pine beetle; and
- Restoring native species that are best adapted to the site promotes resilience to future drought, wildfire, and harmful insects.

See page 5 for more details.

Project Proposal Map




Figure 1. Map of Halfway Pond WMA with highlighted project area.

Climate Consideration Details

MassWildlife has determined that the decision to implement this project is consistent with EEA climate goals and guidelines and agency land management objectives. Carbon and climate change considerations specific to the activities proposed for this project are discussed below.

Proposed Activity	Alignment of Activity with Climate Oriented Strategies and Recommendations
<p>Access improvements (landing improvements, gravel, road grading, ditch maintenance, road widening, straightening, and alteration of intersections).</p>	<p>Roads, landings, and associated infrastructure are critical for access by both the public and natural resource managers. These infrastructure elements are also associated with both vulnerabilities and opportunities in terms of climate change resiliency.</p> <p>Vulnerabilities:</p> <ul style="list-style-type: none"> • Roads occupy areas that would otherwise be carbon rich forest. • Road edges can become avenues for the spread of invasive species. • Roads have the potential for sediment transport into surface water resources. <p>Opportunities</p> <ul style="list-style-type: none"> • A well-designed and well-maintained access system makes all other land management and monitoring activities possible while minimizing impacts. • Roads provide for public access including hiking, hunting fishing, etc. • Roads are critical for both emergency response (injuries, accidents, etc.) and incident stabilization (fire, flood, storm damage, etc.) <p>Given the predicted increase in storm frequency and intensity, improving and maintaining roads, road surfaces, and stormwater infrastructure is imperative.</p> <ul style="list-style-type: none"> • Proper surfacing, grading, and ditching minimize erosion from stormwater and snowmelt. • Periodic maintenance is required to avoid water channelizing within compacted tire paths. • Adding gravel or other material to the road surface helps support the heavy vehicle traffic associated with forestry work, fire operations, and post-storm recovery efforts. Alterations (widening, straightening) are often needed to upgrade old, narrow farm lanes to meet modern vehicle access needs. • Ditching, cross culverts, and relief cuts can be designed with future storm intensities in mind and should minimize, to the greatest degree possible, impacts to surface water resources. <p>Most log landings are temporary in nature. Permanent landings that are properly located and well-built can serve as permanent access infrastructure, concentrating activities and minimizing the non-forested footprint required to conduct agency management. Whether temporary or permanent, the use, maintenance, and stabilization of landings will include considerations of future climate change impacts. Landing BMPs include:</p> <ul style="list-style-type: none"> • Post-harvest stabilization measures such as grading and smoothing to prevent erosion and sedimentation.

	<ul style="list-style-type: none"> • Seeding to provide cover and further stabilize the soil. • Invasive plant survey and control to minimize further infestation risks. • Periodic mowing of permanent landings to allow herbaceous and shrubby vegetation to dominate the site between harvests, adding diverse habitat opportunities for local wildlife.
<p>Invasive plant control, including pre- and/or post-harvest and follow up treatments.</p>	<p>Strong consensus exists among land managers and climate science experts regarding the threat to future forest health posed by the introduction and spread of invasive plants. Invasive plants can:</p> <ul style="list-style-type: none"> • aggressively outcompete native plant species, • dominate understory communities, and even climb, kill, and topple mature trees, • threaten overall biodiversity, and • threaten soil health and long-term carbon storage. <p>Monitoring and controlling invasive and interfering plant populations prior to and following forestry operations is a critical practice for minimizing the risk of further impacts inadvertently (though not unexpectedly) spread by harvesting-related activities.</p>
<p>Habitat restoration and maintenance prescribed fires—heath, shrubland, woodland, or grassland.</p>	<p>Prescribed fire is the planned use of fire in a particular place and time, under established conditions and safety requirements to accomplish resource management goals.</p> <ul style="list-style-type: none"> • Prescribed fire improves habitat for a variety of wildlife and native plants and restores natural communities dependent on fire. • In fire-influenced natural communities, fragmentation of the landscape and the suppression of fires (prescribed or natural) leads to accumulation of volatile hazardous fuels in the surface, mid-story, and canopy vegetation layers. • Excessive vegetation density negatively impacts the habitat quality of the natural community and may eventually lead to fuel buildup and unplanned, catastrophic wildfire. • Prescribed fires that reflect natural return intervals increase below-ground carbon storage and sequestration. <p>The consequences of catastrophic wildfires include:</p> <ul style="list-style-type: none"> • the release of large amounts of carbon including soil carbon, • tree mortality, • severe soil, duff, and below ground vegetation impacts, • potential alteration of soil chemistry, • threats to firefighter safety, human communities, and property damage, and • threats to human health from severe smoke impacts both locally and potentially at long distances.
<p>Establishing and/or maintaining fuel/fire breaks.</p>	<p>Climate models predict drought and wildfire potential increasing in the region due to climate change, and the agency is adopting strategies to both reduce the risk of catastrophic fire spread and maintain fire-adapted habitats. Fuel breaks</p>

	<p>and fire breaks are essential tools for both prescribed burning and wildfire control.</p> <p>Fuel breaks are:</p> <ul style="list-style-type: none"> • vegetated areas, • maintained at lower structure and density, • designed to slow the spread of fire, • designed to control prescribed fire or wildfire, and • opportunities to encourage open woodland, shrubland, or grassland natural communities. <p>Fire breaks may be natural or constructed barriers to the movement of fire, with some examples being:</p> <ul style="list-style-type: none"> • open water, • paved roads, • graveled woods roads, • trails, and • periodically mowed paths (“fire lines”). <p>The fuel and/or fire breaks proposed in this project were designed as part of agency planning efforts for fire control and management for this area within a framework of reducing climate vulnerability.</p>
<p>Tree mowing (for maintaining aspen, e.g.)</p>	<p>Mowing is used to perpetuate both non-forest (grasslands) and young sproutland forest conditions within a given footprint on the landscape. This practice is one dependable way to provide such critical habitats in the absence of discrete localized natural disturbances such as flooding or fire. Mowing to reset the development of a stand that has grown out of the sapling stage is recommended as one climate-smart alternative to achieve the agency’s young forest habitat goals (rather than harvesting mature forest stands over time in a mosaic of large adjacent patches). Although mowing and mulching trees in place aligns with carbon objectives and avoids harvesting older forest, it can be prohibitively costly to implement.</p>
<p>Diffuse overstory removal, partial cut, habitat modification/maintenance.</p>	<p>Open woodlands, savannas, barrens, and heathlands are low tree-density, fire-dependent forests with diverse understory vegetation critical for conserving many state-listed rare species. They are imperiled across Massachusetts due to development and negative ecological alterations resulting from a lack of management primarily decades of fire exclusion. Climate experts recommend prioritizing and maintaining sensitive or at-risk species and habitat, with the expectation that pressure on these will only increase with changing climate. Ecological restoration of these sites ensures continued habitat function and reduces climatic vulnerability.</p> <ul style="list-style-type: none"> • Reducing tree density reduces vulnerability to pests like southern pine beetle and to drought stress. • Restoring native species that are best adapted to the site promotes resilience to future drought, wildfire, and harmful insects. • Reintroducing low-intensity fire promotes resilient native vegetation. • Removing heavy fuel loads reduces vulnerability to wildfire. • Restoration better positions these sites to adapt to climate change. • Restored sites are more reliable carbon sinks in the long term than highly vulnerable dense fire-excluded forests.

	<p>The agency recognizes that this site may store less carbon than denser forests in the short term. But climate models predict an increase in disturbance on these sites including drought, wildfire and range expansion of harmful insects that puts a dense fire suppressed forest at greater risk of becoming a carbon source in the long term. Projects like this are undertaken on federal, state agency, and other conservation lands across the Commonwealth, under the guidance of collaborative teams consisting of biologists, restoration ecologists, foresters, and fire management professionals.</p>
--	---