

**RAPID RESPONSE PLAN FOR
HYDRILLA
(*Hydrilla verticillata*)
IN MASSACHUSETTS**



Prepared for the
Massachusetts Department of
Conservation and Recreation
251 Causeway Street, Suite 700
Boston, MA 02114-2104



Prepared by
ENSR
2 Technology Park Drive
Westford, MA 01886



June 2005

**RAPID RESPONSE PLAN FOR HYDRILLA
(*Hydrilla verticillata*) IN MASSACHUSETTS**

Species Identification and Taxonomy	1
Species Origin and Geography.....	2
Species Ecology.....	3
Detection of Invasion.....	3
Species Confirmation	4
Quantifying the Extent of Invasion.....	5
Species Threat Evaluation.....	6
Communication and Education	8
Quarantine Options	10
Early Eradication Options	12
<i>Hand Harvesting.....</i>	<i>12</i>
<i>Suction Harvesting</i>	<i>13</i>
<i>Benthic Barriers</i>	<i>13</i>
<i>Water Level Drawdown.....</i>	<i>13</i>
<i>Application of Fluridone.....</i>	<i>14</i>
<i>Other Options</i>	<i>15</i>
<i>Recommended Options for Early Eradication.....</i>	<i>15</i>
Deciding Which Technique to Apply	16
Control of Established Infestations	17
Prevention of Re-Infestation	17
Summary.....	18
References.....	19

Species Identification and Taxonomy

Hydrilla verticillata (hydrilla) is a submerged aquatic perennial plant. The roots of hydrilla are long and thin, typically whitish to light brown in color. Roots are usually buried in the hydrosol, but may form adventitiously at the nodes. Stems are ascending and heavily branched near the water surface, and horizontal and creeping under the soil. Stems of hydrilla can reach a length of 8.5 meters. Turions are formed infrequently in the axils of the leaves on the upper part of the stem, and on subsoil stolons. Leaves are narrow, 1-2 cm long, and whorled around the stem in groups of 4-8. On the lower stem, leaves may be opposite in arrangement. The leaf margins are serrated, visibly to the naked eye. Flowers are unisexual, less than 6 mm in diameter, and translucent to white in color. Two biotypes of hydrilla plants occur, dioecious and monoecious. Flowers of only one sex are produced on dioecious plants, while monoecious plants produce both male and female flowers. Male flowers grow on a short stalk and are free floating at maturity. Female flowers are composed of six colorless segments, and are 1.2 to 3.0 mm long. Fruits of hydrilla are cylindrical in shape, and 5 to 10 mm long (Godfrey and Wooten 1979).

According to Crow and Hellquist (2000), the following taxonomic characteristics are used to identify *Hydrilla verticillata* to species:

1. Plants submersed, rooted, or fragments free-floating beneath water surface, leaves basal or cauline, sessile.
2. Leaves cauline, short, opposite or whorled, lacking a lacunae band.
3. Leaves 0.6-1.7 cm long, flowers lacking nectaries.
4. Leaves in whorls of (2) 4-6, leaf margins conspicuously toothed, midvein of lower surface often with spine-like teeth (fresh specimens rough to the touch), spathe of staminate flowers spiny.




Figure 1 – A group of hydrilla photographs, from <http://www.wes.army.mil/el/aqua/apis/biocontrol/html/hydrill1.html>

Species Origin and Geography

The origin of *Hydrilla verticillata* is unclear, but it is currently found on every continent except Antarctica and South America. Schmitz et al. (1991) stated that historical reports support Sri Lanka as the origin of the dioecious biotype. Madeira et al. (1997) used DNA analysis of U.S. samples to conclude the dioecious biotype most likely originated on the Indian mainland, while the monoecious biotype originated in Korea. Some botanists believe the origin of hydrilla is on the African continent (Tarver et al. 1978, Mahler 1979), and some consider it to be a native of Australia (Sainty and Jacobs 1981). The dioecious strain was imported to the United States for use in the aquarium trade during the 1950's (Schmitz et al. 1991). The monoecious strain was not detected in the United States until the 1970's (Haller 1982, Steward et al. 1984). The monoecious form for hydrilla has been detected as far north as Washington in the western U.S., and Pennsylvania and Connecticut in the eastern U.S. (Madeira et al. 2000). More recently, populations were found in Massachusetts and Maine (G. Smith, ACT, pers. comm..).




Figure 2 – A distribution map for *Hydrilla verticillata*, from the center for aquatic and invasive plants website (<http://plants.ifas.ufl.edu>).

Species Ecology

Hydrilla grows most often in freshwater lakes, ponds, rivers, impoundments, canals and ditches, under a wide range of environmental conditions. It usually grows in shallow waters, but can grow at depths greater than 10 m. Mahler (1979) reported that hydrilla can tolerate habitats consisting of 33% seawater. Hydrilla grows in both acidic and alkaline environments, and at various trophic levels ranging from oligotrophic to eutrophic. Although hydrilla grows on all types of substrates, it grows best on sediments with high organic content. Hydrilla is adapted to grow under very low light conditions, and therefore can quickly dominate native vegetation (Bowes et al. 1977). Hydrilla can also tolerate a range of temperatures and is winter-hardy.

Hydrilla is well adapted to rapid spread and growth due to various modes of reproduction. Pollination occurs above the surface of the water and its seeds develop into hypocotyles up to 6 mm in length. The hypocotyle produces a short stem at the node along with 3 leaves and a few roots. Hydrilla can also reproduce from rootstocks, turions (both subsoil and on the stem), and vegetative nodes. Entire colonies can be formed from one single node which can produce adventitious roots and quickly spread. A single tuber can produce more than 6,000 new tubers per square meter (Sutton et al. 1992).

Detection of Invasion

In the great majority of cases, hydrilla enters lakes with flow, boats and birds, and the logical places to look first are the mouths of tributaries, boat ramps and areas of higher bird concentrations. While mature hydrilla growths will usually “top out”, reaching the surface and forming a canopy, new infestations may be less obvious and often require underwater examination for early detection. Hydrilla can grow in deep water, and stems can be over 8 meters (26-27 ft) in length. However, it typically gets its start in shallow waters (<3 m or 10 ft deep) and is likely to be visible from a boat with a viewing tube or by snorkeling. Use of an underwater video system (Aqua-Vu or equivalent) can be very helpful in scanning large areas of variable depth.

Sources may not be obvious, but the pattern of occurrence observed during early detection may provide useful clues. Appearance near boat ramps suggests boats as vectors, while appearance in more remote areas with no direct access or inflows suggests birds as the source. Where plants are detected near the mouth of a tributary, it would be appropriate to check the next upstream waterbody or the stream bed itself if conditions are suitable for rooted plant growth.

There are multiple methods of plant survey, and no truly standardized technique. The object is to be as thorough as time and trained manpower allow, to maximize detection probability. To detect a suspected invasion, or simply to monitor for possible invasion, consider the following steps:

1. Acquire a suitable map of the waterbody, showing shoreline features and reference points, and preferably with water depth contours.
2. Use the taxonomic information supplied here, or supplementary information from taxonomic guides, plant keys, or herbarium sheets to identify hydrilla. Be careful not to confuse it with *Elodea* or *Egeria*, two physically similar genera.

3. Hydrilla growths in shallow water and may be easily detected at the surface in late summer, but surveys should include deep water where plants may not be apparent at the surface.
4. Ideally, space transects around the waterbody, extending from shore to the end of plant growth, with one transect per defined shoreline segment, determining transect location with GPS or readily identified shoreline features. Segments should be of roughly equal length, but this can be based on actual shoreline, straight distance across the water, land use or other features of concern or interest, or encompassed waterbody area. Be sure to cover all boat launch, swimming, inlet, bird congregation, key habitat and intake areas, and any other key access points.
5. Priority can be given to transects of key concern, either based on likely invasion points (access points) or potentially threatened resources (intakes, swimming areas, key habitat) if the number of transects is too great for the manpower and time resources available, but recognize the limitations this will impose on invasion detection.
6. Using a boat with a viewing tube or underwater videocamera, or employing snorkeling or SCUBA gear, examine the plant community along transects between the shore and the maximum depth of plant growth (typically <30 ft, usually <20 ft). Note presence/absence of hydrilla and extent of coverage and density where hydrilla is encountered. Record observations for 2 ft water depth intervals, with each observation representing either a defined area within the depth range or the length of the transect between depth intervals (typically 0-2 ft, 2-4 ft, 4-6 ft, and so on). GPS is particularly useful for both transect and point location for future reference.
7. Tabulate all data in a manner that facilitates future comparisons, typically in a spreadsheet or GIS format. Evaluate presence of any hydrilla, extent of coverage and density, and pattern of occurrence. Map the distribution of hydrilla in the waterbody for visual reference.
8. Repeat the survey at least once every 3 years (about the time for an invasion to have a detectable impact), and preferably every year to allow the earliest possible detection.

Species Confirmation

Unless the invasion is discovered by individuals trained in plant taxonomy, samples should be sent to competent taxonomists for confirmation. In Massachusetts, the Department of Conservation (DCR), the Department of Environmental Protection (DEP), the Massachusetts College of Liberal Arts (North Adams, specifically Dr. Barre Hellquist), and the University of Massachusetts at Amherst (UMASS) have the expertise to assist in plant identification. Many consulting and lake management firms also possess this expertise, but it will be the responsibility of the DCR to determine where specimens should be sent. Therefore, the DCR at 617-626-1411 or 617-626-1395 should be the first point of contact.

Key steps in confirming an invasion include:

1. Collect complete specimens of the suspected hydrilla; try to collect the root system as well. Place the specimen in a clear container with water for easy viewing (clear 2-L soda bottles without labels work well); keep chilled. Alternatively, specimens can be pressed on a sheet of appropriate (absorbent) paper, covered with wax paper and a stack of books or other suitable weight (an actual herbarium press is useful if available).

2. Contact the DCR representative at 617-626-1411 or 617-626-1395 and inform him/her that a suspected occurrence of hydrilla has been detected in the waterbody. The DCR contact will assess past records for the waterbody and will instruct the caller where to send a sample for confirmation, if warranted.
3. As soon as possible, preferably within 2 days, send specimens to the identified DCR representative for confirmation, or to a taxonomic expert as designated by the DCR contact. Note in writing that the enclosed specimen is believed to be hydrilla and include the name of the waterbody, the approximate location in the waterbody (a map is helpful) with water depth and any other site-specific observations, the date and time of collection, and the name, address, phone number and email for the collector or sender.
4. The DCR will confirm the identification or provide an alternative identification either directly or indirectly through a recognized taxonomist, and will be responsible for notifying all appropriate agencies, municipalities and citizen groups either potentially affected or responsible for follow-up actions.

Quantifying the Extent of Invasion

Gaining effective control of hydrilla depends on detecting all growths, as this species can expand rapidly. The initial discovery may be made during a routine mapping exercise, but mapping approaches suitable for overall plant assemblage characterization (e.g., point intercepts on a grid or transects) may not be appropriate for thorough coverage of recent invasions. Where a growth is detected, it is likely that expansion in the first growing season will be by tubers and turions, so viewing each discovered growth in concentric circles moving outward from the apparent center will best facilitate mapping of the growth. Detection of additional growths is best accomplished by a thorough visual inspection of the newly infested area, either using tightly spaced transects radiating out from the first discovered growth or focused in the direction of likely current or wind transport.

If the waterbody is large, effort may have to be limited to the most likely locations for invasion. In this regard, examination of any existing plant maps may be helpful. Look for areas of suitable depth (<30 ft, with emphasis on areas 2-20 ft deep), known plant and bottom disturbance (marinas, boating lanes, windswept shallows), and plant assemblages of lower density and/or lesser canopy formation.

Evaluation of recent hydrilla growths should focus on extent of coverage and degree of dominance. Biovolume or biomass measures are useful but time consuming and are not critical to combating new infestations. Careful stem counts are helpful in assessing the efficacy of possible controls, but are also time consuming. An estimate of stems per unit area and the area covered is more valuable in assessing potential controls for new growths. With regard to dominance, it is important to note other species present, as the presence of protected species and the relative abundance of seed producers vs. vegetative propagators are important to planning management actions. A list of plant species with an approximation of the percent of the community each represents is appropriate.

Assessing the rate of expansion may not be necessary if the invasion is detected early and prompt control actions are implemented. However, where hydrilla has been present for more than a single growing season, information on the rate of expansion may be helpful in planning a control strategy

and in garnering support for rapid action. Isolated plants are likely to signal the first year of growth, while scattered plants are likely to represent the second year of growth and well established beds will normally be more than two growing seasons old.

Useful steps in quantifying the invasion include:

1. Use the data generated by the transect method in the section on Detection of Invasion to get a first impression of the extent of invasion, preferably in mapped format. Where hydrilla is discovered in multiple locations, look for spatial patterns that suggest either transport from the earliest infestation or invasion from multiple sources.
2. If a discovered growth is in a definable cove, examine the entire cove, or at least that portion with a water depth <20 ft.
3. If a discovered growth is associated with a boat ramp, check a suitable area (typically 1-2 acres) associated with that ramp, and check other ramps if present.
4. Where growths occur near a tributary mouth, check area maps for upstream ponds or impoundments on the offending stream and any other tributary and investigate where possible hydrilla sources seem most likely.
5. When the new growth appears associated with areas of bird congregation, check all such areas in the waterbody.
6. In all cases, note which areas have established beds vs. scattered plants vs. a single plant or just a few stems.
7. Identify all other plants in association with hydrilla growths, to the limit of areas likely to be targeted for control. Follow the protocols for species confirmation where specimens of unknown identity are encountered, paying particular attention to possible protected species or other invasive species.

Species Threat Evaluation

Hydrilla verticillata is on the 1979 federal noxious weed list (USDA-NRCS 1999), as well as the noxious weed lists of many U.S. states. The ability to spread quickly and grow in a variety of environmental conditions allows hydrilla to out-compete native vegetation and quickly dominate a waterbody (Bowes et al. 1977). Hydrilla often forms dense mats at the surface of the water, and Haller and Sutton (1975), reported that 20% of the biomass is located in the upper 10 cm of these mats. These large mats displace native vegetation (Haller 1978), reduce biodiversity, and may alter ecosystem balances, including food chains and trophic interactions (Westman 1990, Schmitz and Simberloff 1997). Large mats can also alter water quality features (Smart and Barko 1988).

Aquatic macrophytes can provide food, shelter and spawning habitat for a wide variety of fishes (Lillie and Bud 1992). Intermediate densities of aquatic macrophytes, including hydrilla, enhance fish diversity, feeding, growth and reproduction (Langeland 1996, Pimental et al. 2000). Hydrilla tends to replace native macrophytes, however, creating food shortages for fishes (Engel 1995). Beds of aquatic macrophytes can also impede predation, shelter panfishes, and cover spawning areas, leading to potential decreases in sportfish abundance (Engel 1995). The amount of hydrilla in the water column dictates the presence and size distribution of fish species (Killgore et al. 1993,



Harrel et al. 2001). The depletion of oxygen in waterbodies with dense macrophyte coverage can also result in fish avoidance, and in extreme cases fish kills (Holland and Huston 1984, Engel 1995).

Dense mats of hydrilla limit human uses of the waterbody and can lead to decreased water quality. Dense mats choke channels, clog water intakes, and restrict aquatic activities such as fishing, swimming and boating. Hydrilla can have economic impacts on industrial and recreational companies. Thick mats can reduce flow rates in irrigation operations by as much as 90% (CDFA 2000a). Hydroelectric power generation and irrigation operations are hindered as plant material builds up in trash racks and clogs intakes. Boat marinas have been closed due to thick mats of hydrilla which make waterways impassable by motor boats.

The economic impacts of a hydrilla infestation can be damaging. A California study showed that the cost of management increased linearly and rapidly as the area affected by pest plant infestations increased (Rejmanek and Pitcairn 2002). Due to the rapid growth of hydrilla and its ability to quickly dominate the macrophyte community, costs associated with its control and eradication quickly increase, so responses to identified infestations must be quick.

Potential spread within the waterbody is governed by the physical features of the waterbody (especially water depth and substrate) and the level of activity of potential vectors of spread for hydrilla (especially boats, birds, flow and currents). Hydrilla can grow on nearly any substrate, from rock to loose muck, but it prefers substrates with high organic content. However, all areas of the waterbody are suitable habitat with regards to substrate. Rocky to gravelly substrates typically support lesser densities of macrophytes and very loose muck provides an unstable substrate where growths may be variable over space and time. The depth range for hydrilla is from shore to about 30 ft. Boats and birds can actively transport hydrilla within a waterbody, but hydrilla may also create fragments that drift with currents. The production of tubers and turions with seemingly erratic germination rates further complicates control of this species.

Potential spread outside the waterbody is mainly a function of surface outflow, birds and human activities. Overflow can carry viable fragments downstream to additional waterbodies. Birds may transport fragments, but are more likely to carry seeds, either externally or in their digestive tract. Transport by humans is a known threat, with movement of plant fragments in or on boats and trailers well documented (Johnstone et al. 1985, Bratager et al. 1996).

All of these factors combine to create a site-specific level of threat. Of primary interest are how great an infestation may become, how readily it may be transmitted to new areas (both inside and outside the infested waterbody), what resources may be impacted to what degree, and what the potential is for eradication or control through rapid response to detection of an invasion. In evaluating the potential threat from a new hydrilla infestation in DCR parks on a case by case basis, the DCR staff will consider the following:

1. What portion of the waterbody could be colonized (estimate as the area with water depth <30 ft)?
2. What is the potential for dense bed formation (estimate as the area with stiff muck or sandy silt substrate)?



3. What is the potential for rapid (<3 years) spread of hydrilla (estimate as the common area from #1 and #2 above and not densely covered by native plants)?
4. What is the potential strength of vectors of internal hydrilla spread (boat traffic, flow, currents, open expanses vs. isolated coves)?
5. What is the potential strength of vectors of external hydrilla spread (trailered day-use boats, daily or seasonally mobile bird populations, outlets without screening)?
6. What resources and uses are potentially threatened (water supply, swimming, boating, fishing, aesthetics, sensitive or protected populations)?
7. What is the potential for eradication (based on extent and density of coverage, vectors of spread)?
8. What is the potential for confinement (based on extent and density of coverage, physical isolation of area affected, vectors of spread)?

By answering these questions, one can characterize the threat according to the following matrix, which can then govern the response to detection of an invasion:

FACTOR	YES	NO	THREAT EVALUATION	HIGH	MEDIUM	LOW
A large area could be affected			Extent and speed of possible infestation			
Plant density could be high						
Spread could be rapid						
Water supply may be impacted			Nature of possible impacts			
Swimming may be impacted						
Boating may be impacted						
Fishing may be impacted						
Aesthetics may be impacted						
Sensitive species may be impacted						
Protected species may be impacted						
Spread by water flow likely			Ability to spread			
Spread by birds likely						
Spread by boating likely						
Spread by other human activities likely						
Eradication is possible			Potential success of rapid response			
Confinement is possible						

Communication and Education

The most effective method of controlling hydrilla infestations is through prevention. Public education is the key to preventing the spread of hydrilla and other aquatic nuisance plants. A successful education program would target users of all ages and interest. Included in this list are boaters, lake



managers, aquatic pesticide applicators, anglers, and water gardeners. This program should include introduction information for hydrilla, how to identify hydrilla, what to do and who to contact if you find hydrilla, and the threats of hydrilla in Massachusetts.

Once the presence of hydrilla has been confirmed, the town(s) in which the waterbody is situated should be notified, usually through the Conservation Commission, which will have a chairperson or an agent who is reachable through Town Hall. It would also be appropriate to notify all relevant stakeholder groups, but these need to be identified and many will not have a central clearinghouse contact for notification. Groups who should be informed about the infestation include any active lake association, shoreline property owners, boaters, anglers, swimmers, birdwatchers, and water suppliers. Notification through individual contacts is desirable but may be inefficient. Posting a notice in the local paper will help publicize the problem, but the notice may not receive widespread attention. Posting the waterbody at access points is perhaps the most effective approach, as it is the actual users that should be informed and warned to avoid spreading hydrilla.

It is desirable to post access points with warning signs even before an invasion, displaying a picture or drawing of hydrilla and asking waterbody users to be on the lookout for this invasive plant. Users, particularly boaters, should be asked to inspect their boats and any trailers prior to launching, and to remove any discovered plants with proper disposal in a manner that prevents the plant from reaching the waterbody. A local contact (name and phone number) for notification should be given, typically either a representative of the lake association or the town's Conservation Commission, or both. Users should be advised to mark the location where the plant was observed if at all possible, but not to pull it out unless they can get the whole plant, including the roots. As most users will not be diving or snorkeling, immediate, effective hand harvesting is probably not a realistic expectation.

After an invasion has been discovered, it is even more important that access points be posted with a warning to users to avoid any action that could spread hydrilla. Again, a picture or drawing of hydrilla should be provided, and any known locations of the plant should be shown on a map of the waterbody. Users should be asked to notify a local contact if hydrilla is found in other areas not shown on the map, and to avoid motorized boating in areas with hydrilla. All boats, trailers, fishing equipment, bait buckets or other possible means of transport should be inspected and cleaned prior to leaving the waterbody.

Responsibility for control of hydrilla does not rest with any one entity under the laws of the Commonwealth of Massachusetts. Approval for control actions is governed by the Wetlands Protection Act, which always involves the town Conservation Commission and the Commonwealth's DEP. Approximately 16 states have, or are in the process of developing, laws restricting the transport and/or possession of hydrilla, but Massachusetts has not so far joined this group. Approval for control actions may also involve the Division of Fisheries and Wildlife and/or the Natural Heritage and Endangered Species Program, both agencies of the Commonwealth, depending upon the resources in the waterbody (particularly if protected species are known from the waterbody). Other agencies and approval programs may apply, depending upon the features of the waterbody (naturally large enough to be a statutory Great Pond), the location of the waterbody (e.g., in an Area



of Critical Environmental Concern), or the uses of the waterbody (e.g., as a water supply). However, none of these agencies is charged with controlling invasive species, and there is no legislation in Massachusetts that mandates control of hydrilla. The DCR has taken the lead in Massachusetts with regard to encouraging control of invasive species, and supports control efforts as its budget allows. However, outside of the state parks and reservations, control is largely a function of local desire to protect and maintain the resource.

For waterbodies within DCR parks, the following notification procedures are to be followed when a new infestation by hydrilla has been confirmed:

1. The DCR contact responsible for confirming the hydrilla invasion will notify the DCR Regional Director, Park Supervisor and any regional DCR contact charged with managing water resources. A single letter copied to each party is preferred. The letter should briefly state the problem and outline immediate control steps that are needed, indicating an expected date for a follow up visit by Lakes and Ponds Program staff to begin concerted control measures (see posting procedures below).
2. The DCR contact responsible for hydrilla invasion confirmation will also notify the DEP, the DFW and the NHESP in writing; a copy of the letter sent to DCR parties is sufficient. If a contact for an associated citizens' lake or watershed organization is known, notification should be given to that group as well.
3. The Regional Director or a designated park contact for local affairs will notify the town(s) in which the park and waterbody are situated. The appropriate parties within the town(s) to be notified may vary by town, but should include the Conservation Commission and either the Selectmen, Town Manager or Mayor, depending upon local government structure.

For waterbodies within DCR parks, the following posting procedures are to be followed when a new infestation by hydrilla has been confirmed:

1. All access points to the waterbody (e.g., boat launches, swimming areas, fishing piers or obvious shoreline fishing points) shall be posted with a photograph or drawing of hydrilla and a written notice that this invasive plant has been found in the waterbody.
2. Suggested language is as follows: Warning. Hydrilla (*Hydrilla verticillata*) has been found in this waterbody. This invasive plant represents a threat to this waterbody and its users. Caution should be exercised to avoid the spread of this plant. Do not pick or remove this plant if you encounter it, and be sure all equipment brought to this waterbody is clean before leaving.
3. Include a contact name and phone number on all postings.

Quarantine Options

Both natural processes and human activities can spread hydrilla, both within an invaded waterbody and to other area lakes. Minimizing the spread of hydrilla may require some form of quarantine. Making the waterbody off limits to all users is an extreme action not typically justified for new growths that are likely to be limited in areal coverage. However, keeping people out of infested areas may be a valid option. This may be done by signage, buoys, or an actual sequestration curtain, with cost increasing dramatically in the listed progression.

Where the invasion is occurring at a boat ramp, closure of the ramp may be justified; this will both limit the spread of hydrilla and generate public awareness of the problem and a desire to take action against the hydrilla. A town may take such an action where the public welfare is deemed to be at stake for a boat ramp owned by the town, but it is not clear that such action is legal for private boat ramps, and towns do not have the authority to close ramps owned by the Commonwealth. Consult with private owners or the Public Access Board of Massachusetts when considering closure of a ramp not owned by the town.

Where the invasion is occurring in a swimming area, closure of that area will have much the same effect and limitations as for boat ramps. If the hydrilla growths are localized, it may be possible to partition off the infested area by moving the buoyed ropes that usually delimit swimming areas. If the growths are extensive, it may be appropriate to close the swimming area on the basis of public safety; people can get tangled in hydrilla and drown.

The use of sequestering curtains or screens can both restrict access to an infested area and limit the spread of hydrilla by vegetative fragmentation. This approach, while often expensive, has been very effective in a number of cases, especially for small areas or coves with a narrow connection to the main body of the waterbody.

Possible expansion routes should be considered and addressed to the extent possible.

Sequestration, as noted above, can be highly effective if the infested area is localized and amenable to curtains or screens. Outlets from the waterbody should also be screened to minimize the export of hydrilla fragments with outflow. This may be problematic where leaves or other debris are abundant enough to clog such screens, necessitating frequent cleaning. Rotating screens or other automated outflow restrictors are effective but expensive. Drawdown may also limit hydrilla escape, if an appropriate subsurface outlet exists and hydrilla can be prevented from passing through it. It may be advisable to implement bird controls to limit bird contact with infested areas; scare tactics (e.g., flags or pinwheels on buoys, noisemakers) can be effective for short time periods, which may be all that is necessary for lakes with migratory populations. Greater effort may be needed for lakes with substantial resident bird populations. If boating is allowed, it is advisable to set up a temporary wash station at any ramp; it may be necessary to staff it to maximize use compliance. At the very least, boats and trailers leaving the waterbody should be inspected and cleaned.

Where a hydrilla invasion is confirmed in a waterbody in a DCR park, the following quarantine steps will be evaluated and implemented as warranted:

1. Screen the surface outlet of the waterbody to minimize downstream movement of hydrilla, maintaining the screen as necessary to facilitate outflow.
2. Lower the water level to prevent surface outflow; a subsurface drain may be used to continue outflow, but hydrilla may escape through this exit if not screened, and such screening will require cleaning.
3. Post access points with warnings to avoid the plant and/or certain areas of the waterbody; use marker buoys to identify infested areas.

4. Surround smaller infested areas with sequestration curtain or other enclosing materials that prevent spread and limit access.
5. Curtain off coves or other isolated areas to prevent hydrilla spread and limit access.
6. Use scare tactics or other approaches to limit bird use of the waterbody.
7. Set up a washing station and inspection point for boats taken out of the waterbody; require inspection and cleaning where needed.
8. Close any access point (e.g., boat ramp, beach, other points of active contact) in close proximity to hydrilla, where the potential for internal or external spread is considered high.
9. Close the waterbody to human use.

Early Eradication Options

Timelines for necessary action with regard to hydrilla invasions hinge on stopping the spread of this plant. Expansion from tubers occurs throughout the growing season, so the sooner controls are implemented, the smaller the area that must be addressed. Once the growing season is over (about October), plants are largely dormant and many collapse or are otherwise reduced in biovolume until the following spring. Detecting and effectively removing hydrilla plants by physical means will therefore be more difficult outside the growing season.

Management options for rooted plants are covered in *The Practical Guide to Lake Management in Massachusetts* (Wagner, 2004), a companion guide to the GEIR on Lake Management, available on-line at <http://www.mass.gov/dcr/waterSupply/lakepond/lakepond.htm> and supplied to all towns in the Commonwealth by the DCR in 2004. However, direct experience with hydrilla control in Massachusetts specifically and New England in general is very limited (about four cases as of spring 2005). A summary of control approaches with the potential to eradicate hydrilla during the early stages of an invasion is provided below.

Hand Harvesting

Mode of action: Plants are removed by divers by hand; removal includes root crowns.

Probability of successful control: Where density is <500 plants per acre over a small number of acres, control can be complete, although regrowth from tubers or turions is likely except in the most recent infestations. At higher densities or area of coverage, risk of incomplete harvest or spread by fragment escape increases dramatically.

Potential non-target impacts: Limited; with training, divers recognize hydrilla and avoid other plants; risk to non-target plants increases as density of plant community increases. Temporary turbidity increases are expected.

Permitting needs: Can be approved without Order of Conditions under the Wetlands Protection Act through a Negative Determination of Applicability (WPA regulations deemed not to apply, as only the invasive plant is removed).

Monitoring needs: Critical to delineate target area and provide means for divers to stay on course with complete coverage. Monitoring during harvesting to detect and collect fragments is also very important to successful elimination of hydrilla.

Range of costs: Often done by volunteers, but estimates from professional operations range from \$100 to \$500 per acre.



Other considerations: Use of a fragment barrier around all harvesting areas is highly recommended. Effective hand harvesting requires careful planning and is more difficult than it may appear.

Suction Harvesting

Mode of action: Plants can be pulled directly into the suction apparatus, but for best effect this is a suction aided hand harvesting operation, whereby hand harvested plants are fed into the suction tube and filtered out in an above-water chamber. This speeds up the operation and limits fragment dispersal.

Probability of successful control: High potential for eradication at low to moderate densities of hydrilla; complete removal probability declines at higher densities.

Potential non-target impacts: May pull in non-target plants and plankton by suction, but effects localized and limited. Turbidity plume at surface from filtering chamber may be substantial.

Permitting needs: Generally requires an Order of Conditions under the Wetlands Protection Act, but may be issued a Negative Determination of Applicability where risk to other species and turbidity is expected to be low.

Monitoring needs: Critical to delineate target area and provide means for divers to stay on course with complete coverage.

Range of costs: \$5,000 to \$15,000 per acre, depending upon equipment features, contractor mobilization, hydrilla density, and total area to be harvested.

Other considerations: Turbidity may be unacceptable where a large area is suction harvested. Filtering system must be able to capture tubers and turions, or plant may be spread.

Benthic Barriers

Mode of action: Covers target area with a porous or non-porous blanket, limiting light and physically stressing plants.

Probability of successful control: Usually completely eliminates live vegetation from covered area in 30 to 60 days.

Potential non-target impacts: All plants under the barrier will be killed. Some invertebrates are also killed, but many relocate. Fish find the barriers attractive for cover and foraging area, mainly a function of "edge effect" (creation of edges between plants and open water).

Permitting needs: Often approved through a Negative Determination of Applicability (provisions of WPA do not apply) where hydrilla is the main plant affected. Otherwise permitted with an Order of Conditions with possible restrictions where other species are at significant risk.

Monitoring needs: Careful delineation of areas to be covered is needed. Condition of plant community, and the potential for recolonization through roots and tubers of hydrilla, should be assessed prior to removal.

Range of costs: Materials typically cost \$0.50 to \$1.00 per square foot. With application and maintenance costs, expect \$30,000 to \$50,000 per acre. However, material can be re-used indefinitely, so costs are greatly reduced for subsequent applications.

Other considerations: To enhance performance, benthic barriers should be carefully anchored and periodically cleaned. To minimize hooks and lures getting caught in benthic barriers, mark location with labeled buoys. Barriers may present a safety hazard in swimming areas.

Water Level Drawdown

Mode of action: Lowered water level exposes plants and substrate to drying and freezing action. Ice damage may also be a factor. Where plants can be dried, frozen, or ripped up by ice action, hydrilla may be greatly reduced in abundance. With years of repeated drawdown, exposed substrate tends to be dominated by coarse sediment less hospitable to hydrilla invasion.

Probability of successful control: Very high where drying, freezing and/or ice damage occurs. As this is a function of the weather pattern, uncertainty is high; about one out of three years provides effective drawdown conditions in Massachusetts. Where thick organic sediments, spring activity, or other factors limit freezing and drying, success will be lower. However, hydrilla can grow at significant depths, so the drawdown must be substantial to be effective.

Potential non-target impacts: Other plants that overwinter in vegetative forms are also likely to be harmed. Seed-producing plants may be stimulated. Some invertebrates (especially mollusks), amphibians (most likely frogs), reptiles (particularly wood turtles) and mammals (most probably beaver and muskrat) could be negatively affected. Effects on fish vary, depending upon timing and duration of drawdown and the interaction with feeding and reproduction. Direct water supply and water levels in wells may be affected.

Permitting needs: Requires an Order of Conditions under the Wetlands Protection Act, usually entailing a detailed review of the potential for non-target impacts.

Monitoring needs: Can be extensive. Pre- and post-implementation surveys are needed. Aside from effects on the plant community, effects on susceptible fauna may be required. Water supply must be monitored and a contingency plan is needed if supply is impaired. It should be assumed that at least three years of implementation will be needed to conduct a valid assessment of success and non-target impacts.

Range of costs: Where drawdown is facilitated by existing structures, costs are limited to permitting and monitoring, with potential for mitigation costs if impacts are unacceptable.

Other considerations: A very detailed evaluation of potential drawdown impacts is needed before attempting this technique. Issues of downstream flooding, refill time, and impacts on water supply and non-target organisms must be addressed.

Application of Fluridone

Mode of action: This systemic herbicide is absorbed by vegetative tissues and translocated throughout the plant, inhibiting the synthesis of carotenoid pigments. Lack of these auxiliary (protective) photosynthetic pigments causes susceptible plants to die slowly through reduced food production and damage by sunlight. Uptake must be nearly continuous over an extended period (>60 days preferred), necessitating extended exposure time.

Probability of successful control: Where adequate dose (>6 ppb, preferably up to 20 ppb) and exposure time (60-120 days) are maintained, hydrilla can be eradicated. This has proven difficult to achieve, however, particularly in partial lake treatments. Use of slow release pellet formulations or sequestration of the target area with impervious curtains maximizes exposure time and limits dilution of the dose. Follow up actions, such as hand harvesting, are often necessary. This treatment will not affect the seeds, tubers or turions of the hydrilla, so repeated applications may be needed to achieve eradication or control; it is typically assumed that treatment must be repeated for two or even three consecutive years.



Potential non-target impacts: Susceptibility of other plants to fluridone varies widely, and lowering of the dose can maintain much of the native community. However, doses <10 ppb have a lowered probability of controlling hydrilla, impacts to non-target plants should be expected in an aggressive hydrilla control program employing fluridone. Slow die-off of affected plants limits oxygen reduction. No impacts to fauna or humans are expected at applied doses.

Permitting needs: Requires an Order of Conditions under the Wetlands Protection Act and a License to Apply Chemicals from the DEP.

Monitoring needs: Normally the plant community is monitored before and after treatment. The concentration of fluridone is also commonly tracked on a weekly to monthly basis with an Enzyme Limited Immuno-Sorbent Assay (ELISA).

Range of costs: Costs range from \$500 to \$2,000 per acre, depending upon the form of fluridone applied, any necessary re-treatment to maintain dose, and any sequestration of the target area.

Other considerations: The combination of dose and exposure time is critical to success; the combination of achievable detention time and degree to which non-target plants must be protected will determine the potential for eradication or extended control.

Other Options

Other management options are not listed for one or more of the following reasons:

- impractical on a small scale
- not able to eradicate hydrilla
- could cause hydrilla to spread
- not approved for use in Massachusetts

Recommended Options for Early Eradication

The most commonly recommended early actions are hand harvesting and bottom barriers, each of which has a high potential for success, low cost on a localized basis, and limited permitting needs. Where growths are too dense for effective hand harvesting and too extensive for cost-effective bottom barrier placement, suction harvesting could be considered. However, as expanded growths indicate that tubers and turions have probably been deposited, treatment with fluridone is more commonly recommended, with repeat treatment in a second year and careful follow-up monitoring. Drawdown, where applicable, is a potentially effective preventive control in cases where repeated invasion is expected or documented, but is not applicable in all cases and is unlikely to prevent colonization in deeper waters.


A graphic summary of rapid response actions is provided in Figure 3. Most rapid responses will involve sparse growths over a limited area or small, dense beds in a confined area. While the listed techniques may still be applicable after growths have become widespread, addressing them may not qualify as a rapid response, and additional considerations (e.g., impacts to non-target organisms on a lakewide basis) are likely to become more important in the permitting process. The selection pathways shown in Figure 3 represent logical choices based on general features of the aquatic system, and are not intended to provide infallible rules or inflexible options. Practitioners should use a careful process of option review based on site specific data when selecting a rapid response.



Deciding Which Technique to Apply

The following decision tree is provided as an aid to evaluating control options. Thresholds for application are given as guidelines, not rigid rules. Individual circumstances may affect the choice of approach and outcome. Follow up monitoring is considered essential, and follow up control after an initial application is considered very likely to be necessary.

Figure 3. Decision Tree for the Control of Hydrilla (*Hydrilla verticillata*)



Notes: Hand harvesting and suction harvesting must include root system removal. Benthic barrier should remain in place for 30 to 60 days. Use of diquat or endothall is mainly to minimize spread of the plant; eradication is not expected. Fluridone use may include liquid, pellets, sequestration and repeat (boost or bump) treatments to maximize exposure, with treatments at >10 ppb potentially eradicating hydrilla but also damaging many native plant species. Drawdown use is dependent on many factors, including hydrology and use as a water supply. Moderate to dense growth over an extensive area (>10 acres) may not be appropriate for rapid response consideration.

Control of Established Infestations

This document deals mainly with early invasion and the new infestations that result, but it is important to note that older infestations, where the hydrilla has moved throughout the waterbody into all suitable habitats and probably become the dominant plant, can and should be addressed if continued invasion in the region is to be curtailed. The Practical Guide to Lake Management in Massachusetts (Wagner, 2004), a companion guide to the GEIR on Lake Management, provides a review of all available techniques for combating an invasive species infestation, and many of the techniques described are applicable to hydrilla. On a whole lake basis, herbicide (fluridone) treatment is the most cost-effective means for reducing hydrilla coverage and density to levels that can be controlled by physical techniques like hand harvesting or bottom barriers. Drawdown will reduce hydrilla in the drawdown zone, but it is rare that a waterbody can be drawn down enough to eliminate hydrilla without unacceptable impacts to non-target species. Techniques suitable for combating new growths are seldom practical or effective on a whole lake scale (e.g., hand harvesting, bottom barriers).

Maintenance techniques that limit the impact of hydrilla on waterbody uses, but do not typically result in elimination of hydrilla, include mechanical harvesting, hydroraking, rotovation, and the contact herbicides diquat and endothall. The physical methods may actually spread hydrilla if it is not already everywhere in the waterbody, in which case these are analogous to mowing a lawn. The contact herbicides do not kill the seeds, turions or tubers of hydrilla, allowing regrowth in future growing seasons.

Dredging can remove hydrilla along with all other plants and any remaining seeds or other propagules associated with the dredged sediment. The cost is extremely high, however, and resulting substrate conditions are usually still hospitable to hydrilla growth. With much bare area to be colonized, invasive species such as hydrilla are likely to become dominant if more desirable species are not actively introduced. Only if dredging results in a water depth too great for effective colonization by hydrilla is it likely to be the only method needed to control hydrilla in the target area.

The introduction of herbivorous insects, including hydrilla leaf mining flies or weevils has the potential to reduce hydrilla growths, but is unlikely to eliminate them as a function of classic predator-prey interactions, which usually result in cycles of abundance and scarcity. As fish eat the insects but few herbivores eat the hydrilla, constant inputs of herbivorous insects appear necessary to bolster populations to a level at which they can control hydrilla. In addition, none of these insects have been used to control hydrilla infestations within the United States. Grass carp can eliminate hydrilla (and indeed all other submersed plants) when stocked at sufficient density (Van Dyke et al. 1984), but are not approved for use in Massachusetts at this time.

Prevention of Re-Infestation

Once an invasion has been repulsed through any of the above methods, it should be apparent that the waterbody is susceptible to hydrilla. As the cost of prevention is much less than the cost of rehabilitation of an infested waterbody, steps should be taken to reduce the risk of re-introduction of hydrilla. As hydrilla most often comes from a local source, control activity is encouraged on a

watershed, multi-municipal or regional level. Working across political boundaries with limited funding is difficult, but represents the most sweeping opportunity to limit future invasions. Alternatively, and almost essential as a back-up, steps need to be taken at the individual waterbody to reduce the risk of re-introduction. Key steps may include:

- Education through the lake association or town for all users about the threat of hydrilla, how to avoid introducing it to the waterbody, how to identify it, and who to contact if it is found. See the other sections in this document for relevant information to be provided.
- Posting of all access points with signs warning of the threat, showing how to identify hydrilla, and urging that boats, fishing gear and other recreational equipment be cleaned before and after use in the waterbody. See the section on Communication and Education in this document.
- Provision of wash stations at boat ramps, and/or staffing of ramps with inspectors.
- Drawdown where applicable and permitted to minimize overwintering of introduced hydrilla.
- Monitoring of the plant community to detect hydrilla, with a focus on boat ramps and inlets.

Summary

1. Hydrilla (*Hydrilla verticillata*) is a highly invasive plant normally identified by long, slender, branching stems, and small pointed leaves arranged in whorls of 4-8. The leaf margins are saw-toothed, and the plant is noticeably rough when touched.
2. Hydrilla appears to be native to India and Korea, and possibly several other southeast Asian countries. Today it is found on every continent except Antarctica and South America. It is now found in approximately 20 of 50 states in the USA. It is most often transported on boats or trailers, by birds, and with water flow.
3. Hydrilla can be transported great distances by fragments that can root and grow. It becomes locally abundant by fragmentation and tubers, but also reproduces through seeds and turions.
4. Hydrilla creates canopies that shade out other plant species. At high density it provides poor habitat for most water-dependent fauna, impairs recreational uses, and can have negative impacts on water supply and flood control.
5. Hydrilla is most often detected in the early stages of infestation in water 2-20 ft deep by visual examination (viewing tube from boat or mask and snorkel). Look first in the vicinity of boat ramps, inlets, and areas of bird congregation. One effective long-term monitoring strategy involves setting up transects representing areas of the lake and searching at discrete depth intervals from shore to the maximum depth of plant growth.
6. When detected, map hydrilla coverage with notation of density as beds, scattered plants, or solitary stems. Be thorough with visual coverage of potentially infested areas. Record all other species present and their relative abundance. Confirm identification through the DCR.
7. Educate waterbody users by whatever means practical about the threat and presence of hydrilla. Posting of access points is useful in all cases. Signs should show how to identify hydrilla, urge that all boats, trailers and other recreational equipment be cleaned before and after use in the waterbody, and provide a contact name and phone number for reporting or correspondence.
8. It is advantageous to quarantine infested areas until removal can be attempted. Closing beaches and boat ramps can be problematic, legally and practically, but can promote greater awareness and support for prompt action. Use of curtains or screens both to keep people out of an infested area and to keep hydrilla inside is desirable but expensive.

9. Eradication of hydrilla detected early in an invasion can be accomplished with hand harvesting, suction harvesting, benthic barriers, drawdown, or the herbicides fluridone. Hand harvesting and benthic barriers are often allowable without an Order of Conditions under the WPA, and can therefore be implemented most rapidly. Each method has benefits and drawbacks, and the specific circumstances will affect which option(s) can be applied.
10. A range of additional options are available to combat later stage invasions. Those not mentioned as eradication options for new infestations have some feature that prevents effective, rapid use, but these techniques may have applicability under special circumstances.
11. Drawdown, where feasible, can act as a deterrent to invasion on an annual basis at a relatively low cost, through direct impact on invading hydrilla and by gradually altering the peripheral sediment features to make them less hospitable, but has many possible impacts on aquatic resources and requires a thorough evaluation in each case.
12. Once hydrilla has been removed after an invasion event, steps are necessary to prevent re-infestation. Education of waterbody users, with a focus on boating, and ongoing monitoring to detect new hydrilla plants are critical components. It should be assumed that hydrilla will return, but it is far easier to address new growths than to combat a full infestation.

References

- Bowes, G., A. S. Holaday, T. K. Van, and W. T. Haller. 1977. Photosynthetic and photorespiratory carbon metabolism in aquatic plants, pp. 289-298. *In* Hall, D. O., J. Coombs, and T. W. Goodwin(eds.). *Proceedings of the Fourth International Congress on Photosynthesis*. Reading, Pennsylvania. The Biochemical Society, London.
- Bratager, M., W. Cromwell, S. Enger, G. Montz, D. Perleberg, W. J. Rendall, L. Skinner, C. H. Welling, and D. Wright. 1996. Harmful Exotic Species of Aquatic Plants and Wild Animals in Minnesota. Annual Report. Minnesota Department of Natural Resources, St. Paul. 99 pp.
- CDFA (California Department of Food and Agriculture). 2000a. Hydrilla Program 20.20.01, Program Statement 2000 Season, Fiscal Year 2000/2001. Plant Health and Pest Prevention Services, integrated Pest Control Branch, California Department of Food and Agriculture, Sacramento, California, USA.
- Crow, G. E., and C. B. Hellquist. 2000. Aquatic and wetland plants of northeastern North America. Volume Two: Angiosperms: Monocotyledons. University of Wisconsin Press, Wisconsin.
- Engel, S. 1995. Eurasian watermilfoil as a fishery management tool. *Fisheries* 20(3):20-27.
- Godfrey, R. K. and J. W. Wooten. 1979. *Aquatic Wetland Plants of Southeastern United States: Monocotyledons*. The University of Georgia Press, Athens, Georgia, USA.



- Haller, W. T. 1978. *Hydrilla, a new and rapidly spreading aquatic weed problem*. Agricultural Experiment Station, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, Florida, USA.
- Haller, W. T. 1982. Hydrilla goes to Washington. *Aquatics* 4: 6-7.
- Haller, W. T. and D. L. Sutton. 1975. Community structure and competition between hydrilla and vallisneria. *Hyacinth Control Journal* 13:48-50.
- Harrel, S. L., E. D. Dibble, and K. J. Killgore. 2001. *Foraging behavior of fishes in aquatic plants*. APCRP Technical Notes Collection (ERDC TN-APCRP-MI-06). U.S. Army Engineer Research and Development Center, Vicksburg, Mississippi, USA.
- Holland, L. E., and M. L. Huston. 1984. Relationship of young-of-the-year northern pike to aquatic vegetation types in backwaters of the Upper Mississippi River. *North American Journal of Fisheries Management* 19:18-27.
- Johnstone, I.M., B.T. Coffey, and C. Howard-Williams. 1985. The role of recreational boat traffic in interlake dispersal of macrophytes: a New Zealand case study. *J. Environ. Manage.* 20:263-279.
- Killgore, K. J., E. D. Dibble, and J. J. Hoover. 1993. *Relationships between fish and aquatic plants: a plan of study*. Miscellaneous Paper A-93-1. U.S. Army Engineer Waterways Experiment Station, Vicksburg, Mississippi, USA.
- Langeland, K. A. 1996. *Hydrilla verticillata* (L.F.) Royle (Hydrocharitaceae), "The Perfect Aquatic Weed". *Castanea* 61:293-304.
- Lillie, R. A., and J. Budd. 1992. Habitat architecture of *Myriophyllum spicatum* L. as an index to habitat quality for fish and macroinvertebrates. *Journal of Freshwater Ecology* 7(2):113-125.
- Madeira, P. T., T. K. Van, K. K. Steward, and R. J. Schnell. 1997. Random amplified polymorphic DNA analysis of the phenetic relationships among world-wide accessions of *Hydrilla verticillata*. *Aquatic Botany* 59: 217-236.
- Madeira, P. T., C. C. Jacono, and T. K. Van. 2000. Monitoring hydrilla using two RAPD procedures and the nonindigenous aquatic species database. *Journal of Aquatic Plant Management* 28: 33-40.
- Mahler, M. J. 1979. Hydrilla the number one problem. *Aquatics* 1: 56.

- Pimental, D., L. Lach, R. Zuniga, and D. Morrison. 2000. Environmental and economic costs associated with nonindigenous species in the United States. *Bioscience* 50: 53-65.
- Rejmanek, M., and M.J. Pitcairn. 2002. *When is Eradication of Exotic Pest Plants a Realistic Goal?* Pages 249 - 253 in D. Veitch and M. Clout, eds. *Turning the Tide: Eradication of Invasive Species*. Invasive Species Specialist Group of the World Conservation Union
- Sainty, G. R. and S. W. L. Jacobs. 1981. *Waterplants of New South Wales*. New South Wales Water Resources Commission, New South Wales, Australia.
- Schmitz, D. C., B. V. Nelson, L. E. Nall, and J. D. Schardt. 1991. Exotic aquatic plants in Florida: a historical perspective and review of present aquatic plant regulation program, pp. 303-336. In Center, T. D., R. F. Doren, R. L. Hofstetter, R. L. Myers, and L. D. Whiteaker (eds.). *Proceedings of a Symposium on Exotic Pest Plants*. November 2-4, 1988, Miami, Florida. United States Department of the Interior, National Park Service, Washington, D.C.
- Schmitz, D. C. and D. Simberloff. 1997. Biological invasions: a growing threat. *Issues in Science and Technology* 8: 33-40.
- Smart, R. M. and J. W. Barko. 1988. *Effects of water chemistry on aquatic plants: Interrelationships among biomass production, plant nutrition and water chemistry*. Final Report A-88-5. U.S. Army Engineer Waterways Experiment Station, Vicksburg, Mississippi, USA.
- Steward, K. K., T. K. Van, C. Carter, and A. H. Pieterse. 1984. Hydrilla invades Washington, D. C., and the Potomac. *American Journal of Botany* 71: 162-163.
- Sutton, D. L., T. K. Van and K. M. Portier. 1992. Growth of dioecious and monoecious hydrilla from single tubers. *Journal of Aquatic Plant Management* 30:15-20.
- Tarver, D. P., J. A. Rodgers, M. J. Mahler, and R. L. Lazor. 1978. *Aquatic Wetland Plants of Florida*. Bureau of Aquatic Plant Research and Control, Florida Department of Natural Resources. Tallahassee, Florida, USA.
- Van Dyke, J. M., A. J. Leslie, Jr. and L. E. Nall. 1984. The effects of grass carp on the aquatic macrophytes
- Wagner, K.J. 2004. *The Practical Guide to Lake Management in Massachusetts*. Commonwealth of Massachusetts, Executive Office of Environmental Affairs, Boston, MA.
- Westman, W. E. 1990. Park management of exotic plant species: problems and issues. *Conservation Biology* 4: 251-260.