The Facts About Pediatric Asthma in Lynn



Pediatric asthma has serious consequences for the children and families it affects – and their communities. The City of Lynn has some of the highest rates of pediatric asthma and related health care use and costs in Massachusetts. Pediatric asthma also takes its toll on attendance at school and work. But there are steps we can take to reduce or remove asthma triggers and lessen the impact of asthma on the Commonwealth's children.

Quick Facts



During the 2011–2012 school year, 1,413 of Lynn's elementary and middle school students had asthma – 12.8% of students!



From 2008 to 2012, Lynn children accounted for more than 1,900 asthma-related hospitalizations and Emergency Department (ED) visits, with health care charges totaling **\$7.6 million**.^{2,3}



Lynn's black and Hispanic children had significantly higher rates of asthma-related ED visits than its white children.²

And white children in Lynn had higher rates of asthma-related hospitalizations and ED visits than their peers of the same race or ethnicity statewide.^{2,3}

PEDIATRIC ASTHMA PREVALENCE

In 2012, more Massachusetts children reported that they had ever been diagnosed with asthma (15.3%) and that they still had asthma (10.3%) than in the U.S. in general (13.7% and 8.9%, respectively). In fact, in 2012, 210,000 MA children had asthma. Lynn is home to a disproportionate number of these children. During the 2011–2012 school year, more than three out of every 25 Lynn children (or 12.8%) in Kindergarten through 8th grade had asthma, a significantly higher prevalence than for K–8 students statewide (11.9%).

SCHOOL AND WORK ABSENCES


Nationwide, asthma is the leading cause of school days missed due to a chronic illness with nearly 15 million school absences due to asthma annually.^{5,6} Between 2006 and 2010, over 40% of MA children aged 18 and younger with asthma missed school or day care due to their asthma in the past 12 months.⁷ Working families are especially challenged by pediatric asthma since a missed day of school for a child often means a missed day of work for a parent. Parents' loss of productivity from asthma-related school absences in the U.S. is estimated to be \$719 million annually.⁸

The Facts about Pediatric Asthma in Lynn

HEALTH CARE COSTS AND UTILIZATION

Asthma is the third-ranking cause of hospitalizations for children under 15 years of age in the US.⁶ Nearly 44% of all asthma hospitalizations in this country are for children.⁹ Lynn children have higher rates of hospitalizations and ED visits due to asthma than children statewide. From 2008 to 2012, Lynn children accounted for 1,914 asthma-related hospitalizations and ED visits, 2.6% of the Commonwealth's 72,442 hospitalizations and ED visits for pediatric asthma during that time. Lynn's health care charges associated with pediatric asthma were \$7.6 million. Medicaid was the expected payer for 64.6% of those charges.^{2,3}


In MA, asthma-related health care utilization is significantly higher for racial and ethnic minorities. (See Figures 1 and 2.) The same is true in Lynn. Between 2008 and 2012, among children in Lynn age 19 and under, the asthma-related ED rate for blacks and Hispanics was higher than for Lynn's white children (1.5 and 1.4 times higher, respectively). Black children also had a significantly higher asthma-related hospitalization rate than white children (1.4 times). And while Lynn's white children fare better than their black and Hispanic peers in the City, they had significantly higher rates of asthma-related ED visits and hospitalizations than white children statewide. And the children statewide.



Data Source: CY2008–2012 Massachusetts Emergency Department Discharge Database, Massachusetts Center for Health Information and Analysis (CHIA)

The Facts about Pediatric Asthma in Lynn





Data Source: CY2008–2012 Massachusetts Emergency Department Discharge Database, Massachusetts Center for Health Information and Analysis (CHIA)

ASTHMA TRIGGERS

Asthma prevalence is higher in low-income, urban neighborhoods^{10–12} due to greater exposure to air pollution and a range of indoor "triggers." Pets, rodents, cockroaches, mold, tobacco smoke, gas, and dust mites are well-known asthma triggers. Children's asthma may also be aggravated by exposure to things like air fresheners, cleaning products, pesticides, and perfumes. Reducing exposure to triggers in their homes, schools, and other locations in the community can have a major impact on the well-being of children with asthma. Visit www.mass.gov/dph/asthma for more information.

Working toward solutions in Lynn

The City of Lynn is a Prevention
Wellness Trust Fund partner working
with the Lynn Community Health
Center, Massachusetts Coalition for
the Homeless, and Lynn Housing
Authority and Neighborhood
Development to advance pediatric
asthma initiatives in Lynn.

FOR MORE INFORMATION

The Asthma Prevention and Control Program (APCP) at the Massachusetts Department of Public Health works to improve the quality of life for all Massachusetts residents with asthma, and to reduce disparities in asthma outcomes. APCP also works to reduce exposure to asthma triggers and irritants in homes, licensed child-care centers, schools, workplaces, and senior centers. We support the use of community health worker-led asthma home visits to improve asthma outcomes through the provision of resources and technical assistance around these interventions. The APCP provides Asthma Action Plans for children and adults in seven languages. For more information about the APCP and the work that we do, please call 617-624-5070, email us at prevention.wellness@state.ma.us or visit www.mass.gov/dph/asthma

3

The Facts about Pediatric Asthma in Lynn

SOURCES:

- Pediatric Asthma Surveillance, Bureau of Environmental Health, MA Department of Public Health
- ² CY2008–2012 MA Emergency Department Discharge Database, MA Center for Health Information and Analysis (CHIA).
- ³ CY2008–2012 MA Hospitalization Discharge Database, MA Center for Health Information and Analysis (CHIA).
- ⁴ CY2012 MA Behavioral Risk Factor Surveillance System, MA Department of Public Health
- Wirt, J., Choy, S. & Gerald D, Provasnik, S., Rooney, P., Watanabe, S., Tobin, R. & Glander, M. (2001). Condition of Education: 2001. Washington: National Center for Education Statistics, U.S. Department of Education.
- ⁶ CDC (2005). Asthma's Impact on Children and Adolescents. Atlanta: National Center for Environmental Health, Centers for Disease Control and Prevention. June 8, 2005.
- Source: CY2006–2010 Behavior Risk Factor Surveillance Survey Children Asthma Call-Back Survey.

- Wang, L.Y., Zhong, Y. & Wheeler, L. (2005). Direct and Indirect Costs of Asthma in School-Age Children. Preventing Chronic Disease, 2(1): [serial online], January.
- ⁹ National Hospital Discharge Survey . Hyattesville, MD: National Center for Health Statistics, 2000.
- Weiss, K.B. & Wagener, D.K. (1990). Asthma surveillance in the United States. A review of current trends and knowledge gaps. Chest 98(5 Suppl), 179S–184S.
- Matsui, E.C., Hansel, N.N., McCormack, M.C., Rusher, R., Breysse, P.N. & Diette, G.B. (2008). Asthma in the Inner City and the Indoor Environment. Journal of Immunology and Allergy Clinics North America 28, 665–686.
- O'Connor, G.T., Neas, L., Vaughn, B., Kattan, M., Mitchell, H., Crain, E.F., Evans, R., Gruchalla, R., Morgan, W., Stout, J., Adams, G.K. & Lippmann, M. (2008). Acute respiratory health effects of air pollution on children with asthma in US inner cities. Journal of Allergy and Clinical Immunology, 121(5) online.



For more information about the Asthma Prevention and Control Program, Massachusetts Department of Public Health:

Phone: 617-624-5070

Email: prevention.wellness@state.ma.us Website: www.mass.gov/dph/asthma