

Massachusetts Department of Public Health

POSTED: NOVEMBER 2019



This report contains both confirmed and estimated data through September 2019.

The chart above shows the month-by-month estimates for fatal opioid-related overdoses for all intents from January 2018 through September 2019. In the first nine months of 2019, there are 1,091 confirmed opioid-related overdose deaths and DPH estimates that there will be an additional 332 to 407 deaths.



Figure 2 shows the trend in annual number of confirmed and estimated cases of opioid-related overdose deaths for all intents from 2000 to 2018. In order to obtain timelier estimates of the total number of opioid-related overdose deaths in Massachusetts - confirmed and probable - DPH used predictive modeling techniques for all cases not yet finalized by the Office of the Chief Medical Examiner (OCME). Based on the data available as of October 4, 2019, DPH estimates that there will be an additional 68 to 72 deaths in 2017 and an additional 35 to 37 deaths in 2018, once these cases are finalized.

Opioid-Related Overdose Death Rates, All Intents

In 2018, DPH estimates a 1% decrease in the rate of opioid-related overdose deaths compared with 2017. This follows an estimated 3% decline in the rate of opioid-related overdose deaths from 2016 to 2017. The rate for 2018 represents an estimated 4% decrease from 2016.



Toxicology Analysis: Fentanyl and Other Drugs

Fentanyl is a synthetic opioid that has effects similar to heroin. It can be prescribed for severe pain. According to the U.S. Department of Justice, Drug Enforcement Administration's 2015 Investigative Reporting, while pharmaceutical fentanyl (from transdermal patches or lozenges) is diverted for abuse in the United States at small levels, much of the fentanyl in Massachusetts is due to illicitly-produced fentanyl, not diverted pharmaceutical fentanyl.

The standard toxicology screen ordered by the Office of the Chief Medical Examiner includes a test for the presence of fentanyl. Among the 903 opioid-related overdose deaths in 2019 where a toxicology screen was <u>also</u> available, 838 of them (93%) had a positive screen result for fentanyl. In the second quarter of 2019, heroin or likely heroin was present in approximately 24% of opioid-related overdose deaths that had a toxicology screen. Cocaine was present in approximately 42% of these deaths and benzodiazepines were present in approximately 33%. In the first quarter of 2014, amphetamines were present in 4% of opioid-related overdose deaths that had a toxicology screen. Since then, the presence of amphetamines has been increasing to approximately 7% of opioid-related overdose deaths in the second quarter of 2019. Since 2014, the rate of heroin or likely heroin present in opioid-related overdose deaths has been decreasing while the presence of fentanyl is still trending upward.

While screening tests can be used to note the rate at which certain drugs are detected in toxicology reports, they are insufficient to determine the final cause of death without additional information. The cause of death is a clinical judgment made within the Office of the Chief Medical Examiner.



* Beginning with the November 2019 report, DPH began to use a new method to identify substances present in the toxicology data, which can only be applied from 2017 onward; this new method cannot be applied to the older data

1. This is most likely illicitly produced and sold, not prescription fentanyl

2. Prescription opioids include: hydrocodone, hydromorphone, oxycodone, oxymorphone, and tramadol

Please note that previous estimates may change slightly as DPH routinely receives updated toxicology data from the Office of the Chief Medical Examiner and the Massachusetts State Police.

Technical Notes

- Opioids include heroin, illicitly manufactured fentanyl, opioid-based prescription painkillers, and other unspecified opioids.
- Data for 2017-2019 deaths are preliminary and subject to updates.
- Beginning with the May 2017 report, DPH started reporting opioid-related overdose deaths for all intents, which includes unintentional/undetermined and suicide.
- Beginning with the August 2019 report, DPH updated the case definition used to identify opioid-related overdose deaths to match the CDC's case definition. The following International Classification of Disease (ICD-10) codes for mortality were selected from the underlying cause of death field to identify poisonings/overdoses: X40-X44, X60-X64, X85, and Y10-Y14. All multiple cause of death fields were then used to identify an opioid-related overdose death: T40.0, T40.1, T40.2, T40.3, T40.4, and T40.6.
- This report tracks opioid-related overdoses due to difficulties in identifying heroin and prescription opioids separately. The Department regularly reviews projections as more information becomes available. Information from the Office of the Chief Medical Examiner and the Massachusetts State Police are now incorporated into the predictive model. This additional information has improved the accuracy of the model that predicts the likelihood that the cause of death for any person was an opioid-related overdose. DPH applied this model to death records for which no official cause of death was listed by the OCME. The model includes information from the death certificate, Medical Examiner's notes, and the determination by the State Police of a suspected heroin death. DPH added this estimate to the number of confirmed cases in order to compute the total number of opioid-related overdoses. Should new information become available that changes the estimates to any significant degree, updates will be posted.

Sources

- Massachusetts Registry of Vital Records and Statistics, MDPH
- Massachusetts Office of the Chief Medical Examiner

- Massachusetts State Police
- Population Estimates 2000-2010: National Center for Health Statistics. Postcensal estimates of the resident population of the United States, by year, county, age, bridged race, Hispanic origin, and sex (Vintage 2000-2010).
- Population Estimates 2011-2018: Small Area Population Estimates 2011-2020, version 2017, Massachusetts
 Department of Public Health, Bureau of Environmental Health. Population estimates used for years following the
 decennial census were developed by the University of Massachusetts Donahue Institute (UMDI) in partnership
 with the Massachusetts Department of Public Health, Bureau of Environmental Health.