You received the Johnson & Johnson (Janssen) vaccine Important Information for those receiving the COVID-19 vaccine

Thank you for receiving the COVID-19 vaccine. You are an important part of our effort to stop the spread and keep our communities safe.

You're not fully vaccinated yet!

You will not be fully vaccinated until two weeks after you receive the Johnson and Johnson vaccine. Until then, you can still contract and spread COVID-19. Continue to practice habits to reduce the spread.







Wear a mask



Keep your distance



Avoid groups

Not everyone will be vaccinated when you are. And while we know it dramatically reduces symptoms, we are still learning about the vaccine's protection against COVID-19 in real-life conditions.

Many people experience minor side effects, but major side effects are rare

Serious side effects from vaccines, including the COVID-19 vaccine, are rare. The most common side effects are minor and include:

- Pain at injection site
- Muscle pain and/or joint pain
- Headache

Tiredness

Chills

Fever

These side effects can last several days but are not cause for alarm. If pain at the injection site persists or you are worried, contact your doctor. If you do not have a primary care provider or wish to report any side effects to the CDC, visit cdc.gov/vsafe

Get medical care right away if you develop any of the symptoms below within three weeks after receiving the Johnson & Johnson COVID-19 Vaccine:

- Severe headache or blurred vision
 Chest pain
- Gut pain that does not go away

Shortness of breath

- Leg swelling
- Easy bruising or tiny blood spots under the skin

