THE ESSENTIAL SCHOOL HEALTH SERVICES PROGRAM DATA REPORT

2009 - 2010 School Year



Deval L. Patrick, Governor Timothy P. Murray, Lieutenant Governor JudyAnn Bigby, MD, Secretary of Health and Human Services John Auerbach, Commissioner of Public Health

> Massachusetts Department of Public Health Bureau of Community Health and Prevention Office of Statistics and Evaluation

> > Spring, 2011

This and other Massachusetts Department of Public Health publications can be accessed on the Internet at:

http://www.state.ma.us/dph/pubstats.htm

For additional copies of this report, please contact Robert Leibowitz at:

Massachusetts Department of Public Health Bureau of Community Health and Prevention Office of Statistics and Evaluation 250 Washington Street, 5th Floor Boston, MA 02108-4619

TDD/TTY: (617) 624-5992 (Division for Special Health Needs) or TDD/TTY: (617) 624-6001

To obtain other Department of Public Health data:

Register for the Department's free and internet-accessible data warehouse, MassCHIP: masschip.state.ma.us/beuser.htm or call 1-888-MAS-CHIP (MA only) or (617) 624-5541.

Acknowledgments

This report was prepared by Robert Leibowitz of the Office of Statistics and Evaluation, Bureau of Community Health and Prevention. Anne Sheetz, Director of School Health Services, wrote the introduction, providing the history of the Essential School Health Services model. The authors also acknowledge the contributions of Andy Epstein, Director, Division of Primary Care and Health Access and Special Assistant to the Commissioner. In addition, the authors would also like to thank all those who authored or contributed to earlier editions of this report. Further, we extend our appreciation to Anne Sheetz, Mary Ann Gapinski, Barbara Mackey, Paula Dobrow, Cathy Burgess, Thomas Comerford, and Janet Burke of the School Health Unit for their work with the Essential School Health Services program. At the school district level, we thank the nurse leaders and school nurses for providing their time, clinical expertise, and cooperation.

Table of Contents

| Introduction | 1 |
|---|----|
| Executive Summary | 5 |
| Findings | 8 |
| School Nurse Staffing | 8 |
| Student Demographics | 8 |
| School Health Services Activity | 9 |
| 1. Health Encounters | 10 |
| 2. Injury Reports, Early Dismissals and Referrals for Emergency Health Services | 11 |
| 3. Medication Management | 14 |
| 4. Health Screenings | 17 |
| 5a. Medical Procedures | 20 |
| 5b. Immunizations and H1N1 Flu Clinics | 24 |
| 6. Linkages to health care and insurance providers | 26 |
| 7. Oral Health | 28 |
| 8. Health Education, Tobacco Prevention, and Support Groups | 28 |
| 9. Nursing Case Management | 32 |
| Program Development | 33 |
| Students With Special Health Care Needs | 34 |
| 1. Types of Special Health Care Needs | 34 |
| 2. Students With Do Not Resuscitate (DNR) Orders | 36 |
| 3. Cardiovascular Health and Automated Electronic Defibrillators (AEDs) | 36 |
| Client Satisfaction | 37 |
| Actions to Promote Healthy Weight | 38 |
| References | 45 |
| APPENDIX A | 47 |
| School Districts and Student Enrollment | 47 |
| APPENDIX B | 50 |
| Scope of Service | 50 |

| APPENDIX C | 51 |
|-------------------------|----|
| Data Collection Methods | 51 |
| Data Analysis Methods | 52 |
| Data Limitations | 53 |

Introduction

In recent years, research has highlighted major societal, legal, and medical technological changes and their effect on the demand for school health services. These changes include: (1) increased awareness of the relationship between health and educational achievement; (2) improved medical technology; (3) increase in the number of students with special health care needs combined with an increase in condition severity in these students; (4) rapid restructuring of the health care delivery system; (5) laws requiring inclusion; (6) changes in family structure and patterns of parental employment; (7) rise in social morbidities such as substance abuse, depression, and violence among children; and (8) impact of diverse cultural and linguistic groups.

- Attendance in the early grades is correlated with school achievement and dropout rates. School nurses support attendance by providing needed health services in school. They also provide assessments of illness and injuries. School nurses are significantly less likely to dismiss a student than an unlicensed counterpart (Pennington & Delaney, 2008), and in one study 57% less likely (Wyman, 2005).
- As neonatal intensive care unit survivors enter early intervention services and kindergarten, the need for school health services increases (Clement, Barfield, Ayadi & Wilber, 2007). Data show that the students in the Commonwealth's schools require increasingly complex health care during the school day. The current (FY10) Essential School Health Data Report indicates that 28% of the students in ESHS and partner schools have at least one special health care need. Children with special health care needs (CSHCN) are defined by the Department of Health and Human Services, Health Resources and Services Administration, Maternal and Child Health Bureau (MCHB) as: "...those who have or are at increased risk for a chronic physical, developmental, behavioral, or emotional condition and who also require health and related services of a type or amount beyond that required by children generally" (McPherson et al., 1998).
- Nationally, the incidence of diabetes among adults 18 79 has almost doubled in the last 10 years (CDC, 2008), and diabetes is increasingly being diagnosed in children and adolescents (Hannon, Rao, and Arslanian, 2005). In Massachusetts the percentage of children prescribed epinephrine for life threatening anaphylaxis more than doubled between 2001 and 2010, rising from .72% to 2.05%. In addition, the Cedar Rapids v. Garret Supreme Court decision of 1999 clarified the extent to which school districts are required to provide school nursing services for medically fragile children.
- Children assisted with medical technology, e.g. catheterizations, tracheostomies, ventilators, etc., are now attending school. Likewise terminally ill children are in the Commonwealth's classrooms, necessitating end of life planning.
- The rapid restructuring of the health care delivery system has dramatically impacted school health service programs. With reduced hospitalizations and/or reduced lengths of stay, school nurses are now often responsible for supervising the care of children who have illnesses such

as acute asthma and diabetes, formerly managed in a hospital setting (Chabra et al., 2000; Coffman et al., 2008; Leslie et al., 1998; Schutte et al., 1997).

- Social attitudes that promote inclusion, as well as state and national laws, such as the Individuals with Disabilities Act and Section 504 of the Rehabilitation Act of 1973 specify disability rights and access to education, resulting in more children requiring nursing care and other health-related services in school (Palfrey et al., 1992; Raymond, 2009; Small et al., 1995).
- With more working parents, children who are sick with mild or chronic conditions are less likely to be monitored at home on school days, and more likely to be sent to the school nurse for assessment and a determination as to whether they need to see a physician (Smolensky and Gootman, 2003; Thurber et al., 1991; Uphold & Graham, 1993; U.S. Census Bureau, 2000; Wold, 2001). In Massachusetts, 36.7% of health encounters in 2009-2010 were for the purpose of illness assessment.
- Students spend a large part of their day at school; therefore, the school has become an important site where health and education risks, e.g. depression, absenteeism, substance use, may be identified and timely interventions initiated. One in five young people between that ages of 9 and 17 experiences symptoms of mental health problems, and one in ten children and adolescents has a mental illness severe enough to cause some level of impairment; yet in any given year, only about one-fifth of children in need of mental health services actually receive them. (US Surgeon General's Conference on Children's Mental Health, 2000). This disproportion can result in increased demands for professional health services in the schools (Thurber et al., 1991).
- Massachusetts schools have many "newcomer" groups, both immigrants and refugees, as
 well as those families who move between different communities. Often such families rely on
 the school for information about what services or providers are available in the community.
 They may not know how to obtain care elsewhere because of language or cultural barriers
 and, therefore, look to the school health service for assistance.

The Massachusetts Department of Public Health (MDPH) recognizes the need for quality school health services and provides consultation to all of the Commonwealth's school districts. Since 1993, the Department of Public Health has extended to a number of school systems the opportunity to expand on the basic school health services model by establishing the Essential School Health Services Program (ESHS). (The Essential School Health Services Program was originally entitled the Enhanced School Health Service Program.)

In 1993, thirty-six school districts were funded for three and half years to: (a) strengthen the infrastructure of school health services in the areas of personnel and policy development, programming, and interdisciplinary collaboration; (b) incorporate health education programs, including tobacco prevention and cessation programs, into the existing school health programs; and (c) develop linkages between school health service programs and community health care providers.

In October 1997, the Department funded 19 school districts under the Essential model (Essential School Health Services, ESHS) and 8 school districts with experience in developing the Essential model to provide consultation to approximately 42 additional school districts ("recipient schools") across the Commonwealth (Essential School Health Services with Consultation, ESHSC). These recipient school districts were interested in developing similar school health service programs.

In November, 1999, the Massachusetts legislature allocated additional funding to the Essential School Health Service Programs (ESHS and ESHSC). School systems for both models were selected for participation through a competitive bid process based on a Request for Response (RFR) developed by MDPH. As a result of the 1999 RFR process, a total of 77 school districts (or affiliated school systems)¹ received awards in 2000: 11 Essential School Health Services with Consultation and 66 basic Essential Programs. An added component of the 1999 RFR was that each applicant public school district was required to provide some elements of basic school health services (vision/hearing screening, immunization review, etc.) to all non-public and charter schools within the community (77 award recipients in 2000 served 253 non-public and charter schools)². An additional 32 school districts received awards in 2001; all of these were basic Essential Programs (Sheetz, 2003).

In February 2003, midyear budget reductions eliminated most funding for the ESHS programs for the remainder of the fiscal year. Because of this, three programs decided to withdraw from the ESHS grant, thus reducing the number of participants to 106 school districts in the spring of 2003. Three more schools withdrew from the grant in 2004, and one additional school withdrew in 2006, leaving 102 districts in the ESHS program.

In 2009 a new funding cycle started and 80 school districts were funded (see **Appendix A**). Of these 80 funded districts, 68 (85%) had been funded during the previous cycle. Thirty-four districts in the previous funding cycle (33% of the 102 districts included in the earlier funding cycle) were not included in the new funding cycle. The number of funded districts was reduced because some funds were freed to establish an extension of the ESHS programs, namely mentored/partnered schools. Each of the 68 experienced programs (with the exception of the large cities) was required to mentor or partner with two other school districts in order to increase adoption of the standards established in the ESHS program initiative. Therefore 146 additional mentored/partnered school districts, each with a limited amount of funding, were added to the model. These school districts were required to meet a specified scope of service. Of note is that in the FY10 school year, these mentored/partnered school districts will begin to submit some data, consistent with ESHS requirements.

¹ ESHS funding was awarded to local public school systems, regional academic school systems, independent vocational systems, vocational-technical regional systems, and school unions.

² 223 non-public (private and parochial) schools, 30 charter schools.

³ All public school districts were invited to join this program. A number of vocational schools, educational collaboratives and charter schools were also invited to participate in this program when an opening in a geographic area was available.

In addition to the Mentor/Partner School Program component of the 2009 grant cycle, a Regional Consultation program was also included in the funding. These six regional ESHS programs (based on the EOHHS defined regions) were selected to provide consultation to ESHS programs within their general geographical area. Regional consultation school districts must have been previously awarded the Essential School Health Service (ESHS) or Essential School Health Service with Consultation programs (ESHSC). The general goal of the ESHS Regional Consultation grant is to maximize the existing school nursing expertise, leadership and infrastructure to provide additional consultation to ESHS programs (including their mentored/partnered school districts and community public schools as appropriate) within a general region.

In October 2009, 9C cuts to the ESHS programs resulted in the reduction to 50% funding for 13 programs. These reductions impacted data collection efforts in these school districts. Therefore, the FY10 report has fewer districts reporting on certain indicators.

Throughout this report, comparison data from previous years are presented. Because the mix of school districts included in the program has changed over the years, caution should be exercised when interpreting these data, as differences may be the result of the changing composition of school districts in the program.

The staff of the School Health Unit, Division of Primary Care and Health Access in the MDPH Bureau of Community Health and Prevention administers the programs.

Executive Summary

The information collected by the Essential School Health Services Program provides a valuable snapshot of school nursing practice in a diverse cohort of Massachusetts public schools. The data reveal that school nurses perform a wide array of duties -- direct care, health education, administrative case management, and policy/program development and oversight -- on behalf of students whose health needs range from routine to serious and complex. In addition, some school nurses provide services to school staff.

Analysis of the ESHS program data for the school year beginning September, 2009 and ending June, 2010 showed the following:

- 933 schools in 78 ESHS school districts reported a total of 4,946,757 student health encounters, and 99,903 staff encounters.
- In a typical district, students visited the school nurse an average of 1.1 times per month.⁴ There was substantial variability among school districts, with the encounter rate ranging from 0.7 to 1.9 visits per month.
- After assessment and/or treatment by a school nurse, the majority (92.2%) of the students visiting the nurse's office with an illness or injury complaint were returned to the classroom to continue their studies.
- 9.5% of the more serious injuries to students were classified as intentional. These include injuries resulting from assaults (e.g. physical fighting) and those that were self-inflicted (e.g. intentional drug overdose, suicide attempts).
- School nurses in ESHS districts referred students to urgent health care services a total of 7,086 times, 1,805 of which involved 9-1-1 ambulance calls. In the remaining cases, parents or others were called to transport the student to health services.
- The majority (92.2%) of the prescriptions managed by the school nurse were for medications dispensed on a PRN, or "as needed" basis.⁵
 - Among students taking PRN medications, asthma medications were the most common (34.5 prescriptions per 1,000 enrolled students).
 - The prescription rate for "as needed" epinephrine increased from 7.2 per 1,000 students in 2003 to 20.5 per 1,000 in 2010.
 - Among students on scheduled prescription medications, psychotropic medications (drugs affecting perception, emotion or behavior) were by far the most common (5.3 per 1,000 enrolled students).
- In the ESHS districts, school nurses administered an average of 119,106 doses of prescription medication to students per month. Fifty-seven percent of the scheduled doses were for psychotropic medication, and 56% of the PRN prescription doses were for asthma medication.

⁵ PRN is an abbreviation for "pro re nada," a Latin term meaning "as needed." PRN medications are not scheduled for set times, but given as needed, based on a nursing assessment.

⁴ "Typical" is defined in this report as the median district. It is the district lying in the middle of the group, with half the districts having higher values and half having lower values.

- School nurses in 133 ESHS and partner districts conducted Body Mass Index screenings on 163,509 students in grades 1, 4, 7 and 10. Overall, 33.4% of the students screened were overweight or obese (16.3% obese, 17.1% overweight).
- Diabetes care procedures account for an increasing amount of nurses' time. Insulin pump care increased from 5.2 procedures per 1,000 students the prior year to 7.0 procedures per 1,000 students this year. Blood glucose testing, the most common medical procedure, increased from 56.7 procedures per 1,000 students each month the prior year to 66.0 procedures per 1,000 students. While the proportion of students requiring glucose testing may be relatively small, the number of daily tests on those students requires considerable nursing time and assessment, as each child usually requires glucose monitoring several times a day.
- In response to the H1N1 flu pandemic, school nurses conducted flu clinics in cities and towns throughout the state. Over 10 times more immunizations were administered in schools in 2009-2010 than in the prior school year. Immunizations were administered to a total of 51,408 students and 12,880 staff.
- 13,138 students received an oral health screening from a school nurse, and 25,155 were screened by a dentist or hygienist.
- Tobacco prevention and cessation programs reached substantial numbers of individuals, although activity levels varied widely across districts.
 - Individual tobacco cessation counseling sessions increased from 1,675 students the prior year to 8,833 students this year (43 districts).
 - 13,387 students participated in group tobacco prevention activities.
- A total of 176,034 students with special health care needs were reported to school nurses in ESHS and partner districts (28% of enrollment).
 - The most common physical/developmental condition reported to school nurses was asthma (124.5 per 1,000 enrolled students).
 - The most commonly reported behavioral/emotional condition was Attention-Deficit/Hyperactivity Disorder (57.0 per 1,000 enrolled students).
- Almost 78% of the ESHS school districts have at least one AED in all of their school buildings, up from 29.7% in 2003-2004 (Table 20) and 68.4 % last year. All ESHS districts have deployed AEDs in at least one school building. Only 17.8% of school buildings in ESHS districts do not have an AED
- Parent satisfaction with school health services was measured through a survey mailed to a sample of parents with a child who received nursing services. The response rate was 38.8% (1,009 questionnaires were returned out of 2,600 distributed). Satisfaction rates on the 6 measured criteria ranged from 93 to 97 percent.

Continued refinements in data collection and analysis will more accurately capture school nursing and school health activity, improve our ability to monitor the health needs and status of the school age population, and identify areas for improvements in services and quality of care.

Identifying trends in school health encounters and student health indicators may assist school nursing staff in improving the delivery of prevention, education, and intervention services to the school community. Future data collection efforts will seek to increase our knowledge of health needs in the school setting and in the school age population, explore the relationship between student health status and educational outcomes, and investigate ways in which health services and prevention activities in schools can help children live healthier lives.

Findings

School Nurse Staffing

In the ESHS program, 1,157 full-time school nurses (or full time equivalents) provided health care services to students and staff in the 80 ESHS funded public school districts. The student-to-nurse ratio was 412 students per nurse (compared to 404 the prior year). An additional 321 school nurses provided care in 57 partner school districts. In the partner districts, the student-to-nurse ratio was 466. Finally, 26 additional school nurses provided care in 2 partner charter schools and 6 partner educational collaboratives. Nearly 30 percent of ESHS RN school nurses have an advanced degree (Table 1).

| TABLE 1. Educational Level of RN School Nurses in ESHS and Partner Districts | | | | | | | |
|--|----------|----------------|---------------|-----------------------|-----------|-----------|--|
| | | (Percent of to | otal RN FTEs, | , <i>2009-2010</i>)) | | | |
| | Total RN | Diploma | Associate | Bachelor's | Advanced | | |
| Type of | FTEs | RN | Degree | Degree | Degree | Other | |
| District | (Number) | (Percent) | (Percent) | (Percent) | (Percent) | (Percent) | |
| ESHS | 996.7 | 5.6 | 4.3 | 58.3 | 29.6 | 2.2 | |
| Partner | 290.1 | 8.3 | 4.5 | 64.2 | 22.0 | 1.0 | |

Includes Nurse Leaders. "Advanced Degree" includes Master's and Doctoral degrees. Source: 77 ESHS and 58 partner districts.

Student Demographics

In 2009-2010, 49.9 percent of Massachusetts public school students were enrolled in an ESHS-funded school district. The racial and ethnic composition of the ESHS student population is different than that found in the Massachusetts public school population, however. There is a higher percentage of Black and Hispanic students in ESHS-funded districts (Table 2). In addition, a higher percentage of students in ESHS-funded districts are low income, have limited English proficiency, and have a first language that is not English (Table 3).

| TABLE 2. Race/Ethnicity of Students in ESHS Districts and Massachusetts Public Schools (2009-2010) | | | | | | |
|--|---------|---------|--|--|--|--|
| ESHS Schools State Public Schools | | | | | | |
| Race/Ethnicity | Percent | Percent | | | | |
| African American or Black | 12.1 | 8.2 | | | | |
| Asian | 6.8 | 5.3 | | | | |
| Hispanic or Latino | 22.2 | 14.8 | | | | |
| Multi-race, Non-Hispanic | 2.4 | 2.2 | | | | |
| Native American | 0.3 | 0.3 | | | | |
| Native Hawaiian or Pacific Islander | 0.1 | 0.1 | | | | |
| White | 56.0 | 69.1 | | | | |

⁶ These statistics include data from the ESHS districts, but do not include data from any associated districts. The count of "School Nurses" includes only Registered Nurses (RNs) and nurse leaders, but excludes other health support staff which may have been funded by the ESHS contract.

⁷ Enrollment numbers for educational collaboratives are not available.

Source: Massachusetts Department of Elementary and Secondary Education.

| TABLE 3. Selected Characteristics of Students in ESHS Districts and Massachusetts Public Schools (2009-2010) | | | | | | | |
|--|-----------------------------------|------|---------|------|--|--|--|
| | ESHS Schools State Public Schools | | | | | | |
| Characteristic Number Percent Number Per | | | | | | | |
| First Language Not English | 111,238 | 23.3 | 149,300 | 15.6 | | | |
| Limited English Proficient | 48,350 | 10.1 | 59,337 | 6.2 | | | |
| Low Income | 212,923 | 44.6 | 314,870 | 32.9 | | | |
| Total Population | 477,163 | | 957,053 | | | | |

Source: Massachusetts Department of Elementary and Secondary Education.

Of the 304,049 students in 64 ESHS funded districts whose health insurance status was reported, 64% had private insurance, 35% had public insurance, and 1% had no insurance (Table 4). The status of 16% of students in ESHS funded districts and 8% of partner districts was unknown.

| TABLE 4. Health Insurance Status of Students in ESHS and Partner Districts | | | | | | | |
|--|-----------|-----------------------------|-----------|-----------|--|--|--|
| (2009-2010) | | | | | | | |
| Type of Insurance | | | | | | | |
| | Number of | Private Public No Insurance | | | | | |
| District Type | Students | (Percent) | (Percent) | (Percent) | | | |
| ESHS funded | 304,049 | 64.1 | 34.6 | 1.2 | | | |
| Partner | 88,682 | 81.6 | 17.6 | 0.7 | | | |

Source: Status Reports submitted by 64 ESHS and 38 partner districts. Districts reporting insurance status for less than 30% of their student enrollment were excluded. Percentages may not add up to 100 due to rounding error.

School Health Services Activity

The primary goal of the Essential School Health Services Program is to improve the delivery of health services to students by reinforcing the school health service infrastructure. Toward that end, program participants were required to report throughout the year the type and scope of school nursing activity in their districts. These activities were divided into nine categories of data:

- 1) Health encounters, including dispositions following assessment
- 2) Injury reports, early dismissals, and referrals for emergency health services
- 3) Medication management
- 4) Screenings
- 5) Medical procedures
- 6) Linkages to health care and insurance providers
- 7) Oral health
- 8) Health education, tobacco prevention, and support groups
- 9) Nursing case management

1. Health Encounters

Each month, districts reported the total number of student health encounters. An "encounter" was defined as any contact with a student during which the school nurse provided counseling, treatment, or aid of any kind. Casual conversations fall outside this definition and were not counted. In addition, mandatory screenings (such as vision, hearing, BMI and postural) were not counted as encounters because these are routine population-based activities. Screenings were tracked separately, however.

During FY2006, the ESHS Evaluation Committee refined the monthly and annual data collection tools. As a result, the FY07, FY08, and FY09 encounter categories are not comparable to those used in previous years. In addition to changes in encounter categories, districts no longer report secondary reasons for an encounter. The major impact of that change is that the multifaceted nature of the health encounter, which often includes health education and mental health counseling components, is not fully reflected in these data: The following rules are used to help define encounter categories:

- Every encounter includes nursing assessment and health education. An encounter is recorded as an Individual Health Education encounter only when the primary issue is health education and there is no illness or injury involved. Individual Health Education encounters previously made up a large percentage of the reported secondary issues.
- An illness encounter may include illness assessment, acute illness, chronic health condition, etc. It excludes scheduled medication administrations (e.g. daily medication administration for ADHD) and scheduled procedures (ostomy care, scheduled glucose testing).
- Mental/Behavioral Health Support includes any encounter requiring active listening, anticipatory guidance, stress management, behavior modification/program support or evaluation of altered mental status. The primary reason for the encounter is related to a mental/behavioral health need. Mental/behavioral health services tend be underreported as nurses will often categorize an encounter according to the presenting complaint (e.g., headache) even if it is determined that the complaint has an underlying mental/behavioral health origin.

Between September 1, 2009 and June 30, 2010, 78 ESHS school districts reported a combined total of 4,946,757 student health encounters. In a typical district, 84 percent of the student enrollment visited the health room at least once during the school year. "Illness assessment," "Injury/first aid," and "Scheduled medication administration" were the most common reasons for visits to the school nurse (Table 5). The number of encounters reported per district varied widely, with individual districts averaging between 240.4 and 36,896.1 encounters per month. These differences were largely due to district size. In a typical district, each student visited the school nurse an average of 1.1 times per month, although the encounter rate varied across the districts

⁸ While the goal of recording secondary reasons for an encounter was to capture the mental health services being provided, this goal was not achieved. Nurses frequently categorize the encounter with the presenting symptom, e.g., headache, when, upon further assessment, the underlying cause relates to behavioral health. An exploratory study by the Massachusetts School Nurse Research Network is underway to address this issue.

⁹ 77 districts reporting.

from 0.7 to 1.9 visits per month. While some students are seen several times each month, many others are never seen. The school nurse workload, measured by the number of encounters a full time nurse logs each month, varied greatly across the districts, with the school nurse workload in a typical district being 455.1 student encounters per month¹⁰.

Health services were also provided to school staff (i.e., teachers and administrators). School nurses in 78 ESHS districts reported a total of 99,903 staff health encounters. Across all districts, monthly averages ranged from 1.5 to 1,172 staff health encounters per month.

| TABLE 5. Number of Student and Staff Health Encounters | | | | | | | |
|--|-----------|---------|--------|---------|--|--|--|
| September 1, 2009 - June 30, 2010 | | | | | | | |
| Students Staff | | | | | | | |
| Type of Encounter | Number | Percent | Number | Percent | | | |
| Illness Assessment | 1,815,581 | 36.7 | 35,295 | 35.3 | | | |
| Injury/First Aid | 1,018,622 | 20.6 | 14,529 | 14.5 | | | |
| Scheduled Medication Administration | 720,809 | 14.6 | 1,048 | 1.0 | | | |
| Scheduled Medical Procedures* | 590,416 | 11.9 | 10,595 | 10.6 | | | |
| Individual Health Education | 139,196 | 2.8 | 11,871 | 11.9 | | | |
| Mental/Behavioral Health Support | 65,906 | 1.3 | 3,828 | 3.8 | | | |
| Other | 596,227 | 12.1 | 22,737 | 22.8 | | | |
| TOTAL | 4,946,757 | 100.0 | 99,903 | 100.0 | | | |

^{*&}quot;Scheduled Medical Procedures" are those performed for preexisting conditions, which usually require an MD order. Source: *Monthly Activities Reports* submitted by 78 districts in the Essential School Health Services program.

An additional 1,589,821 student encounters and 2,057 staff encounters were reported by 50 partner districts (including charter school districts), 5 partner collaboratives, and 6 private schools. The typical nurse workload in partner districts was 482.8 student encounters per month, slightly higher than the workload in funded districts.

2. Injury Reports, Early Dismissals and Referrals for Emergency Health Services

An important function of school nursing practice is to provide on-site health services to students who are sick, injured, or experiencing a serious health emergency. Each month, districts tallied the number of on-campus injury reports, early dismissals due to illness, and referrals for emergency health services. After assessment and/or treatment by a school nurse, the majority (92.5%) of students visiting the nurse's office with an illness or injury complaint returned to the classroom to continue their studies (Table 6 and Figure 1). These on-site services provide major benefits. Students who are treated on-site can be returned to the classroom with minimal interruption of their educational activities; working parents do not have to take time off from work to provide care; and the high cost of treatment in a doctor's office is avoided.

-

¹⁰ For these calculations, "school nurses" includes only RNs. The "typical" district workload was the workload that fell in the middle of the group (Half the ESHS districts had a higher workload, and half a lower workload).


| TABLE 6. Disposition After Illness/Injury Assessment September 1, 2009- June 30, 2010 | | | | | | | |
|--|----------------|------|--------|------|--|--|--|
| _ | Students Staff | | | | | | |
| Disposition Number Percent Number Percent | | | | | | | |
| Returned to Class | 3,968,866 | 92.5 | 65,352 | 92.2 | | | |
| Dismissals | 265,516 | 6.2 | 2,899 | 18.5 | | | |
| Other* | 55,207 | 1.3 | 2,603 | 3.7 | | | |
| Total | 4,289,589 | | 70,854 | | | | |

^{*} Includes "Stayed in health office" and "Referred to counselor's office". Source: *Monthly Activities Reports* submitted by districts in the Essential School Health Services program.

When students had to be dismissed, it was usually the result of illness (93.2%) rather than injury (6.8%).

The returned-to-class rate for student health encounters reported by 50 partner districts (which have a higher student-to-nurse ratio than funded districts) was 89.0%, which was lower than that reported by funded districts, and the dismissal rate was 8.6%, higher than that reported by funded districts.

FIGURE 1. Disposition After Nursing Assessment Student Health Encounters September 1, 2009- June 30, 2010



^{*} Includes "Stayed in health office" and "Referred to counselor's office". Source: *Monthly Activities Reports* submitted by districts in the Essential School Health Services program.

For injuries of a more serious nature, school nurses filed *injury reports* according to state and local policy. For the 2009-2010 School Year, districts reported a total of 23,381 student injury reports and 2,194 staff injury reports (Table 7):

| TABLE 7. Number of Student and Staff Injury Reports | | | | | | | | |
|---|---------------|---------|--------|---------|--|--|--|--|
| September 1, 2009 - June 30, 2010 | | | | | | | | |
| | Student Staff | | | | | | | |
| Intent | Number | Percent | Number | Percent | | | | |
| Unintentional | 18,214 | 77.9 | 1549 | 70.6 | | | | |
| Intentional | 2,223 | 9.5 | 244 | 11.1 | | | | |
| Unknown intent | 2,944 | 12.6 | 401 | 18.3 | | | | |
| Total | 23,381 | | 2,194 | | | | | |

Source: Monthly Activities Reports submitted by districts in the Essential School Health Services program.

Of the student injury reports filed by school nurses, 9.5% involved the intentional infliction of injury (Table 7). These include injuries resulting from assaults (e.g. physical fighting) and those that were self-inflicted (e.g. intentional drug overdose, suicide attempts).

In addition, school nurses in the 78 districts referred students to *urgent health care services* a total of 7,086 times.

- In 1,805 (25.5%) of these events, 9-1-1 or ambulance services were called.
- In the remaining 5,281 (74.5%) events, parents or others were called to transport the student to health services.

3. Medication Management

In 1993, the Massachusetts Department of Public Health promulgated regulations governing the administration of medications in public and private schools. The purpose of these regulations (105 CMR 210.000) is to provide minimum safety standards for the administration of prescription medications to students during the school day.

The school nurse's role in managing the medication administration program for the district is broad in scope. In addition to developing district-wide medication policies in collaboration with the school committee, school administration, and school physician, the school nurse:

- administers medications to students (including monitoring students' response to medications);
- delegates the administration of selected medications to appropriately trained school staff (if the district is registered with the MDPH to do so);
- ensures the proper training and supervision of these designated staff; and
- establishes a formal record-keeping system for the district's medication administration program.

Implicit in the description of medication administration is the nurse's responsibility for the following: development of the medication administration plan; assessment of the child prior to administering each medication; follow-up evaluation of medication efficacy and side effects; and ongoing communication with parents and providers.

ESHS districts tracked the number of *prescriptions* that had been ordered for their students. Throughout the year, the total number of prescriptions reported to school nurses averaged 83,182.4 per month for the 78 districts (Table 8). Note that because some students had more than one prescription, the number of prescriptions is larger than the number of students with prescriptions. Among prescriptions taken on a scheduled basis, psychotropic medications were the most common, while among prescriptions taken on an "as-needed" (PRN) basis, analgesics and asthma medications were the most common.

TABLE 8. Number of Student Prescriptions Reported to School Nurses (Monthly Average)

September 1, 2009 - June 30, 2010

| | Medication Schedule | | | | | |
|-----------------------------|------------------------------|---------------------------------|---------------------------------------|--|--|--|
| Medication Class | Scheduled (All Districts) | PRN (As needed) (All Districts) | Total (Daily & PRN) Medications | | | |
| Analgesics | 21.4 | 30,103.2 | 30,124.6 | | | |
| Antibiotics | 289.2 | 976.4 | 1,265.6 | | | |
| Anticonvulsants | 158.2 | 621.7 | 779.9 | | | |
| Antihypertensive | 63.8 | 33.6 | 97.4 | | | |
| Antihistamines | 41.2 | 6,671.0 | 6,712.2 | | | |
| Asthma Medications | 444.8 | 17,329.5 | 17,774.3 | | | |
| Epinephrine | 0.0 | 8,817.3 | 8,817.3 | | | |
| Insulin | 991.4 | 756.6 | 1,748.0 | | | |
| Psychotropic | 3,269.7 | 513.7 | 3,783.4 | | | |
| Other Prescription/OTC Meds | 1,140.3 | 10,939.4 | 12,079.7 | | | |
| Total | 6,420.0 | 76,762.4 | 83,182.4 | | | |
| Row Percent | 7.7% | 92.3% | 100.0% | | | |

Source: Monthly Activities Reports submitted by 78 districts in the Essential School Health Services program.

Tables 8a and 8b show the at-school prescription rates reported by the ESHS districts. The atschool prescription rate reflects the medications that are to be administered at school, during school hours, by the school nurse (or under the supervision of the school nurse). These rates understate the actual number of students taking prescription medications, however. There are two reasons for this. First, students who self-administer at school without the knowledge of the nurse are not counted in the nurse's data reports. 11 This type of "counting error" may disproportionately lower reported prescription rates for certain categories of students. Middle and high school students, for example, might be more likely to self-administer than elementary school students, and, therefore, would be less likely to be counted in the numbers reported by the school nurse. Second, medications taken only at home, as some types of daily medications are, are unlikely to be reported to school nurses. For example, the decrease in the at-school psychotropic prescription rate over the last few years (from 21.0 per 1,000 students in 2001 to 5.1 per 1,000 students in 2009) may be due to the use of new one-dose slow-release psychostimulant drugs, which are administered at home and are not reported to school nurses. On the other hand, PRN medications (medications prescribed for administration on an 'as needed' basis) such as medications taken to treat asthma attacks or allergic reactions, are more likely to be reported to the school nurse because of the potential need for administration during the school day. As a result, prescription rates for these medications may be better estimates of the true overall prescription rate for the school age population.

¹¹ Regulations require that students inform nurses about self-administered medications. If students do not comply with regulations, these medications may not come to the attention of school nurses.

| TABLE 9a. Prescription Medication Rate for Scheduled Medication | | | | | | | | |
|---|--------------|-------------|-------------|---------|-------------|--------|--|--|
| (Prescriptions Per 1,000 Students) | | | | | | | | |
| School | | Asthma | | | Anti- | | | |
| Year | Psychotropic | Medications | Antibiotics | Insulin | Convulsants | Others | | |
| 2000-2001 | 21.0 | 1.5 | 1.4 | 0.2 | | 1.9 | | |
| 2001-2002 | 13.2 | 1.0 | 1.2 | 0.3 | | 2.0 | | |
| 2002-2003* | 7.0 | 0.5 | 0.8 | 0.3 | 0.2 | 0.9 | | |
| 2003-2004 | 7.3 | 0.9 | 0.8 | 0.6 | 0.5 | 1.3 | | |
| 2004-2005 | 5.6 | 0.4 | 0.8 | 0.6 | 0.3 | 1.1 | | |
| 2005-2006 | 5.8 | 0.3 | 0.7 | 0.8 | 0.3 | 1.2 | | |
| 2006-2007 | 5.5 | 0.6 | 0.8 | 1.0 | 0.3 | 1.4 | | |
| 2007-2008 | 5.0 | 0.5 | 0.8 | 1.3 | 0.2 | 1.5 | | |
| 2008-2009 | 5.1 | 0.6 | 0.6 | 1.5 | 0.2 | 1.6 | | |
| 2009-2010 | 5.3 | 0.6 | 0.6 | 1.7 | 0.2 | 1.6 | | |

While the scheduled medication rate for insulin increased (from 0.2 per 1,000 students in 2001 to 1.7 in 2010), rates for most other classes of scheduled medications decreased from 2000-2001 levels, including psychotropic medications, asthma medications, and antibiotics (Table 9a). In contrast, for "as needed" medications, rates for a number of medication classes have increased. For example, the epinephrine prescription rate increased from 7.2 per 1,000 students in 2001 to 20.5 per 1,000 in 2010 (Table 9b). Similarly, "as needed" prescription rates increased for insulin and anti-convulsants.

| T | ABLE 9b. | Prescript | ion Med | lication | Rate for | As Needed | (PRN) Me | dication | | |
|------------|------------------------------------|-----------|---------|-----------------|----------|-----------|------------------|----------|--------|--|
| | (Prescriptions Per 1,000 Students) | | | | | | | | | |
| School | Asthma Medi- | Epi- | Anal- | Anti- hista- | | Psycho- | Anti- Convul- | Anti- | | |
| Year | cations | nephrine | gesic | mines | Insulin | tropic | sants | biotics | Others | |
| 2000-2001 | 25.2 | 7.2 | | | 0.5 | 0.5 | | 0.1 | 10.1 | |
| 2001-2002 | 26.3 | 8.3 | | | 0.7 | 0.4 | | 0.1 | 9.3 | |
| 2002-2003* | 22.7 | 8.1 | 4.5 | | 1.0 | 0.2 | 0.1 | 0.1 | 12.6 | |
| 2003-2004 | 30.2 | 9.8 | 15.6 | | 1.2 | 1.4 | 0.4 | 0.2 | 3.7 | |
| 2004-2005 | 28.0 | 12.1 | 4.2 | | 1.3 | 1.2 | 0.3 | 0.1 | 3.5 | |
| 2005-2006 | 30.9 | 12.8 | 4.4 | | 1.4 | 1.1 | 0.4 | 0.1 | 3.3 | |
| 2006-2007 | 32.2 | 15.3 | 5.7 | 4.8 | 1.5 | 0.8 | 0.7 | 0.0 | 6.4 | |
| 2007-2008 | 33.4 | 16.9 | 6.7 | 5.7 | 1.6 | 1.1 | 0.7 | 0.0 | 6.4 | |
| 2008-2009 | 35.3 | 18.8 | 6.2 | 8.1 | 1.5 | 1.0 | 1.1 | 0.0 | 6.3 | |
| 2009-2010 | 34.5 | 20.5 | 6.2 | 9.5 | 1.6 | 1.0 | 1.2 | 0.0 | 5.6 | |

^{*} The 2002-2003 school year report only included data for 4 of the 10 months of the school year. The 2000-2001 school year had 74 districts reporting as compared to 103 districts in 2003-2004, and 80 districts in 2008-2009.

Rates shown are those reported by the typical (median) district in the ESHS program.

Source: Monthly Activities Reports submitted by districts in the Essential School Health Services program

School nurses in the 78 ESHS districts administered an average of 119,106 doses of medication to students per month. Psychotropic medication was the most commonly administered type of scheduled prescription medication, and asthma medication was the most commonly administered type of PRN prescription medication. Among medications administered per school protocol, analgesic medication was the most common. (Table 10).¹²

| TABLE 10. Average Number of Medication Doses by Type |
|--|
| Administered to Students by School Nurses* Per Month |
| September 1, 2009- June 30, 2010 |

| | Medication Schedule | | | | | |
|-------------------------|---------------------|---|----------|-----------------------------|----------|-------|
| Medication Class | Schedule | Scheduled Doses PRN Doses per Prescription | | PRN Doses per Protocol** | | |
| | N | % | N | % | N | % |
| Analgesic | 74.5 | 0.1 | 2,077.6 | 12.0 | 18,036.1 | 59.2 |
| Antibiotic | 1,088.6 | 1.5 | 50.5 | 0.3 | 638.0 | 2.1 |
| Anticonvulsant | 1,730.8 | 2.4 | 33.7 | 0.2 | 0.5 | 0.0 |
| Antihypertensive | 819.1 | 1.1 | 3.3 | 0.0 | 7.1 | 0.0 |
| Antihistamine | 203.4 | 0.3 | 318.6 | 1.8 | 562.9 | 1.8 |
| Asthma | 2,489.6 | 3.5 | 9,617.6 | 55.7 | 315.8 | 1.0 |
| Epinephrine | 0.0 | 0.0 | 50.1 | 0.3 | 4.3 | 0.0 |
| Insulin | 13,301.6 | 18.6 | 2,896.5 | 16.8 | 66.1 | 0.2 |
| Psychotropic | 40,413.4 | 56.6 | 340.1 | 2.0 | 62.0 | 0.2 |
| Other | 11,243.8 | 15.8 | 1,863.8 | 10.8 | 10,798.1 | 35.4 |
| TOTAL | 71,364.8 | 100.0 | 17,251.8 | 100.0 | 30,490.9 | 100.0 |

^{*} Includes supervised self-administration ** These are protocols for non-prescription medications written by school physicians. Source: *Monthly Activities Reports* submitted by districts in the Essential School Health Services program.

School nurses also administered an average of 3,166 doses of medication to school staff per month, including 2,977 monthly doses of OTC/PRN medications, and 190 monthly doses of other prescription medications.

4. Health Screenings

Public schools in Massachusetts are required by law to conduct postural, hearing, vision, and height/weight screening on all students. Some school systems conduct additional health

¹² "PRN doses administered per protocol" refers to medication orders, signed by the school physician, which permit school nurses to administer over-the-counter (non-prescription) medications to students, according to guidelines provided by the Board of Registration in Nursing. "PRN doses per prescription" refers to medication orders written for prescription medications, which are to be administered to specific students.

¹³ The law permits waivers of these screening requirements in certain circumstances. Postural screenings of students in grades 5 through 9 may not be waived, however. Beginning in FY11, all public schools will be required to complete BMI screenings for students in grades 1, 4, 7, and 10. See 105 CMR 200 for further changes in screening requirements.

screenings based on the particular health needs of their students. School nurses are responsible for screening students and making referrals for follow-up care when needed. Parents are responsible for making appointments for the follow up care specified in the referral, and for ensuring that students keep the appointments. During the school year, school nurses at 77 districts conducted the following number of required and voluntary student health screenings (Table 11). These numbers represent *initial* screenings, and do not include *re-screenings*.

| TABLE 11. Yearly Student Health Screenings and Referrals School Year 2009-2010 | | | | | | | | |
|---|--|----------------------|--------|------------------------|--------|---------------------------|--|--|
| | Screenings Referrals Completed Referra | | | | | | | |
| Type of Screening | Number | % of All Students | Number | % of Screened Students | Number | % of Referred Students | | |
| Hearing | 222,811 | 47.2 | 4,555 | 2.0 | 1,979 | 43.4 | | |
| Height/Weight | 186,719 | 39.5 | 23,238 | 12.4 | 2,857 | 12.3 | | |
| Postural | 136,743 | 28.9 | 4,237 | 3.1 | 1,802 | 42.5 | | |
| Vision | 264,831 | 56.0 | 28,254 | 10.7 | 11,006 | 39.0 | | |

Source: Status Reports submitted by 77 districts in the Essential School Health Services program.

Body Mass Index (BMI) Screenings

The Centers for Disease Control and Prevention recommends the use of Body Mass Index (BMI) measurement to screen for obesity in children. BMI is a number calculated from height and weight, and is considered a reliable indicator of body fat in most people. For children and teens, BMI is age and sex specific. The measure is plotted on BMI growth charts to reveal the child's percentile ranking, which indicates the relative position of the child's BMI among children of the same age and sex. The BMI percentile can then be used as a screen for overweight or underweight. BMI percentiles derived from direct measurements should be more accurate than those derived from self-reports in student surveys. Nurses were asked to complete BMI screenings for all students in grades 1, 4, 7 and 10. In order to ensure the results would be representative of the district, if the district did not screen at least 60% of enrollment for a grade, data for that grade were excluded. School nurses in 133 districts (77 ESHS and 56 partner districts) met this criteria for 1 or more grade levels, for a total of 163,509 students (Table 12). Nurses in 104 districts met the screening criteria for all 4 grade levels.

| TABLE 12. Number of School Districts Providing Universal BMI Screening September 1, 2009- June 30, 20010 (n = 133 districts) | | | | | | | |
|--|-----|-----------|----------|----------|--|--|--|
| Grade | | Districts | Students | Screened | | | |
| | n | % | n | % | | | |
| 1 | 122 | 91.7 | 43,761 | 95.0 | | | |
| 4 | 124 | 93.2 | 43,828 | 95.0 | | | |
| 7 | 122 | 91.7 | 41,633 | 92.2 | | | |
| 10 | 117 | 88.0 | 34,287 | 86.3 | | | |
| All grades | 133 | 100.0 | 163,509 | 89.0 | | | |

Notes: Includes 77 ESHS districts and 56 partner districts. A total of 104 local districts, 23 regional academic districts, 3 regional vocational districts, and 2 charter districts. If district data for a grade did not include at least 60% of the grade enrollment it was excluded. *Percent of enrollment in districts included.

^{*} A "completed" referral is one in which an appointment for follow-up care has been made and kept.

Overall, 33.4% of the students screened were overweight or obese (16.3% obese, 17.1% overweight). In each of the 4 grade levels, at least 28% of the students screened were overweight or obese, with males in all 4 grades more likely to be overweight or obese than females (Table 13). BMI screening results may be sent to a student's physician or parents, depending on local policy.

TABLE 13. Percentage of Under- and Overweight Students in Grades 1, 4, 7, and 10 in ESHS and Partner Districts as Reported by School Nurses Conducting Universal BMI Screenings
(133 Massachusetts Public School Districts, 2009-2010 School Year)


| | Grade 1 | | de 1 | Grade 4 | | Grade 7 | | Grade 10 | |
|----------------------|--------------------------|--------|--------|----------|----------|----------|----------|----------|----------|
| | | Male | Female | Male | Female | Male | Female | Male | Female |
| | Total students screened: | 22,439 | 21,322 | 22,504 | 21,324 | 21,180 | 20,453 | 17,134 | 17,153 |
| | BMI Percentile | | | | | | | | |
| Weight category* | Range | % | % | % | % | % | % | % | % |
| | Less than the 5th | | | | | | | | |
| Underweight | percentile | 2.3 | 2.4 | 1.8 | 2.2 | 2.5 | 2.3 | 2.1 | 1.6 |
| | 5th percentile to less | | | | | | | | |
| Healthy Weight | than the 85th | 67.0 | 68.7 | 60.6 | 63.8 | 60.4 | 63.8 | 64.2 | 67.3 |
| | 85th to less than the | | | | | | | | |
| Overweight | 95th percentile | 15.8 | 15.2 | 18.1 | 17.9 | 17.5 | 17.9 | 16.5 | 18.1 |
| | Equal to or greater | | | | | | | | |
| Obese | than the 95th | 14.9 | 13.7 | 19.4 | 16.1 | 19.5 | 16.0 | 17.3 | 13.0 |
| Total | | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 |
| | | | | | | | | | |
| Subtotal: Overweight | t or Obese | 30.7 | 28.9 | 37.6 | 34.0 | 37.1 | 33.9 | 33.8 | 31.1 |

^{*} For children and adolescents, the CDC uses the term "overweight" instead of "obese" and the term "at risk of overweight" instead of "overweight." We have chosen to use the same labels that are used with adults to avoid confusion over the terminology in line with recommendations recently released by a committee of experts representing 15 medical and health organizations (Expert Committee, 2007).

5a. Medical Procedures

School enrollment of children assisted by medical technology has increased in recent years. This phenomenon presents multiple challenges for school administrators, parents and guardians, school health services personnel, teachers, and students. ESHS school districts collected information on the number and type of procedures that involved medical technology, as well as other medical procedures performed by school nurses. Consistent trends in the school health data may be associated with emergent public health issues. For example, the increase in Blood Glucose Testing and Insulin Pump Care over the past 5 years may be a consequence increasing diabetes prevalence in face of the current obesity/diabetes epidemic. Monthly medical procedure rates per 1,000 enrolled students are shown in Figures 2 and 3.


FIGURE 2. Medical Procedure Rates (Students) Sepember 1, 2009 - June 30, 2010



Source: *Monthly Activities Reports* submitted by districts in the Essential School Health Services program. Note: Rates were calculated from those districts performing the procedure at least once.

The procedures listed in Figure 2 required differing amounts of nursing time. Those procedures identified with an asterisk (*) require significant amounts of professional nursing care, health education and monitoring. Many of these procedures were formerly performed in a hospital setting.

FIGURE 3. Procedure Rates per 1,000 Students per Month* School Years 2000-2001 through 2009-2010



^{*}Among those districts performing the procedure at least once.

Note that in 2002-2003, data was available for only 4 out of 10 months. If there are no data points then data was not available for that year. Rates shown are those reported by the typical (median) district in the ESHS program.

Source: Monthly Activities Reports submitted by districts in the Essential School Health Services program

While some procedure rates have declined (blood pressure monitoring, wound care), procedures related to diabetes management (blood glucose monitoring and insulin pump are) have increased.

^{**} The definition of Wound Care was changed in 2007, so that dressing changes are no longer counted.

Monthly medical procedure totals are summarized in Table 14:

| TABLE 14 Medical Procedure Types and Totals | | | | | | | |
|---|--------------------------------|-------|--|--|--|--|--|
| September 1, 2009- June 30, 2010 | | | | | | | |
| | Number of Procedures Per Month | | | | | | |
| Type of Procedure | Students | Staff | | | | | |
| Administer Immunizations | 5,141 | 1,288 | | | | | |
| Auscultate Lungs | 14,216 | 261 | | | | | |
| Blood Glucose Testing | 30,013 | 81 | | | | | |
| Blood Pressure Monitoring | 2,805 | 1,735 | | | | | |
| Carbohydrate/Insulin Calculation | 11,655 | 4 | | | | | |
| Catheter Care | 2,307 | 3 | | | | | |
| Central Line Care (a) | 89 | 1 | | | | | |
| Check Ketones | 1,408 | 2 | | | | | |
| Device Adjustment (e) | 1,571 | 9 | | | | | |
| Insulin Pump Care | 11,047 | 185 | | | | | |
| IV Infusion Care | 4,474 | 3 | | | | | |
| Nebulizer Treatment | 35 | 3 | | | | | |
| Ostomy Care (c) | 1,079 | 6 | | | | | |
| Oxygen Administration | 408 | 2 | | | | | |
| Oxygen Saturation Check | 190 | 3 | | | | | |
| Peak Flow Monitoring | 3,993 | 100 | | | | | |
| Physical Therapy | 1,279 | 26 | | | | | |
| Suctioning | 786 | 5 | | | | | |
| Tracheostomy Care | 182 | 0 | | | | | |
| Tube Care or Usage (b) | 88 | 1 | | | | | |
| Weight measurement (d) | 3,484 | 1 | | | | | |
| Wound Care | 458 | 187 | | | | | |

a) Central Line Care: Monitor infusion or administration, Pump monitoring, IV Bag Change, dressing change.

Source: Monthly Activities Reports submitted by districts in the Essential School Health Services program.

b) Naso-Gastric, Gastronomy or Other Feeding Tube Care or Usage

c) Ostomy Care- Colostomy/Ileostomy/Urostomy

d) Weight management for medical conditions not related to screening

e) Includes orthotic or prosthetic device adjustment, wheelchair assistance, and crutch walking instructions.

In addition to medical procedures, school nurses performed head checks for pediculosis at a rate of 15.2 per 1,000 students per month.

5b. Immunizations and H1N1 Flu Clinics

In response to the H1N1 flu pandemic, school nurses conducted flu clinics in cities and towns throughout the state (See map, following page). Over 10 times more immunizations were administered in ESHS-funded schools in 2009-2010 than in the prior school year. In ESHS and partner schools, immunizations were administered to a total of 67,128 students and 15,224 staff. This number understates the total number of immunizations provided, as immunizations administered by school nurses in town flu clinics are not captured in the school totals. The majority of immunizations were administered in the fall, particularly in December (Figure 4a). Dismissals due to illness declined after the flu clinics were held (Figure 4b).






FIGURE 4a. Student and Staff Immunizations by Month





Source: Monthly Activities Reports submitted by 77 ESHS districts and 56 partner districts.



6. Linkages to health care and insurance providers


ESHS school systems identified students without a primary care provider and, in consultation with their families, referred them to appropriate health care services. A referral is reported whenever an actual appointment has been set up with a provider or agency. School systems also referred many students to their existing primary care providers. During the 2009-2010 school year, participating districts reported the following:

- A total of 101,856 students requiring primary care services were identified and referred to primary care providers. Those students without primary care providers were referred to new providers. Referrals included:
 - 6,424 referrals to new primary care providers (6.3% of total primary care referrals). In a typical district, monthly referrals to new primary care providers averaged 1.5 students, a rate of 0.4 referrals per 1,000 enrolled students per month.
 - 95,432 referrals to existing primary care providers (93.7% of total primary care referrals). In a typical district, monthly referrals to existing primary care providers averaged 60.6 students, a rate of 15.2 referrals per 1,000 enrolled students per month.

-

¹⁴ Prior to 2006-2007, a referral was counted whenever the student was advised to follow-up with a provider.

FIGURE 4. Primary Care Provider Referrals Median Monthly Rate Per 1,000 Students School Years 2003-2004 to 2009-2010



Source: Monthly Activities Reports submitted by districts in the Essential School Health Services program.

In addition, districts in the ESHS program provided the following referrals for students during 2008-2009:

- 5,604 referrals to insurance providers.
- 12,054 referrals for mental/behavioral health services.

Each month, school nurses receive Massachusetts Asthma Action Plans (MAAPs) from health care providers. These written plans provide individualized instructions for managing asthma episodes and administering asthma medications. During the school year, 77 ESHS districts reported receiving MAAPs for 5,904 students. Individual districts received between 0 and 1,515 action plans.

¹⁵ This section refers only to Standard Triplicate Form Massachusetts Asthma Action Plans.

7. Oral Health

School nurses are increasingly performing oral health related activities. Table 13 summarizes these activities for the 2009-2010 school year.

The typical district participating in oral health screening activities screened students at an annual rate of 45.4 per 1,000 students. There was considerable variability across districts, with the range being 0.5 to 443 screenings per 1,000 students. Slightly more than one-third of oral health screenings were performed by school nurses (Table 15).

| TABLE 15. Number of Students Receiving Oral Health Services September 1, 2009 - June 30, 2010 | | | | | | |
|--|------------------------------------|--|--|--|--|--|
| Type of Oral Health Activity | % of Districts Performing Activity | Number of Students (All Districts) | | | | |
| Oral health screenings by a school nurse | 39.0 | 13,138 | | | | |
| Oral health screenings by a dentist or hygienist | 55.8 | 25,155 | | | | |
| Referrals to a dental provider | 63.6 | 8,291 | | | | |
| Referrals completed | 48.1 | 3,078 | | | | |
| Screenings of third grade students | 50.6 | 5,572 | | | | |
| Dental sealants applied in school | 37.7 | 12,691 | | | | |
| Flouride rinse treatments applied in school | 54.5 | 26,778 | | | | |

Source: Monthly Activities Reports submitted by 77 districts in the Essential School Health Services program.

8. Health Education, Tobacco Prevention, and Support Groups

School nurses are often called upon to provide health education and deliver presentations. In this teaching role they provide information to students, staff, and community members on topics such as nutrition education, life threatening allergies, and human growth and development. Throughout the 2009-2010 school year, school nurses in 78 ESHS districts reported making 13,752 classroom presentations. In addition, 2,055 presentations were made in 50 partner districts, 55 presentations were made in 5 collaboratives, and 181 presentations were made in 6 private schools. In a typical ESHS funded district, each full-time school nurse delivered 1.1 presentation every month (range: 0 to 8.3 presentations per nurse per month). The types of presentations given most frequently in funded districts were fitness/nutrition/wellness, life threatening allergies, and oral health/hygiene (Table 16). During the school year, school nurses in funded districts made an average of 12.2 presentations per nurse, while the average in partner districts was 9.8 presentations per nurse, the average in collaboratives was 4.6 presentations per nurse, and the average in private schools was 2.9 presentations per nurse.

Rate is based on those districts that performed one or more oral health screening activities.

| TABLE 16. Number of Wellness/Safety Presentations |
|---|
| and Number of Participants, by Topic Area |
| September 1, 2009- June 30, 2010 |

| | | Number of Participants Per Month | | | | |
|----------------------------|---|----------------------------------|---------|-----------|--|--|
| Topic Area | Number of Presentations Per Month | Students | Staff | Community | | |
| Blood Borne Pathogens | 59.1 | 415.6 | 1,682.5 | 90.6 | | |
| CPR/AED Programs | 42.1 | 104.3 | 399.1 | 20.1 | | |
| Crisis Team | 23.8 | 157.0 | 245.4 | 2.4 | | |
| Environmental Health | 44.4 | 1,993.8 | 371.0 | 144.9 | | |
| Fitness/Nutrition/Wellness | 248.8 | 7,674.3 | 1,414.5 | 287.1 | | |
| Growth/Development | 99.8 | 2,308.0 | 84.3 | 130.0 | | |
| Life Threatening Allergies | 188.8 | 808.1 | 2,521.3 | 60.7 | | |
| Mental Health/Wellness | 62.3 | 1,076.9 | 194.2 | 41.8 | | |
| Oral Health/Hygiene | 301.0 | 8,291.8 | 360.9 | 80.3 | | |
| Other | 305.1 | 12,104.5 | 1,930.0 | 1,054.5 | | |

Source: Monthly Activities Reports submitted by 78 districts in the Essential School Health Services program.

Health education was also promoted through the preparation of flyers and mailings. During the school year, school nurses in funded districts were involved in the creation of a total of 17,291 health promotion / education flyers or mailings. In the typical funded district, each nurse was involved in the creation of 1.1 flyer or mailing per year.

During the school year, school nurses in ESHS districts provided the following tobacco prevention/cessation and substance abuse services:

- 64 districts reported a total of 1,841 assessments of students for suspected substance abuse.
- A total of 465 tobacco group prevention meetings were held in 21 districts, in which attendance summed to 13,387 students and 538 adults.
- A total of 135 tobacco group cessation meetings were held in 19 districts, in which attendance summed to 494 students and 98 adults.
- Individual tobacco cessation counseling sessions increased from 1,675 students the prior year to 8,833 students this year (43 districts). Counseling sessions were also provided to 149 adults.¹⁷

-

¹⁷ Trainings of School Nurse Interventions to Assist Students to Stop Smoking resumed in FY10. Each ESHS district is required to have at least one high school nurse trained and implementing the program.

• In 24 districts, students were referred to other tobacco prevention/cessation services 243 times, and adults were referred to outside sources 46 times.

During the 2002-2003 school year, the MDPH School Health Unit collaborated with the University of Massachusetts, Department of Preventive and Behavioral Medicine, in conducting a randomized controlled trial (RCT) to determine if school-nurse intervention could help individual students stop using tobacco. The intervention consisted of a series of scheduled appointments with content designed to address tobacco triggers, barriers to quitting, and helpful techniques. The student was required to designate a quit date. The study was implemented in 71 Massachusetts schools. The results demonstrated the feasibility and potential efficacy of this intervention in increasing self-reported short term (6 week and 3 month) quit rates among adolescent smokers who wished to quit.

Based on these outcomes, the National Institutes of Health (NIH) awarded the University of Massachusetts Medical School (UMMS) a four-year grant to test this intervention in a randomized controlled trial, designed to be delivered by the school nurse in the course of her/his routine clinical duties through four individual 15 to 20 minute sessions with individual teens. As a result of the partnership with the UMMS Department of Preventive and Behavioral Medicine and the MDPH School Health Unit, thirty-six public high schools with an enrollment of at least 350 students are currently participating in this NIH grant study. Prior to the NIH study, the Northeastern School Health Institute had been offering trainings to school nurses based on the results of the 2002-2003 study. These trainings have been resumed in FY10.

.

¹⁸ It is anticipated that approximately 1,000 teens will be recruited during the course of two years with baseline assessments including salivary cotinine (metabolic of nicotine) and follow-up assessments 3 and 12 months following baseline. Cotinine validation and 12 month follow-up assessment is considered the gold standard of tobacco research.

¹⁹ The Northeastern School Health Institute is the continuing education vendor tor the MDPH School Health Unit, providing relevant programs for approximately 2,000 school nurses a year.

Support Groups

Table 17 summarizes participation in student support group activities led or assisted by school nurses. It does not include tobacco-related support groups which were discussed previously. Across all topic areas, a total of 352 support group meetings were conducted every month.

| TABLE 17. Participation in Support Group Activities, by Topic Area September 1, 2009- June 30, 2010 (n=78 districts) | | | | | | | | |
|--|---------------------|-------------------|----------------------|-------|----------------------|--|--|--|
| | % of ESHS Districts | Monthly | Monthly Participants | | | | | |
| Topic Area | Offering Group | Group Meetings | Students | Staff | Parent/ Community | | | |
| Alcohol or Substance Abuse | 28.2 | 22.3 | 120.9 | 17.1 | 21.3 | | | |
| Anger/Conflict/Violence Management | 24.4 | 22.0 | 189.5 | 32.0 | 5.7 | | | |
| Asthma | 24.4 | 7.3 | 27.8 | 16.5 | 6.8 | | | |
| Diabetes | 25.6 | 25.6 | 36.2 | 26.7 | 10.8 | | | |
| Emotional / Psychosocial Support | 41.0 | 86.6 | 171.4 | 74.4 | 10.0 | | | |
| Food Allergy | 29.5 | 18.9 | 29.3 | 69.8 | 11.9 | | | |
| Gay/Bisexual/Lesbian/ Transgender | 9.0 | 7.2 | 44.0 | 8.0 | 0.8 | | | |
| Health Careers | 24.4 | 9.3 | 135.8 | 8.3 | 9.8 | | | |
| Nutrition/Physical Activity | 50.0 | 65.7 | 320.9 | 132.4 | 20.6 | | | |
| Peer Leadership | 21.8 | 13.8 | 202.6 | 23.6 | 11.9 | | | |
| Other | 62.8 | 73.6 | 590.4 | 171.2 | 54.6 | | | |
| Total* | | 352.3 | 1,868.8 | 580.0 | 164.2 | | | |

Source: Monthly Activities Reports submitted by districts in the Essential School Health Services program.

The type of support group most likely to be offered was "Nutrition/Physical Activity." This type of group was offered by 50% of districts and attracted the highest number of participants, among both students and staff. The second most common type of support group was "Emotional/psychosocial," offered by 41% of districts. Support groups in the "Emotional/psychosocial" area met more frequently than the other types of support groups.

^{*} Those participants that are in more than 1 group may be counted twice.

During the school year, nurses in funded districts provided an average of 8.3 meetings per 1,000 students, while nurses in partner districts provided an average of 7.9 meetings per 1,000 students.

In nutrition programs, school nurse support can extend beyond making support groups available. Some students come to school without adequate breakfasts or lunches, and school nurses provide food and/or snacks. During the school year, school nurses reported they provided snacks a total of 104,589 times.

9. Nursing Case Management

Data from the monthly activities report revealed that, beyond providing direct care to students, school nurses spent a significant portion of their day performing case management duties that included communication with families, other school staff, and community health care providers about student health concerns. During the school year, school nurses from 78 districts conducted:

- a total of 913,325 health counseling and education communications with parents (including phone calls and letters, but excluding meetings and home visits), with the typical district reporting 814.1 communications per month (range: 10.6 to 8,040.4 communications per month);
- a total of 965 home visits, with the typical district reporting 0.2 home visits per month (range: 0.0 to 16.4 home visits per month);
- a total of 325,120 communications with other school staff about student health issues, with the typical district reporting 260.0 communications per month (range: 4.7 to 4,107.4 meetings per month);
- a total of 79,726 communications with other agencies and health providers about student health issues, with the typical district reporting 29.9 communications per month (range: 0.0 to 1,870. phone calls per month).
- a total of 26,046 case management meetings, with the typical district reporting 14.6 meetings per month (range: 0.0 to 484.8 meetings per month).

The following table shows median case-management activity levels per school nurse FTE per month across the 78 participating districts:

TABLE 18. Nursing Case Management Activities: Student-Health Related Activities Per Month Per Nurse FTE September 1, 2009 - June 30, 2010

| | Activities Per Month |
|--|-----------------------------|
| Type of Activity | Per FTE |
| Communications with parents | 73.1 |
| Communications with staff | 23.5 |
| Communications with community agencies/providers | 3.3 |
| Case management meetings | 1.4 |

Source: Monthly Activities Reports submitted by districts in the Essential School Health Services program.

For children with special health care needs, nursing case management involves the development of Individual Health Care Plans (IHCPs) designed to maximize their potential for learning. An IHCP, usually developed by the school nurse in conjunction with the student's family, the school physician, other school staff, and relevant community health care providers, is an individualized care plan that stipulates a student's specific medical, nursing, emergency care, and educational needs while in school during the school day. IHCPs are reviewed on a regular basis to ensure that students receive the appropriate health care they need during the school day.

During the 2009-2010 school year, 77 ESHS districts reported:

- a total of 33,680 IHCPs for the year, with the median district reporting 170 IHCPs (range: 0 to 4,550 IHCPs);
- a median rate of 22.9 IHCPs per full-time school nurse (range: 0 to 170.2 IHCPs per full-time school nurse).

Program Development

School nurses perform program planning and development activities in coordination with other school district professionals, in areas such as environmental health, policy development, crisis management, and emergency preparedness. In addition, nurses attend meetings that contribute to their professional development. Meetings may be held at a specific school building or at the school district level. During the 2009-2010 school year, school nurses in 78 districts attended 1,166.3 program and professional development meetings per month (Table 19). Partner districts, partner collaboratives, and private schools conducted an additional 282.4 meetings per month.

| TABLE 19. Number of Program Development Meetings Attended by School Nurses, by Topic Area | | | | |
|--|------------------------|--|--|--|
| September 1, 2009 - Jun | ne 30, 2010 | | | |
| | Number of Meetings Per | | | |
| Topic Area | Month (All Districts) | | | |
| Crisis Management | 115.9 | | | |
| Emergency Preparedness | 86.7 | | | |
| Environmental | 16.1 | | | |
| Mental Health | 84.1 | | | |
| Policy Development | 107.0 | | | |
| Professional Development | 379.0 | | | |
| Other | 377.5 | | | |
| Total | 1,166.3 | | | |

Source: Monthly Activities Reports submitted by 78 districts in the Essential School Health Services program.

Students With Special Health Care Needs

1. Types of Special Health Care Needs

School nurses provide care for students with a wide variety of special health care needs. Table 19 shows the rates by type of condition. These rates are based on information provided to the school nurse by the student's primary care provider, who conducts a physical examination and submits a School Health Record once every 3 to 4 years. This information is supplemented by parent reports (on emergency cards and health information forms) submitted annually. Conditions not requiring special nursing care in school may be less likely to be reported to school nurses. For those conditions, these data may under-count the true rate in the student population. In the ESHS funded and partner schools that reported these data (136 funded and partner districts, and 5 collaboratives), the total enrollment was 622,368 (65% of the total public school enrollment in Massachusetts). In these schools, a total of 176,034 students with special health care needs were reported to school nurses (28% of enrollment). The most commonly reported physical/developmental condition is asthma (Table 19). The asthma rate among the schools reporting increased from 97.7 in 2006-2007 to 124.5 per 1,000 students in 2009-2010. Other common conditions include allergies, migraine headaches, seizure disorder, and cardiac conditions. The most commonly reported behavioral/emotional condition is Attention-Deficit/Hyperactivity Disorder (ADHD).

TABLE 20: Number of Students With Special Health Care Needs Reported to School Nurses in Selected Massachusetts Districts (Number and Rate Per 1,000 Enrolled Students) September 1, 2009 - June 30, 2010

| September 1, 2007 June 30 | | Rate Per 1,000 |
|--|-----------------|-----------------|
| | Number | Students |
| | (All Districts) | (All Districts) |
| Physical/Developmental Conditions | | |
| Allergies: | | |
| Bee Sting Allergies | 3,471 | 5.6 |
| Food Allergies | 26,712 | 42.9 |
| Latex Allergies | 1,293 | 2.1 |
| Asthma | 77,507 | 124.5 |
| Autoimmune Disorders (Arthritis, Lupus, etc.) | 1,177 | 1.9 |
| Blood Dyscrasias: | | |
| Hemophilia | 133 | 0.2 |
| Sickle Cell Disease | 684 | 1.1 |
| Other Blood Dyscrasias | 1,651 | 2.7 |
| Cancer | 543 | 0.9 |
| Cardiac Conditions | 4,986 | 8.0 |
| Celiac Disease | 840 | 1.3 |
| Cystic Fibrosis | 226 | 0.4 |
| Diabetes Type I | 1,863 | 3.0 |
| Diabetes Type II | 290 | 0.5 |
| Inflammatory Bowel Disease (IBS, Crohn's, etc) | 2,320 | 3.7 |
| Migraine Headaches | 7,280 | 11.7 |
| Neurologic Conditions: | | |
| Cerebral Palsy | 1,066 | 1.7 |
| Spina Bifida | 210 | 0.3 |
| Seizure Disorder | 5,230 | 8.4 |
| Neuromuscular Degenerative Disorder | 879 | 1.4 |
| Other Physical/ Developmental conditions | 16,806 | 27.0 |
| Behavioral/Emotional Conditions | | |
| ADHD/ADD | 35,460 | 57.0 |
| Autism | 6,617 | 10.6 |
| Depression | 7,044 | 11.3 |
| Eating Disorders | 1,133 | 1.8 |
| Other Behavioral/Emotional conditions | 15,596 | |
| Total Students With Special Health Care Needs | 176,034 | 282.8 |

Source: Status Reports submitted by 77 ESHS funded districts, 59 partner districts, and 5 collaboratives.

2. Students With Do Not Resuscitate (DNR) Orders

For some students who are terminally ill, parents and medical providers may determine that cardio pulmonary resuscitation should not be performed, and a Comfort Care/Do Not Resuscitate order will be prepared. During the school year, 6 students with DNR orders were reported to school nurses.

3. Cardiovascular Health and Automated Electronic Defibrillators (AEDs)

An automated external defibrillator (AED) is a portable device used to restore normal heart rhythm to patients in cardiac arrest. If cardiac arrest is not treated within a few minutes, the condition is fatal. AEDs located in ESHS districts were used 4 times during the school year (2 times with a student, and 2 times with staff). In 2 of those cases, use of the AED successfully restored a heart rhythm and the patient had a pulse when Emergency Medical Services (EMS) arrived.

Almost 78% of the ESHS school districts have at least one AED in all of their school buildings, up from 29.7% in 2003-2004 (Table 21) and 68.4 % last year. All ESHS districts have deployed AEDs in at least one school building. Only 17.8% of school buildings in ESHS districts do not have an AED.

| TABLE 21. Deployment of Automated External Defibrillators (AEDs) | | | | | | | | |
|--|--------|------|-------|------|--|--|--|--|
| in ESHS School Buildings and Districts | | | | | | | | |
| | 2003-2 | 2004 | 2009- | 2010 | | | | |
| | n % | | n | % | | | | |
| Total buildings | 870 | | 918 | | | | | |
| AED Status of Building | | | | | | | | |
| No AEDs | 596 | 68.5 | 163 | 17.8 | | | | |
| One AED | 218 | 25.1 | 598 | 65.1 | | | | |
| More than One AED | 56 | 6.4 | 157 | 17.1 | | | | |
| Total districts | 91 | | 77 | | | | | |
| AED Status of District | | | | | | | | |
| No AEDs in any building | 30 | 33.0 | 0 | 0.0 | | | | |
| At least one AED in all buildings | 27 | 29.7 | 60 | 77.9 | | | | |
| At least one building with more than one AED | 36 | 39.5 | 71 | 92.2 | | | | |


Source: Status Reports submitted by districts in the Essential School Health Services program.

Note: Since the group of districts participating in the ESHS program is not the same as it was in 2003-2004, the number of buildings is greater than it was in 2003-2004 even though the number of districts is smaller.

Client Satisfaction

In order to assess parents' perceptions of the quality of care that their students receive at school, a client satisfaction survey was conducted. Parents of students who received school health services were asked to complete a brief questionnaire. Each district is surveyed once every three years. In these districts, parents of approximately 100 students receiving health services are mailed a questionnaire and then requested to complete the questionnaire and return it to DPH. Parents of students at all grade levels are included in the sample. In the 2009-2010 school year, 1,009 parents returned completed questionnaires (38.8% of the 2,600 parents who were mailed questionnaires). Parental satisfaction rates on the measured criteria ranged from 93 to 97 percent (Figure 5).

FIGURE 5. Percentage of Parents Satisfied with School Nursing Services in ESHS Districts 2005-06 (n = 1,323), 2006-07 (n = 1,663), 2007-08 (n = 1,599), 2008-09 (n = 1,193), 2009-10 (n=1,009)



Detailed description of the Satisfaction Criteria:

- 1. I am very satisfied with the care my child receives from the school nurse.
- 2. If I have a question or concern, I can reach the school nurse for help without any problem.
- 3. The school nurse does his or her best to keep me informed about my child.
- 4. In an emergency at school, my child can get nursing care quickly.
- 5. The school nurse treats my child with respect.
- 6. I value the advice given by the school nurse

^{*} Parents were deemed "Satisfied" if they "Agreed" or "Strongly Agreed" with the statement.

Actions to Promote Healthy Weight

Previously Funded Essential School Health Services Districts

As part of the Essential School Health Service Performance Improvement Program, portions of the questionnaire entitled "Nutrition, Exercise, and Obesity: What's happening in Your School Districts?" are distributed annually by SurveyMonkey to the nurse leaders in the Essential School Health Service program. While each respondent represents an entire district, not all schools in a district would necessarily answer the same way. For example, actions which may be fully in place among elementary schools may not be in place among high schools. Shown in the tables below are the FY2011 responses of the 61 school districts whose funding was renewed in FY2009, and provides comparison figures for districts funded by the ESHS program in prior years. In Table 22 below is the percentage of respondents reporting obesity concerns. Shown in tables 23 through 24 below is the percentage of respondents who reported the school action or policy as being either fully or partially in place.

Obesity Concern in the Community

Most Nurse Leaders (93%) report that obesity is a concern in the community. Most (90%) report some school efforts to improve the nutritional quality of meals and snacks available to students. While 90% report school staff support these efforts, parents were seen as somewhat less supportive of these efforts (69%).

Table 22. Obesity concerns

| Obesity and Nutrition | 05-06 | 06-07 | 07-08 | 08-09 | 09-10 | 10-11 |
|---|-------|-------|-------|-------|-------|-------|
| 1. Obesity is a concern in the community? | 60% | 70% | 74% | 87% | 91% | 93% |
| 2. Efforts are being made in school to improve the nutritional quality of meals and snacks available to students? | 68% | 84% | 83% | 96% | 95% | 97% |
| 3. School staff support efforts to improve the nutritional quality of meals and snacks, for example, reduce fat and/or caloric content or replace sugared drinks with water or 100% juices? | 56% | 70% | 71% | 87% | 86% | 90% |
| 4. Parents support efforts to improve the nutritional quality of meals and snacks, for example, reduce fat and/or caloric content or replace sugared drinks with water or 100% juices? | 37% | 42% | 50% | 59% | 58% | 69% |

²⁰ The survey includes a sample of questions from the CDC's School Health Index.

38

* Percentages of respondents reporting affirmatively on a 1-5 point scale with 1 = yes, a lot; 5 = no, not much.

Physical Activity

Less than half of the respondents report that all students receive at least 150 minutes of PE per week. In addition, while less than half of the respondents report that their schools spread PE over at least 3 days per week, this number has almost doubled over the past five years.

Table 23. School actions undertaken to increase physical activity

| Increased Physical Activity | 05-06 | 06-07 | 07-08 | 08-09 | 09-10 | 10-11 |
|-------------------------------------|-------|-------|-------|-------|-------|-------|
| 5. Providing at least 20 minutes of | 85% | 93% | 93% | 94% | 91% | 92% |
| recess each day | | | | | | |
| 5a. Monitors encouraging students | 80% | 84% | 88% | 89% | 89% | 95% |
| to be active at recess | | | | | | |
| 6. Using a sequential PE curriculum | 90% | 95% | 96% | 98% | 98% | 97% |
| that is consistent with state or | | | | | | |
| national standards | | | | | | |
| 7. All students receiving at least* | 21% | 24% | 32% | 32% | 35% | 41% |
| 150 minutes of PE per week | | | | | | |
| 7a. Spreading PE over at least 3 | 23% | 28% | 33% | 38% | 48% | 39% |
| days (preferably 5 days) per week | | | | | | |
| 8. Promoting walking /biking to | 35% | 40% | 46% | 58% | 54% | 59% |
| school | | | | | | |

Nutrition

School actions to improve nutrition are reported in Table 24. Most respondents reported their schools provided a variety of foods on menus and offered low-fat and skim milk every day, but a much smaller percentage of respondents reported that their schools offered appealing low fat items in vending machines, parties, and after-school programs.

Table 24. School actions to improve nutrition

| Improved nutrition | 05-06 | 06-07 | 07-08 | 08-09 | 09-10 | 10-11 |
|---------------------------------------|-------|-------|-------|-------|-------|-------|
| 9. Providing a variety of foods on | 86% | 89% | 94% | 92% | 97% | 93% |
| school menus | | | | | | |
| 10. Offering low-fat and skim milk | 98% | 99% | 100% | 100% | 100% | 98% |
| every day | | | | | | |
| 11. Offering at least one appealing | 91% | 94% | 96% | 91% | 95% | 98% |
| low fat item from each of the | | | | | | |
| following food groups every day: | | | | | | |
| fruits, vegetables, grains, and dairy | | | | | | |
| products? | | | | | | |
| 12. Allowing ample time for lunch | 79% | 82% | 91% | 85% | 95% | 90% |

^{*} Please note in late FY07 the MDPH issued the Comprehensive Growth Screening Guidelines which will facilitate school districts in addressing these issues.

| and breakfast | | | | | | |
|-----------------------------------|-----|-----|-----|-----|-----|-----|
| 13. Vending machines exist in | 94% | 96% | 92% | 88% | 92% | 95% |
| school? | | | | | | |
| 14. Restricting access to vending | 91% | 93% | 96% | 88% | 88% | 93% |
| machines (among districts with | | | | | | |
| vending machines) | | | | | | |
| 15a. Offering appealing low fat | 32% | 42% | 54% | 56% | 40% | 38% |
| items in vending machines | | | | | | |
| 15b. Offering appealing low fat | 25% | 35% | 51% | 66% | 63% | 43% |
| items at parties | | | | | | |
| 15c. Offering appealing low fat | 27% | 35% | 45% | 66% | 65% | 62% |
| items at after school programs | | | | | | |

School Nurse

School nurse actions to improve physical activity and nutrition are reported in Table 25. Compared to five years ago, school nurses are now more likely to collaborate to improve physical activity and nutrition, and to have a system in place for measuring BMIs . In addition, there was a large increase in the percentage of respondents reporting BMIs to families and physicians, and in the percentage of respondents using a written protocol for managing students identified as at risk for weight.*

Table 25. School nurse actions to improve physical activity and nutrition

| Improved physical activity and nutrition | 05-06 | 06-07 | 07-08 | 08-09 | 09-10 | 10-11 |
|--|-------|-------|-------|-------|-------|-------|
| 16. Promoting physical activity | | | | | | |
| through: | | | | | | |
| Educational materials | 83% | 83% | 92% | 92% | 91% | 97% |
| Individual advice | 95% | 95% | 100% | 100% | 100% | 98% |
| Small groups | 53% | 50% | 67% | 68% | 58% | 54% |
| Presentations | 55% | 67% | 75% | 64% | 65% | 61% |
| 17. Promoting healthy eating | | | | | | |
| through: | | | | | | |
| Educational materials | 85% | 89% | 94% | 88% | 95% | 95% |
| Individual advice | 96% | 96% | 100% | 97% | 98% | 100% |
| Small groups | 57% | 54% | 62% | 48% | 57% | 51% |
| Presentations | 56% | 68% | 70% | 65% | 66% | 66% |
| 18. Collaborating to promote healthy | | | | | | |
| eating and physical activity through: | | | | | | |
| Policy development | 66% | 89% | 87% | 89% | 91% | 95% |
| Curriculum development | 47% | 55% | 67% | 82% | 77% | 75% |
| Unit and lesson planning | 44% | 49% | 62% | 72% | 62% | 69% |
| Special events/planning | 56% | 59% | 72% | 77% | 78% | 84% |

| In service training | 40% | 50% | 63% | 58% | 60% | 62% |
|-------------------------------------|-----|-----|-----|-----|------|------|
| Having a system in place to measure | 90% | 93% | 99% | 98% | 100% | 98% |
| student BMI's | | | | | | |
| Reporting BMI's to students family | 49% | 46% | 55% | 66% | 94% | 100% |
| Reporting BMI's to students | 19% | 17% | 20% | 17% | 52% | 43% |
| physicians | | | | | | |
| Managing students identified as at | 13% | 12% | 21% | 14% | 36% | 36% |
| risk for weight using a written | | | | | | |
| protocol | | | | | | |

Newly Funded Essential School Health Service Districts

Shown in the tables below are the FY2011 responses of the 9 school districts whose ESHS funding began in FY09 (and who had not been funded the prior year). The percentages reported below may be expected to fluctuate from year to year due to the small number of respondents in this group.

Shown in Table 26 below is the percentage of respondents reporting obesity concerns. Shown in tables 27 through 29 below is the percentage of respondents who reported the school action or policy as being either fully or partially in place.

Obesity Concern in the Community

As shown by Table 26, concerns about obesity are reported to have increased in the communities served by newly funded districts. Most of the respondents report that school and school staff are making efforts to improve the nutritional quality of meals and snacks available to students. Support for these efforts among parents has increased.

Table 26. Obesity concerns

| Questions | 08-09 | 09-10 | 10-11 |
|---|-------|-------|-------|
| 1. Obesity is a concern in the community? | 73% | 91% | 89% |
| 2. Efforts are being made in school to improve the nutritional quality of meals and snacks available to students? | 91% | 100% | 100% |
| 3. School staff support efforts to improve the nutritional quality of meals and snacks, for example, reduce fat and/or caloric content or replace sugared drinks with water or 100% juices? | 91% | 91% | 89% |
| 4. Parents support efforts to improve the nutritional quality of meals and snacks, for example, reduce fat and/or caloric content or replace sugared drinks with water or 100% juices? | 55% | 73% | 56% |

Physical Activity

The action these schools were least likely to undertake was "all students receiving at least 150 minutes of PE per week."

Table 27. School actions undertaken to increase physical activity

| Increased Physical Activity | 08-09 | 09-10 | 10-11 |
|---|-------|-------|-------|
| 5. Providing at least 20 minutes of recess | 100% | 91% | 100% |
| each day | | | |
| 5a. Monitors encouraging students to be | 82% | 91% | 89% |
| active at recess | | | |
| 6. Using a sequential PE curriculum that is | 91% | 91% | 100% |
| consistent with state or national standards | | | |
| 7. All students receiving at least 150 | 27% | 36% | 56% |
| minutes of PE per week | | | |
| 7a. Spreading PE over at least 3 days | 36% | 45% | 67% |
| (preferably 5 days) per week | | | |

Nutrition

All of these schools offer low fat items on menus. Only about half offer low fat items in vending machines, parties, or after school programs.

Table 28. School actions to improve nutrition

| Improved nutrition | 08-09 | 09-10 | 10-11 |
|---|-------|-------|-------|
| 9. Providing a variety of foods on school | 91% | 100% | 100% |
| menus | | | |
| 10. Offering low-fat and skim milk every | 100% | 100% | 100% |
| day | | | |
| 11. Offering at least one appealing low fat | 91% | 91% | 100% |
| item from each of the following food | | | |
| groups every day: fruits, vegetables, | | | |
| grains, and dairy products? | | | |
| 12. Allowing ample time for lunch and | 82% | 73% | 100% |
| breakfast | | | |
| 13. Vending machines exist in school? | 91% | 82% | 89% |
| 14. Restricting access to vending machines | 90% | 91% | 89% |
| (among districts with vending machines) | | | |
| 15a. Offering appealing low fat items in | 27% | 45% | 44% |
| vending machines | | | |
| 15b. Offering appealing low fat items at | 36% | 55% | 56% |
| parties | | | |
| 15c. Offering appealing low fat items at | 36% | 55% | 44% |
| after school programs | | | |

School Nurse

Compared to the prior year, more school districts reported promoting physical activity and healthy eating through small group activities, and more school districts reported collaborating to promote these activities. In addition, all of the respondents now report a system in place for measuring BMI's. The percentage of districts that have procedures for reporting BMIs to families and physicians and for managing students at risk for weight increased over the prior year,.²¹

²¹ Please note: In late FY07 the MDPH issued the Comprehensive Growth Screening Guidelines which will facilitate school districts in addressing these issues.

Table 28. School nurse actions to improve physical activity and nutrition

| 1 able 28. School nurse actions to improve | pnysicai ac | uvity and i | 141111011 |
|---|-------------|-------------|-----------|
| Improved physical activity and nutrition | 08-09 | 09-10 | 10-11 |
| 16. Promoting physical activity through: | | | |
| Educational materials | 73% | 91% | 100% |
| Individual advice | 82% | 100% | 89% |
| Small groups | 18% | 27% | 67% |
| Presentations | 27% | 45% | 33% |
| 17. Promoting healthy eating through: | | | |
| Educational materials | 82% | 82% | 100% |
| Individual advice | 100% | 100% | 100% |
| Small groups | 18% | 45% | 56% |
| • Presentations | 45% | 55% | 67% |
| 18. Collaborating to promote health eating | | | |
| and physical activity through: | | | |
| Policy development | 73% | 64% | 89% |
| Curriculum development | 55% | 64% | 78% |
| Unit and lesson planning | 27% | 45% | 78% |
| Special events/planning | 36% | 55% | 89% |
| In service training | 18% | 55% | 44% |
| Having a system in place to measure | 82% | 100% | 100% |
| student BMI's | | | |
| Reporting BMI's to students' families | 18% | 64% | 100% |
| Reporting BMI's to students' physicians | 0% | 36% | 44% |
| Managing students identified as at risk for | 0% | 9% | 33% |
| weight using a written protocol | | | |

References

Chabra, A. & Chavez, G. (2000). A comparison of long pediatric hospitalizations in 1985 and 1994. Journal of Community Health, 25(3), 199-210.

Cedar Rapids Community School District v. Garret F, 119 S.Ct. 992, 29 IDELR 966 (U.S. 199). To view the full decision, go to: http://supct.law.cornell.edu/supct/html/96-1793.ZS.html

Centers for Disease Control and Prevention (nd). Number (in Millions) of Civilian/Noninstitutionalized Persons with Diagnosed Diabetes, United States, 1980–2006. Retrieved January 11, 2010 from http://www.cdc.gov/diabetes/statistics/incidence/fig1.htm

Clements KM, Barfield WD, Ayadi F, Wilber N. Preterm birth-associated cost of early intervention services: an analysis by gestational age. <u>Pediatrics</u> 2007;119:e866–e874.

Coffman JM, Cabana MD, Halpin HA, Yelin EH. (2008) Effects of asthma education on children's use of acute care services: a meta-analysis. Pediatrics. 121(3):575-86.

Expert Committee Recommendations on the Assessment, Prevention and Treatment of Child and Adolescent Overweight and Obesity (2007). Retrieved January 8, 2008 from http://www.ama-assn.org/ama1/pub/upload/mm/433/ped_obesity_recs.pdf.

Hannon, TS; Rao, G; Arslanian, SA (2005). Childhood Obesity and Type 2 Diabetes Mellitus. Pediatrics, 116 (2), 473-480

Leslie, L., Sarah, R., & Palfrey, J. S. (1998). Child health care in changing times. <u>Pediatrics</u>, <u>101</u>(4), 746-751.

Massachusetts Department of Elementary and Secondary Education (2008). State Profile: Enrollment Data. Retrieved October 7, 2008 from http://profiles.doe.mass.edu/.

Massachusetts Department of Public Health (2009). Data Health Brief: Epinephrine Administration In Schools (School Year 2007-2008).

McPherson M, Arango P, Fox H, Lauver C, McManus M, Newacheck P, Perrin J, Shonkoff J, Strickland B. (1998) A new definition of children with special health care needs. <u>Pediatrics</u>, 102(1):137–140.

Palfrey, J.S., Haynie, M., Porter, S., Bierle, T., Cooperman, P., Lowcock, J. (1992). Project school care: Integrating children assisted by medical technology into educational settings. Journal of School Health, 62(2), 50-54.

Pbert, L., Osganian, S., Gorak, D., Druker, S., Reed, G., O'Neill, K., Sheetz, A. (2006). School Nurse-delivered Adolescent Smoking Cessation Intervention: A Randomized Controlled Trial, Preventive Medicine: 43: 313-320.

Pennington N & Delaney E. (2008). The number of students sent home by school nurses compared to unlicensed personnel. <u>Journal of School Nursing</u> 24(5):290-7.

Raymond JA. (2009). The integration of children dependent on medical technology into public schools. <u>Journal of School Nursing</u>;25(3):186-94. Epub 2009 Apr 10.

Schutte, E. B., Price, D. L., & James, S. R. (1997). <u>Thompson's Pediatric Nursing.</u> Philadelphia: W. B. Saunders.

Sheetz, A, Developing School Health Services in Massachusetts: A Public Health Model. <u>Journal of School Nursing</u>. 2003; 19(4): 204-211.

Small, M.L., Majer, L.S., Allensworth, D.D., Farquhar, B.K., Kann, L., & Pateman, B.C. (1995). School health services. Journal of School Health, 65(8), 319-326.

Smolensky, E. & Gootman, JA. (Eds.). (2003). <u>Working Families and Growing Kids: Caring for Children and Adolescents.</u> Washington: National Academies Press. Retrieved February 8, 2011 from http://www.nap.edu/catalog.php?record_id=10669.

Thurber, F., Berry, B., & Cameron, M.E. (1991). The role of school nursing in the United States. <u>Journal of Pediatric Health Care</u>, 5(3), 135-140.

Uphold, C.R. & Graham, M.V. (1993). Schools as centers for collaborative services for families: A vision for change. <u>Nursing Outlook</u>, 41(5), 204-211.

U.S. Bureau of the Census. (2000). <u>Statistical Abstract of the United States</u> (120th ed.). Washington, D.C.; 60 & 655.

U.S. Department of Health and Human Services. (2000). U.S. Public Health Service, <u>Report of the Surgeon General's Conference on Children's Mental Health</u>. Washington, DC: Author. Retrieved January 8, 2010, from http://www.surgeongeneral.gov/topics/cmh/cmhreport.pdf

Wold, S.J. (2001). School health services: History and trends. In N.C. Schwab & M.H.B. Gelfman (Eds.), <u>Legal issues in school health services</u> (pp. 7-54). North Branch, MN: Sunrise River Press.

Wyman, L. (2005). Comparing the number of ill or injured students who are released early from school by school nursing and non-nursing personnel. <u>Journal of School Nursing</u>, 21(6), 350-355.

APPENDIX A

School Districts and Student Enrollment Essential School Health Services Program: 2009-2010

| DISTRICT NAME | REGION | ADMINISTRATION | ENROLLMENT |
|---------------------------------|------------|-------------------|------------|
| 1 Acton-Boxborough | Metro West | Regional Academic | 6,039 |
| 2 Amesbury | Northeast | City or Town | 2,474 |
| 3 Andover | Northeast | City or Town | 6,163 |
| 4 Arlington | Metro West | City or Town | 4,713 |
| 5 Ashburnham-Westminster | Central | Regional Academic | 2,388 |
| 6 Attleboro | Southeast | City or Town | 5,933 |
| 7 Barnstable | Southeast | City or Town | 5,421 |
| 8 Belchertown | Western | City or Town | 2,610 |
| 9 Berkshire Hills (Stockbridge) | Western | Regional Academic | 1,377 |
| 10 Billerica | Northeast | City or Town | 5,940 |
| 11 Boston | Boston | City or Town | 55,371 |
| 12 Braintree | Metro West | City or Town | 5,377 |
| 13 Bridgewater Raynham | Southeast | Regional Academic | 5,804 |
| 14 Brockton | Southeast | City or Town | 15,502 |
| 15 Brookline | Boston | City or Town | 6,472 |
| 16 Cambridge | Metro West | City or Town | 5,950 |
| 17 Canton | Metro West | City or Town | 3,125 |
| 18 Central Berkshire (Dalton) | Western | Regional Academic | 1,987 |
| 19 Chelsea | Boston | City or Town | 5,638 |
| 20 Chicopee | Western | City or Town | 7,845 |
| 21 Douglas | Central | City or Town | 1,771 |
| 22 East Longmeadow | Western | City or Town | 2,850 |
| 23 Fall River | Southeast | City or Town | 9,886 |
| 24 Fitchburg | Central | City or Town | 4,997 |
| 25 Framingham | Metro West | City or Town | 8,153 |
| 26 Gardner | Central | City or Town | 2,600 |
| 27 Gateway (Huntington) | Western | Regional Academic | 1,202 |
| 28 Georgetown | Northeast | City or Town | 1,688 |
| 29 Gill-Montague | Central | Regional Academic | 1,085 |
| 30 Gloucester | Northeast | City or Town | 3,372 |
| 31 Granby | Western | City or Town | 1,125 |
| 32 Hadley | Western | City or Town | 714 |
| 33 Hampden Wilbraham | Western | Regional Academic | 3,600 |
| 34 Hampshire | Western | School Union | 1,865 |

| DISTRICT NAME | REGION | ADMINISTRATION | ENROLLMENT |
|---------------------------------|------------|-------------------|------------|
| 35 Harwich | Southeast | City or Town | 1,334 |
| 36 Haverhill | Northeast | City or Town | 6,845 |
| 37 Holyoke | Western | City or Town | 5,901 |
| 38 Hudson | Metro West | City or Town | 3,071 |
| 39 Lawrence | Northeast | City or Town | 12,284 |
| 40 Leominster | Central | City or Town | 6,290 |
| 41 Lexington | Metro West | City or Town | 6,182 |
| 42 Lowell | Northeast | City or Town | 13,331 |
| 43 Ludlow | Western | City or Town | 3,050 |
| 44 Lynn | Northeast | City or Town | 13,373 |
| 45 Mansfield | Southeast | City or Town | 4,888 |
| 46 Marblehead | Northeast | City or Town | 3,232 |
| 47 Marshfield | Southeast | City or Town | 4,746 |
| 48 Medford | Northeast | City or Town | 4,854 |
| 49 Middleborough | Southeast | City or Town | 3,506 |
| 50 Nashoba | Central | Regional Academic | 3,433 |
| 51 Natick | Metro West | City or Town | 4,734 |
| 52 Needham | Metro West | City or Town | 5,311 |
| 53 New Bedford | Southeast | City or Town | 12,636 |
| 54 Newburyport | Northeast | City or Town | 2,251 |
| 55 Newton | Metro West | City or Town | 11,765 |
| 56 North Andover | Northeast | City or Town | 4,614 |
| 57 North Attleborough | Southeast | City or Town | 4,750 |
| 58 North Berkshire (Clarksburg) | Western | School Union | 332 |
| 59 Northampton | Western | City or Town | 2,692 |
| 60 Northboro Southboro | Metro West | School Union | 4,848 |
| 61 Northbridge | Central | City or Town | 2,539 |
| 62 Pittsfield | Western | City or Town | 6,072 |
| 63 Plymouth | Southeast | City or Town | 8,240 |
| 64 Provincetown | Southeast | City or Town | 152 |
| 65 Quincy | Metro West | City or Town | 8,969 |
| 66 Randolph | Metro West | City or Town | 2,851 |
| 67 Rockport | Northeast | City or Town | 977 |
| 68 Sandwich | Southeast | City or Town | 3,579 |
| 69 Scituate | Metro West | City or Town | 3,278 |
| 70 Somerville | Metro West | City or Town | 4,842 |
| 71 Springfield | Western | City or Town | 25,591 |
| 72 Stoughton | Southeast | City or Town | 3,776 |
| 73 Taunton | Southeast | City or Town | 7,920 |
| 74 Walpole | Metro West | City or Town | 3,954 |

| | DISTRICT NAME | REGION | ADMINISTRATION | ENROLLMENT |
|----|------------------|------------|----------------|------------|
| 75 | Waltham | Metro West | City or Town | 4,763 |
| 76 | West Bridgewater | Southeast | City or Town | 1,292 |
| 77 | Weston | Metro West | City or Town | 2,388 |
| 78 | Weymouth | Metro West | City or Town | 6,919 |
| 79 | Wilmington | Metro West | City or Town | 3,783 |
| 80 | Worcester | Central | City or Town | 23,988 |
| | TOTAL | | | 477,163 |

Notes:

Source: Massachusetts Department of Elementary and Secondary Education (DESE) ESHS-funded districts may include schools not included in DESE -defined districts, so the enrollment numbers shown above may differ from those provided by DESE.

[&]quot;Region" refers to the six geographic regions defined by the Executive Office of Health and Human Services (EOHHS).

APPENDIX B

Scope of Service
Essential School Health Services Program

COMPONENTS

Each program must meet or continue to meet the following seven components as described below:

- 1. School health service program infra-structure
- 2. Collaboration with the comprehensive, coordinated health education program, tobacco control program, etc.
- 3. Plan for linkage of students with primary care providers, dental providers, behavioral/mental health programs (as needed), community prevention programs, and health care insurance.
- 4. Development of a management information system.
- 5. Implementation of performance improvement (continuous quality improvement) and evaluation programs.
- 6. Services to private schools located in the applicant's community
- 7. Collaboration/consultation/networking among school nurses.

For a more complete description of each of these components, please contact the School Health Unit.

APPENDIX C

Data Collection Methods

Contractual obligations require districts in the ESHS programs to submit a monthly report to MDPH. This report, the ESHS **Monthly Activities Report**, provides a detailed, standardized summary of the health services activities that took place in the district during the prior month. It includes a count of the number of encounters, medications administered, medical procedures, and other types of services provided.

Information for these reports is gathered from each school nurse. In most districts, school nurses enter health encounter data into a computer database loaded on a computer located in the school health office. The database facilitates data reporting as well as helps the nurse maintain systematic records and schedule follow-ups. ²² Nurses are encouraged to enter information during or directly after a health encounter. Each district in the ESHS program selects its own database software. Across the program, ten or more different software products are used, although the majority of districts use one of two popular applications. Within a district, all school nurses usually use the same software product. The software products operate differently. Many districts use a networked database that links all schools to the same database and permits the data coordinator to run district-wide data reports, while other districts use stand-alone databases in which data reports must be run separately at each school before being compiled at the district level. Due to resource constraints, nurses in a few school districts maintain paper logs and manually tabulate the data. Although districts use different software applications and some districts tabulate data manually, all districts are required to tabulate their data the same way and to submit a standard data report to MDPH. In any event, information is gathered from each school nurse in the district, tabulated, and entered into the Monthly Activities Report form in summary (or aggregate) form.

In addition, districts in the ESHS programs submit **status reports** once a year. This report measures progress in meeting program objectives, and includes performance measures relating to health services infrastructure, MIS development, linkages to all aspects of the health delivery system, and quality evaluation. It also summarizes the number of health screenings performed and health surveys administered during the school year. The mentored school districts in the program submit this report once a year, beginning in 2009-2010.

The statistics in this report were derived from the monthly activities reports submitted by districts participating in the ESHS program. Over the course of the 2009-2010 school year, monthly encounter data were collected successfully from 78 of the 80 ESHS award recipients. For these school systems, MDPH received 752 (96%) of the 780 expected monthly reports.

For the 78 districts that form the basis of this report, the median student enrollment was 14,014, with a range of 152 to 55,371 students. This sample includes school districts from many areas of

²² Paper logs are still used to record data elements that are not typically included in most school health software programs. For example, one item that is usually logged by hand is "Number of support group meetings."

the state. It includes urban, suburban, and rural districts; city, town, and regional school systems; and large, medium, and small districts.

Data Analysis Methods

In order to reduce the potential for confusion, the statistical concepts and terms used in this report are described below.

For each measurement or "indicator," a *district-level statistic* is determined in each district by calculating a monthly average for the 10-month evaluation period. The **monthly average** for a particular district is calculated by adding the total number of events or encounters that occurred in a particular district during the evaluation period and dividing that total by the number of months included in that evaluation period. Because it is awkward to refer constantly to the "monthly average for the district" or the "district-based monthly average," these data are referred to as the **district average**. These two terms--the monthly average and district average--are used interchangeably in this report. All monthly averages in this report were calculated over the same ten-month period (September through June).

Wherever possible, standard units of analyses (rates) are used, as they facilitate both crossdistrict and historical comparisons, which can provide context and meaning to the statistics. The standard units of analysis that were used most frequently in this report are the monthly rate per 1,000 student health encounters, the monthly rate per 1,000 enrolled students, and the monthly rate per full-time equivalent (FTE) nurse. The monthly rate per 1,000 student health encounters is calculated by dividing the monthly average for that indicator by the total number of student health encounters in that district and multiplying the result by 1,000. Similarly, the monthly rate per 1,000 enrolled students is calculated by dividing the monthly average by the total number of enrolled students in that district and multiplying the result by 1,000. Rates per thousand enrolled students were calculated utilizing October student enrollment figures provided by the Massachusetts Department of Education (see Appendix A). Finally, the **monthly rate per** full-time equivalent (FTE) nurse is calculated by dividing the monthly average by the total number of Registered Nurse FTEs in that district. Sometimes the rate is not based on an average of monthly data but on aggregate data for the full year. For example, the rate of health screenings per 1,000 students is determined by dividing the total number of screenings for the whole year by the number of students enrolled and multiplying the result by 1,000.

Program-wide statistics describe not individual districts, but the ESHS program as a whole. In these calculations, each district represents a data point that is used in calculating summary statistics. For example, if averages are calculated for 100 districts, the result is a collection of 100 district averages that can be arrayed from lowest to highest along a frequency distribution. When frequency distributions are *skewed* (that is, the values tend to clump around either the lowest or highest value, rather than around the middle), the *median*, rather than the *average*, is used to measure central tendency. *Because most of the ESHS frequency distributions were skewed, the median is used throughout this report.* The **median** represents the number above and below which exactly 50% of the districts fall. It is a better measure of central tendency than the *average* for skewed data, because the average tends to be more affected by extreme values. The most common use of median in this report is with district-based monthly averages; for a

particular indicator, the median for the group of ESHS districts (a *program-level* statistic) is the district average (or monthly average) above and below which exactly 50% of the individual district averages fell. The **range** of a set of district averages refers to the lowest and highest values across the entire group of ESHS districts. The district with the median value for an indicator is sometimes referred to as the **median district** or the **typical district**. The median value across all the monthly district averages is also referred to as the **median district average**.

Medians can also be calculated for rates. For example, the **median Emergency Referral rate** (i.e., Emergency Referrals per 1,000 health encounters) is calculated by first putting the total number of Emergency Referrals in the form of a rate (for each district, dividing the total number of Emergency Referrals by the number of student health encounters and multiplying by 1,000), and then finding the median of these rates.

Data Limitations

This report focuses on the delivery of school health services by nursing staff. Project sites do not serve as a representative sample of the Commonwealth's schools. Therefore this report should not be used to make generalized statements about health services in all Massachusetts public schools. Furthermore, caution should be exercised when comparing ESHS statistics across years. Each year the set of districts that report data changes to some degree, which creates somewhat different sample sets. For example, in the 2000-2001 school year, 74 districts reported data, whereas in the school year 2003-2004, 103 districts reported data. In addition, in years prior to 2001, the number of districts that reported data (approximately 25) was drastically lower than in more recent years (approximately 100). Due to this difference in data sets, comparisons to data from years prior to 2001 would be considerably less valid. Also, data has not always been available for all months of the school year. Most notably, in the 2002-2003 school year, only the months September through December were reported. This noted, after 2001 the core group of districts has been relatively stable, and the sample size is large enough such that comparisons are not without merit. Where statistical differences are large, and trends continue for several years, comparisons are more likely to be meaningful.

The descriptive data presented here also do not capture the dynamic and multi-faceted nature of health services delivery in a school system, which would require in-depth qualitative analysis of the program participants. Differences in data collection and data tabulation procedures may account for some of the variability observed across districts. Furthermore, a small percentage of the school districts in the program did not have computerized records of office visits and relied on paper logs and hand tallying of data by individual nurses. In these cases, it is impossible to control for factors such as data-entry errors at the district level, consistent misinterpretation of data elements, and numerical "guesstimates" provided by participants. Some of these data quality problems can lead to significant under- or over-counting. Finally, interpretation of the data is limited because we have not attempted to analyze the influence of school district demographics or other participant differences.

Participating districts were required to implement, in a short period of time, both program innovations that entailed major organizational change and, in most cases, the development of an internal data collection system. Therefore, this report represents a preliminary attempt to measure the health services activity in participating school systems. Improvements in data collection

procedures, data collection tools, and data collection instructions and training occur on a continuing basis, leading to corresponding improvements in data validity and reliability.