### Table 14A Cancer Incidence in Hingham, MA 1982-1994

| 1982-1994 |     |       | Total |      |      |     |       | Male | 1    |     |     |      | Female |    |      |
|-----------|-----|-------|-------|------|------|-----|-------|------|------|-----|-----|------|--------|----|------|
|           | Obs | Exp   | SIR   | 95   | % CI | Obs | Exp   | SIR  | 95 % | CI  | Obs | Exp  | SIR    | 95 | % CI |
| Bladder   | 37  | 46.6  | 79    | 56   | 109  | 28  | 34.1  | 82   | 55   | 119 | 9   | 12.6 | 72     | 33 | 136  |
| Brain     | 20  | 17.1  | 117   | 71   | 181  | 8   | 9.4   | 85   | 37   | 168 | 12  | 7.7  | 156    | 80 | 272  |
| Kidney    | 17  | 26.6  | 64    | 37   | 102  | 12  | 16.4  | 73   | 38   | 128 | 5   | 10.2 | 49     | 16 | 114  |
| Leukemia  | 15  | 22.0  | 68    | 38   | 112  | 8   | 12.5  | 64   | 27   | 126 | 7   | 9.5  | 74     | 30 | 152  |
| Liver     | 2   | 5.6   | NC    |      | NC   | 1   | 4.0   | NC   | NC   | 2   | 1   | 1.6  | NC     | ]  | NC   |
| Lung      | 145 | 175.0 | 83    | * 70 | 97   | 75  | 108.6 | 69   | * 54 | 87  | 70  | 66.4 | 105    | 82 | 133  |
| NHL       | 39  | 39.5  | 99    | 70   | 135  | 20  | 20.8  | 96   | 59   | 149 | 19  | 18.8 | 101    | 61 | 158  |
| Pancreas  | 14  | 24.4  | 57    | * 31 | 96   | 5   | 11.9  | 42   | * 14 | 98  | 9   | 12.5 | 72     | 33 | 137  |

### Notes:

Expected number of cases presented are rounded to the nearest tenth.

SIRs are calculated based on the exact number of expected cases.

SIRs and 95% Confidence Interval are not calculated when fewer than 5 cases are observed.

Obs = Observed number of cases

Exp = Expected number of cases

SIR = Standardized Incidence Ratio

NC = Not Calculated

95% CI = 95% Confidence Interval

\* Indicates statistical significance (P<.05)

Data source: Massachusetts Cancer Registry, Bureau of Health Statistics, Research and Evaluation, Mass. Dept. of Public Health

Table 14B Cancer Incidence in Hingham, MA 1982-1986 & 1987-1994

| 1982-1986 |     |      | Total |    |      |     |      | Male | 3      |    |     |      | Femal | e       |
|-----------|-----|------|-------|----|------|-----|------|------|--------|----|-----|------|-------|---------|
|           | Obs | Exp  | SIR   | 95 | % CI | Obs | Exp  | SIR  | 95 % ( | Ί  | Obs | Exp  | SIR   | 95 % CI |
| Bladder   | 11  | 17.8 | 62    | 31 | 111  | 9   | 12.8 | 70   | 32 13  | 33 | 2   | 4.9  | NC    | NC      |
| Brain     | 8   | 5.8  | 138   | 59 | 272  | 4   | 3.2  | NC   | NC     |    | 4   | 2.6  | NC    | NC      |
| Kidney    | 2   | 7.8  | NC    |    | NC   | 2   | 4.7  | NC   | NC     |    | 0   | 3.1  | NC    | NC      |
| Leukemia  | 6   | 8.3  | 73    | 27 | 158  | 3   | 4.6  | NC   | NC     |    | 3   | 3.7  | NC    | NC      |
| Liver     | 0   | 1.8  | NC    |    | NC   | 0   | 1.2  | NC   | NC     |    | 0   | 0.6  | NC    | NC      |
| Lung      | 57  | 60.2 | 95    | 72 | 123  | 26  | 39.0 | 67   | * 43 9 | 8  | 31  | 21.1 | 147   | 100 209 |
| NHL       | 11  | 12.5 | 88    | 44 | 158  | 5   | 6.4  | 78   | 25 18  | 32 | 6   | 6.1  | 98    | 36 214  |
| Pancreas  | 6   | 9.2  | 65    | 24 | 142  | 1   | 4.3  | NC   | NC     |    | 5   | 4.8  | 103   | 33 241  |

| 1987-1994 |     |       | Total |      |      |     |      | Male | 1       |     |      | Female |         |
|-----------|-----|-------|-------|------|------|-----|------|------|---------|-----|------|--------|---------|
|           | Obs | Exp   | SIR   | 95   | % CI | Obs | Exp  | SIR  | 95 % CI | Obs | Exp  | SIR    | 95 % CI |
| Bladder   | 26  | 28.7  | 91    | 59   | 133  | 19  | 21.1 | 90   | 54 141  | 7   | 7.6  | 92     | 37 190  |
| Brain     | 12  | 11.3  | 106   | 55   | 185  | 4   | 6.2  | NC   | NC      | 8   | 5.1  | 157    | 68 310  |
| Kidney    | 15  | 18.9  | 79    | 44   | 131  | 10  | 11.8 | 85   | 41 156  | 5   | 7.1  | 70     | 23 164  |
| Leukemia  | 9   | 13.7  | 66    | 30   | 124  | 5   | 7.9  | 63   | 20 147  | 4   | 5.8  | NC     | NC      |
| Liver     | 2   | 3.9   | NC    |      | NC   | 1   | 2.8  | NC   | NC      | 1   | 1.0  | NC     | NC      |
| Lung      | 88  | 114.3 | 77    | * 62 | 95   | 49  | 69.1 | 71   | * 52 94 | 39  | 45.3 | 86     | 61 118  |
| NHL       | 28  | 27.0  | 104   | 69   | 150  | 15  | 14.3 | 105  | 59 173  | 13  | 12.7 | 103    | 55 175  |
| Pancreas  | 8   | 15.2  | 53    | 23   | 104  | 4   | 7.5  | NC   | NC      | 4   | 7.7  | NC     | NC      |

### Notes:

Expected number of cases presented are rounded to the nearest tenth.

SIRs are calculated based on the exact number of expected cases.

SIRs and 95% Confidence Interval are not calculated when fewer than 5 cases are observed.

Obs = Observed number of cases

Exp = Expected number of cases

SIR = Standardized Incidence Ratio

NC = Not Calculated

95% CI = 95% Confidence Interval

\* Indicates statistical significance (P<.05)

Data source: Massachusetts Cancer Registry, Bureau of Health Statistics, Research and Evaluation, Mass. Dept. of Public Health

# Table 15A Cancer Incidence in Hingham, MA CT 5011.01 1982-1994

| 1982-1994 |     |      | Total |         |     |      | Male |         |     |      | Female |         |
|-----------|-----|------|-------|---------|-----|------|------|---------|-----|------|--------|---------|
|           | Obs | Exp  | SIR   | 95 % CI | Obs | Exp  | SIR  | 95 % CI | Obs | Exp  | SIR    | 95 % CI |
| Bladder   | 10  | 8.8  | 113   | 54 208  | 8   | 6.5  | 123  | 53 241  | 2   | 2.3  | NC     | NC      |
| Brain     | 3   | 3.0  | NC    | NC      | 2   | 1.7  | NC   | NC      | 1   | 1.4  | NC     | NC      |
| Kidney    | 1   | 4.9  | NC    | NC      | 1   | 3.0  | NC   | NC      | 0   | 1.9  | NC     | NC      |
| Leukemia  | 4   | 4.0  | NC    | NC      | 2   | 2.3  | NC   | NC      | 2   | 1.7  | NC     | NC      |
| Liver     | 0   | 1.0  | NC    | NC      | 0   | 0.8  | NC   | NC      | 0   | 0.3  | NC     | NC      |
| Lung      | 28  | 32.7 | 86    | 57 124  | 19  | 20.3 | 93   | 56 146  | 9   | 12.3 | 73     | 33 138  |
| NHL       | 6   | 7.2  | 83    | 30 181  | 3   | 3.8  | NC   | NC      | 3   | 3.4  | NC     | NC      |
| Pancreas  | 3   | 4.6  | NC    | NC      | 2   | 2.3  | NC   | NC      | 1   | 2.3  | NC     | NC      |

### Notes:

Expected number of cases presented are rounded to the nearest tenth.

SIRs are calculated based on the exact number of expected cases.

SIRs and 95% Confidence Interval are not calculated when fewer than 5 cases are observed.

Obs = Observed number of cases

Exp = Expected number of cases

SIR = Standardized Incidence Ratio

NC = Not Calculated

95% CI = 95% Confidence Interval

\* Indicates statistical significance (P<.05)

Data source: Massachusetts Cancer Registry, Bureau of Health Statistics, Research and Evaluation, Mass. Dept. of Public Health

## Table 15B Cancer Incidence in Hingham, MA CT 5011.01

1982-1986 & 1987-1994

| 1982-1986 |     |      | Total |         |     |     | Male |         |     |     | Female |         |
|-----------|-----|------|-------|---------|-----|-----|------|---------|-----|-----|--------|---------|
|           | Obs | Exp  | SIR   | 95 % CI | Obs | Exp | SIR  | 95 % CI | Obs | Exp | SIR    | 95 % CI |
| Bladder   | 3   | 3.4  | NC    | NC      | 1   | 2.5 | NC   | NC      | 2   | 0.9 | NC     | NC      |
| Brain     | 1   | 1.0  | NC    | NC      | 1   | 0.6 | NC   | NC      | 0   | 0.5 | NC     | NC      |
| Kidney    | 0   | 1.4  | NC    | NC      | 0   | 0.9 | NC   | NC      | 0   | 0.6 | NC     | NC      |
| Leukemia  | 1   | 1.5  | NC    | NC      | 1   | 0.8 | NC   | NC      | 0   | 0.7 | NC     | NC      |
| Liver     | 0   | 0.3  | NC    | NC      | 0   | 0.2 | NC   | NC      | 0   | 0.1 | NC     | NC      |
| Lung      | 11  | 11.3 | 97    | 48 174  | 8   | 7.4 | 108  | 46 213  | 3   | 3.9 | NC     | NC      |
| NHL       | 2   | 2.3  | NC    | NC      | 1   | 1.2 | NC   | NC      | 1   | 1.1 | NC     | NC      |
| Pancreas  | 1   | 1.7  | NC    | NC      | 0   | 0.8 | NC   | NC      | 1   | 0.9 | NC     | NC      |

| 1987-1994 |     |      | Total |         |     |      | Male |         |     |     | Female | 3       |
|-----------|-----|------|-------|---------|-----|------|------|---------|-----|-----|--------|---------|
|           | Obs | Exp  | SIR   | 95 % CI | Obs | Exp  | SIR  | 95 % CI | Obs | Exp | SIR    | 95 % CI |
| Bladder   | 7   | 5.5  | 128   | 51 264  | 7   | 4.1  | 173  | 69 356  | 0   | 1.4 | NC     | NC      |
| Brain     | 2   | 2.0  | NC    | NC      | 1   | 1.1  | NC   | NC      | 1   | 0.9 | NC     | NC      |
| Kidney    | 1   | 3.5  | NC    | NC      | 1   | 2.2  | NC   | NC      | 0   | 1.3 | NC     | NC      |
| Leukemia  | 3   | 2.5  | NC    | NC      | 1   | 1.4  | NC   | NC      | 2   | 1.0 | NC     | NC      |
| Liver     | 0   | 0.7  | NC    | NC      | 0   | 0.5  | NC   | NC      | 0   | 0.2 | NC     | NC      |
| Lung      | 17  | 21.4 | 79    | 46 127  | 11  | 13.0 | 85   | 42 152  | 6   | 8.5 | 71     | 26 155  |
| NHL       | 4   | 4.9  | NC    | NC      | 2   | 2.6  | NC   | NC      | 2   | 2.3 | NC     | NC      |
| Pancreas  | 2   | 2.9  | NC    | NC      | 2   | 1.4  | NC   | NC      | 0   | 1.4 | NC     | NC      |

### Notes:

Expected number of cases presented are rounded to the nearest tenth.

SIRs are calculated based on the exact number of expected cases.

SIRs and 95% Confidence Interval are not calculated when fewer than 5 cases are observed.

Obs = Observed number of cases

Exp = Expected number of cases

SIR = Standardized Incidence Ratio

NC = Not Calculated

95% CI = 95% Confidence Interval

\* Indicates statistical significance (P<.05)

Data source: Massachusetts Cancer Registry, Bureau of Health Statistics, Research and Evaluation, Mass. Dept. of Public Health

# Table 16A Cancer Incidence in Hingham, MA CT 5011.02 1982-1994

| 1982-1994 |     |      | Total |      |      |     |      | Male |         |     |      | Female | •         |
|-----------|-----|------|-------|------|------|-----|------|------|---------|-----|------|--------|-----------|
|           | Obs | Exp  | SIR   | 95   | % CI | Obs | Exp  | SIR  | 95 % CI | Obs | Exp  | SIR    | 95 % CI   |
| Bladder   | 7   | 16.0 | 44    | * 18 | 90   | 4   | 11.6 | NC   | NC      | 3   | 4.4  | NC     | NC        |
| Brain     | 10  | 5.9  | 170   | 81   | 312  | 4   | 3.2  | NC   | NC      | 6   | 2.7  | 220    | 81 480    |
| Kidney    | 5   | 9.1  | 55    | 18   | 128  | 5   | 5.5  | 91   | 29 212  | 0   | 3.6  | NC     | NC        |
| Leukemia  | 9   | 7.6  | 119   | 54   | 226  | 6   | 4.2  | 142  | 52 309  | 3   | 3.3  | NC     | NC        |
| Liver     | 1   | 1.9  | NC    |      | NC   | 0   | 1.4  | NC   | NC      | 1   | 0.6  | NC     | NC        |
| Lung      | 63  | 59.9 | 105   | 81   | 135  | 26  | 36.6 | 71   | 46 104  | 37  | 23.4 | 158    | * 112 218 |
| NHL       | 12  | 13.7 | 88    | 45   | 153  | 4   | 7.1  | NC   | NC      | 8   | 6.6  | 121    | 52 238    |
| Pancreas  | 6   | 8.4  | 71    | 26   | 155  | 1   | 4.0  | NC   | NC      | 5   | 4.4  | 113    | 37 265    |

### Notes:

Expected number of cases presented are rounded to the nearest tenth.

SIRs are calculated based on the exact number of expected cases.

SIRs and 95% Confidence Interval are not calculated when fewer than 5 cases are observed.

Obs = Observed number of cases

Exp = Expected number of cases

SIR = Standardized Incidence Ratio

NC = Not Calculated

95% CI = 95% Confidence Interval

\* Indicates statistical significance (P<.05)

Data source: Massachusetts Cancer Registry, Bureau of Health Statistics, Research and Evaluation, Mass. Dept. of Public Health

### Table 16B Cancer Incidence in Hingham, MA CT 5011.02

1982-1986 & 1987-1994

| 1982-1986 |     |      | Total |         |     |      | Male |         |     |     | Femal | e         |
|-----------|-----|------|-------|---------|-----|------|------|---------|-----|-----|-------|-----------|
|           | Obs | Exp  | SIR   | 95 % CI | Obs | Exp  | SIR  | 95 % CI | Obs | Exp | SIR   | 95 % CI   |
| Bladder   | 1   | 6.3  | NC    | NC      | 1   | 4.5  | NC   | NC      | 0   | 1.8 | NC    | NC        |
| Brain     | 5   | 2.0  | 249   | 80 581  | 2   | 1.1  | NC   | NC      | 3   | 0.9 | NC    | NC        |
| Kidney    | 2   | 2.7  | NC    | NC      | 2   | 1.6  | NC   | NC      | 0   | 1.1 | NC    | NC        |
| Leukemia  | 4   | 2.9  | NC    | NC      | 2   | 1.6  | NC   | NC      | 2   | 1.3 | NC    | NC        |
| Liver     | 0   | 0.6  | NC    | NC      | 0   | 0.4  | NC   | NC      | 0   | 0.2 | NC    | NC        |
| Lung      | 26  | 21.1 | 123   | 80 180  | 8   | 13.5 | 59   | 25 117  | 18  | 7.6 | 237   | * 140 374 |
| NHL       | 3   | 4.4  | NC    | NC      | 1   | 2.2  | NC   | NC      | 2   | 2.2 | NC    | NC        |
| Pancreas  | 5   | 3.3  | 153   | 49 358  | 1   | 1.5  | NC   | NC      | 4   | 1.7 | NC    | NC        |

| 1987-1994 |     |      | Total |         |     |      | Male |         |     |      | Female |         |
|-----------|-----|------|-------|---------|-----|------|------|---------|-----|------|--------|---------|
|           | Obs | Exp  | SIR   | 95 % CI | Obs | Exp  | SIR  | 95 % CI | Obs | Exp  | SIR    | 95 % CI |
| Bladder   | 6   | 9.7  | 62    | 23 135  | 3   | 7.0  | NC   | NC      | 3   | 2.6  | NC     | NC      |
| Brain     | 5   | 3.9  | 129   | 42 301  | 2   | 2.1  | NC   | NC      | 3   | 1.8  | NC     | NC      |
| Kidney    | 3   | 6.4  | NC    | NC      | 3   | 3.9  | NC   | NC      | 0   | 2.5  | NC     | NC      |
| Leukemia  | 5   | 4.7  | 107   | 34 249  | 4   | 2.7  | NC   | NC      | 1   | 2.0  | NC     | NC      |
| Liver     | 1   | 1.3  | NC    | NC      | 0   | 0.9  | NC   | NC      | 1   | 0.4  | NC     | NC      |
| Lung      | 37  | 38.7 | 96    | 67 132  | 18  | 23.0 | 78   | 46 124  | 19  | 15.8 | 120    | 73 188  |
| NHL       | 9   | 9.3  | 97    | 44 184  | 3   | 4.8  | NC   | NC      | 6   | 4.4  | 135    | 49 294  |
| Pancreas  | 1   | 5.2  | NC    | NC      | 0   | 2.5  | NC   | NC      | 1   | 2.7  | NC     | NC      |

### Notes:

Expected number of cases presented are rounded to the nearest tenth.

SIRs are calculated based on the exact number of expected cases.

SIRs and 95% Confidence Interval are not calculated when fewer than 5 cases are observed.

Obs = Observed number of cases

Exp = Expected number of cases

SIR = Standardized Incidence Ratio

NC = Not Calculated

95% CI = 95% Confidence Interval

\* Indicates statistical significance (P<.05)

Data source: Massachusetts Cancer Registry, Bureau of Health Statistics, Research and Evaluation, Mass. Dept. of Public Health

# Table 17A Cancer Incidence in Hingham, MA CT 5012 1982-1994

| 1982-1994 |     |      | Total |      |      |     |      | Male |         |     |      | Female |      |      |
|-----------|-----|------|-------|------|------|-----|------|------|---------|-----|------|--------|------|------|
|           | Obs | Exp  | SIR   | 95   | % CI | Obs | Exp  | SIR  | 95 % CI | Obs | Exp  | SIR    | 95 9 | % CI |
| Bladder   | 20  | 21.8 | 92    | 56   | 142  | 16  | 15.9 | 100  | 57 163  | 4   | 5.9  | NC     | N    | IC   |
| Brain     | 7   | 8.2  | 85    | 34   | 176  | 2   | 4.6  | NC   | NC      | 5   | 3.6  | 138    | 44   | 321  |
| Kidney    | 11  | 12.6 | 87    | 43   | 156  | 6   | 7.9  | 76   | 28 166  | 5   | 4.7  | 105    | 34   | 246  |
| Leukemia  | 2   | 10.5 | NC    |      | NC   | 0   | 6.0  | NC   | NC      | 2   | 4.5  | NC     | N    | IC   |
| Liver     | 1   | 2.7  | NC    |      | NC   | 1   | 1.9  | NC   | NC      | 0   | 0.7  | NC     | N    | IC   |
| Lung      | 54  | 82.3 | 66    | * 49 | 86   | 30  | 51.6 | 58   | * 39 83 | 24  | 30.7 | 78     | 50   | 116  |
| NHL       | 21  | 18.6 | 113   | 70   | 172  | 13  | 9.9  | 131  | 70 225  | 8   | 8.7  | 92     | 39   | 181  |
| Pancreas  | 5   | 11.4 | 44    | 14   | 102  | 2   | 5.6  | NC   | NC      | 3   | 5.8  | NC     | N    | IC   |

### Notes:

Expected number of cases presented are rounded to the nearest tenth.

SIRs are calculated based on the exact number of expected cases.

SIRs and 95% Confidence Interval are not calculated when fewer than 5 cases are observed.

Obs = Observed number of cases

Exp = Expected number of cases

SIR = Standardized Incidence Ratio

NC = Not Calculated

95% CI = 95% Confidence Interval

\* Indicates statistical significance (P<.05)

Data source: Massachusetts Cancer Registry, Bureau of Health Statistics, Research and Evaluation, Mass. Dept. of Public Health

### Table 17B Cancer Incidence in Hingham, MA CT 5012

1982-1986 & 1987-1994

| 1982-1986 |     |      | Total |         |     |      | Male |         |     |     | Female |         |
|-----------|-----|------|-------|---------|-----|------|------|---------|-----|-----|--------|---------|
|           | Obs | Exp  | SIR   | 95 % CI | Obs | Exp  | SIR  | 95 % CI | Obs | Exp | SIR    | 95 % CI |
| Bladder   | 7   | 8.1  | 86    | 34 177  | 7   | 5.8  | 120  | 48 247  | 0   | 2.3 | NC     | NC      |
| Brain     | 2   | 2.7  | NC    | NC      | 1   | 1.5  | NC   | NC      | 1   | 1.2 | NC     | NC      |
| Kidney    | 0   | 3.6  | NC    | NC      | 0   | 2.2  | NC   | NC      | 0   | 1.4 | NC     | NC      |
| Leukemia  | 1   | 3.9  | NC    | NC      | 0   | 2.1  | NC   | NC      | 1   | 1.7 | NC     | NC      |
| Liver     | 0   | 0.8  | NC    | NC      | 0   | 0.6  | NC   | NC      | 0   | 0.3 | NC     | NC      |
| Lung      | 20  | 27.7 | 72    | 44 112  | 10  | 18.1 | 55   | 26 102  | 10  | 9.6 | 104    | 50 191  |
| NHL       | 6   | 5.8  | 104   | 38 226  | 3   | 3.0  | NC   | NC      | 3   | 2.8 | NC     | NC      |
| Pancreas  | 0   | 4.2  | NC    | NC      | 0   | 2.0  | NC   | NC      | 0   | 2.2 | NC     | NC      |

| 1987-1994 |     |      | Total |      |      |     |      | Male | •    |     |     |      | Female |    |      |
|-----------|-----|------|-------|------|------|-----|------|------|------|-----|-----|------|--------|----|------|
|           | Obs | Exp  | SIR   | 95   | % CI | Obs | Exp  | SIR  | 95 % | CI  | Obs | Exp  | SIR    | 95 | % CI |
| Bladder   | 13  | 13.5 | 96    | 51   | 164  | 9   | 10.0 | 90   | 41 1 | 172 | 4   | 3.6  | NC     | ]  | NC   |
| Brain     | 5   | 5.5  | 92    | 30   | 214  | 1   | 3.0  | NC   | NC   |     | 4   | 2.4  | NC     | ]  | NC   |
| Kidney    | 11  | 9.0  | 122   | 61   | 218  | 6   | 5.7  | 105  | 38 2 | 229 | 5   | 3.3  | 150    | 48 | 351  |
| Leukemia  | 1   | 6.6  | NC    | ]    | NC   | 0   | 3.8  | NC   | NC   |     | 1   | 2.7  | NC     | ]  | NC   |
| Liver     | 1   | 1.8  | NC    | ]    | NC   | 1   | 1.4  | NC   | NC   |     | 0   | 0.5  | NC     | ]  | NC   |
| Lung      | 34  | 54.3 | 63    | * 43 | 88   | 20  | 33.2 | 60   | * 37 | 93  | 14  | 21.0 | 67     | 36 | 112  |
| NHL       | 15  | 12.8 | 117   | 65   | 193  | 10  | 6.9  | 145  | 70 2 | 267 | 5   | 5.9  | 84     | 27 | 197  |
| Pancreas  | 5   | 7.2  | 70    | 23   | 163  | 2   | 3.6  | NC   | NC   |     | 3   | 3.5  | NC     | ]  | NC   |

### Notes:

Expected number of cases presented are rounded to the nearest tenth.

SIRs are calculated based on the exact number of expected cases.

SIRs and 95% Confidence Interval are not calculated when fewer than 5 cases are observed.

Obs = Observed number of cases

Exp = Expected number of cases

SIR = Standardized Incidence Ratio

NC = Not Calculated

95% CI = 95% Confidence Interval

\* Indicates statistical significance (P<.05)

Data source: Massachusetts Cancer Registry, Bureau of Health Statistics, Research and Evaluation, Mass. Dept. of Public Health