



**Massachusetts Division of Marine Fisheries
Technical Report TR-7**

Technical Report

Massachusetts Coastal Commercial Lobster Trap Sampling Program May—November, 2000


B. T. Estrella and R. P. Glenn

**Massachusetts Division of Marine Fisheries
Department of Fisheries, Wildlife and Environmental Law Enforcement
Executive Office of Environmental Affairs
Commonwealth of Massachusetts**

July 2001



Massachusetts Division of Marine Fisheries
Technical Report TR-7



Massachusetts Coastal Commercial Lobster Trap Sampling Program May-November, 2000

Bruce T. Estrella and Robert P. Glenn

Massachusetts Division of Marine Fisheries
South Shore Field Station
50A Portside Drive
Pocasset, MA 02559

July 2001

Massachusetts Division of Marine Fisheries
Paul Diodati, Director
Department of Fisheries, Wildlife and Environmental Law Enforcement
Dave Peters, Commissioner
Executive Office of Environmental Affairs
Bob Durand, Secretary
Commonwealth of Massachusetts
Jane Swift, Governor

TABLE OF CONTENTS

ABSTRACT	ii
INTRODUCTION	1
STUDY AREA	1
SAMPLING PROCEDURE	1
ANALYTICAL PROCEDURES	3
RESULTS AND DISCUSSION	4
Commercial Lobster Sampling	4
Water Temperature Time Series	7
ACKNOWLEDGEMENTS	11
LITERATURE CITED	12
APPENDIX	13

ABSTRACT

This is the Massachusetts Division of Marine Fisheries twentieth annual assessment of the status of the American lobster resource in Massachusetts coastal waters. During the period of May through November, 2000, eighty-three (83) sampling trips were made aboard commercial lobster vessels. A total of 38,390 lobster was sampled from 17,251 trap hauls. The catch rate of marketable lobster, 0.885 lobster per trap, was 2% lower than the 1999 index, 0.902. The proportion of females ovigerous, 15.4%, was lower than in the previous year (17.4%). The coastwide fishing mortality estimate, 1.38, was higher than the 1999 index of 1.22. Exploitation rate, 0.68, increased fractionally, while mean carapace length of marketable lobster, 89.4 mm, and mean size of egg-bearing females, 88.1 mm, were similar and remained unchanged, respectively. The cull rate, 18.2%, decreased about 2% from the 1999 estimate of 20.8%. Less than 1% of the lobster sampled from traps were dead.

A time series of data from our bottom water temperature monitoring program is presented for seven locations in Buzzards Bay, Cape Cod Bay, and Massachusetts Bay for the period 1985-2000. The locations of three shallow water sites (<20'), added during summer 2000, are depicted.

INTRODUCTION

This is the Massachusetts Division of Marine Fisheries (DMF) nineteenth annual assessment of the status of the American lobster resource in Massachusetts coastal waters. Since the lobster resource supports the most economically important single-species fishery in Massachusetts coastal waters, a long-term coastwide lobster monitoring program yielding biological and catch per unit effort data was devised and initiated in Massachusetts in May, 1981. A sea sampling/survey design was chosen by which both catch per unit effort and biological data could be collected temporally and areally with sufficient precision for stock assessments. The objective was to assess variations in population parameters due to environmental factors, fishing pressure, and regulatory changes.

Data collected during the 2000 coastwide commercial lobster trap sampling program are summarized below. Parameter trends occurring during the 1981-2000 study period are presented.

STUDY AREA

The study area is primarily defined by the Massachusetts territorial sea, except where lobstering activities of cooperating commercial lobstermen exceeded territorial boundaries (Figure 1). Territorial waters total 5,322 sq km (2,055 sq n mi), of which an estimated 60% is considered major lobster habitat. Six sampling regions, Cape Ann, Beverly-Salem, Boston Harbor, Cape Cod Bay, outer Cape Cod, and Buzzards Bay, were chosen for coverage of the major lobstering regions of the state. For convenience, these regions are depicted in Figure 1 as generalized hatch-marked areas wherein lobster gear sampled may be discontinuously distributed.

SAMPLING PROCEDURE

Sampling of coastal waters was accomplished by monitoring catches during the normal lobstering operations of volunteer commercial lobstermen in each designated region. Multiple lobstering operations were observed to reduce bias from varying degrees of lobstering skill and to enhance areal coverage. Pot-sampling trips were day trips, conducted a minimum of once per month per region during the major lobstering season, May-November.

Utilizing portable cassette tape recorders, sea samplers recorded carapace length (to the nearest mm); sex; and condition, including the degree of shell hardness, culls and other shell damage, external gross pathology, mortality, and presence of extruded ova on females (ovigerous). Catch in number of lobster, number of trap hauls, set-over-days, trap and bait type were also recorded. Trap locations were recorded from LORAN and plotted on nautical charts. Depth information was then acquired from the charts as a coastwide standard to avoid variability from tidal fluctuations.




Figure 1. Map of Massachusetts with six sampling regions and trap/trawl locations sampled during 2000.

ANALYTICAL PROCEDURES

Data were computer coded and keypunched with a microcomputer data entry program. The data base was subsequently transferred for analysis to the Massachusetts Executive Office of Environmental Affairs' (EOEA) Digital Equipment Corporation VAX-11/780 computer system. A computer auditing process was used to uncover keypunch and recording errors and statistical analyses were performed with SPSS (Nie 1983) statistical sub-programs.

Because parameter means exhibit significant regional and monthly variation, an areal and temporal data weighting scheme was incorporated into analytical software. As a result, each month's data contribute equally to regional parameter means which are weighted by area in square nautical miles to generate coastwide means.

Unless specified otherwise, the terms "legal" or "legal sized" lobster include all lobster in the carapace length category ≥ 82.6 mm. The marketable segment of this category, which excludes ovigerous females, is analyzed separately and referred to as "marketable lobster". The sublegal length category includes all lobster < 82.6 mm.

The catch rates of marketable lobster are expressed as CTH'_3 . This is catch per trap haul standardized to 3 set-over-days (Estrella and McKiernan 1989).

Estimates of total instantaneous mortality (Z) and total annual mortality ($A=1-e^{-Z}$) were computed by two methods which produce extremes in the possible range of estimates. The method of Gulland (1969) requires computation of the regression line slope of natural log-transformed numbers at estimated age (15% molt groups, 14% for Buzzards Bay, were derived from tagging data). Beverton and Holt's (1956) process employs von Bertalanffy Growth Equation parameters (from Fair 1977) and mean and minimum length of exploitable sizes.

Estimates of fishing mortality (F) were calculated with cohort analysis (Pope 1972, Jones 1974). Rates of exploitation were calculated with the equation $u=FA/Z$, where F = fishing mortality, A = total annual mortality, and Z = total instantaneous mortality.

Lobster landings data were derived from lobstermen's catch reports which are compiled annually by the DMF Commercial Fisheries Statistics Project.

Since current management strategy stresses uniform coastwide regulations, all data are grouped for a coastwide analysis. However, the uniqueness of the Massachusetts coastline, its role as a temperature barrier which profoundly affects many marine species (Colton 1964), and the influence of offshore lobster stocks on the inshore resource mandate a regional data treatment as well.

RESULTS AND DISCUSSION

Commercial Lobster Sampling

During the period of May through November, 2000, eighty-three (83) sampling trips were made aboard commercial lobster vessels in Massachusetts coastal waters. A total of 38,390 lobster was sampled from 17,251 trap hauls.

The 2000 coastwide mean catch per unit effort index (CTH₃), 0.885 marketable lobster per trap, was only 2% lower than the 1999 index, 0.902 (Appendix Table 1). Total Massachusetts commercial landings, 14,853,390 lbs, decreased by 7% from 1999. Landings from territorial waters, 9,865,733 lbs, increased by 3%. Landings and catch rate trends are depicted in Figure 2. The coastwide mean catch rates of sublegal lobster increased by 19% (CTHSOD) and 11% (CTHAUL) from 1999 (Appendix Tables 2 and 3).




Figure 2. Catch per unit effort of marketable American lobster from commercial trap sampling and Massachusetts lobster landings from territorial water, 1981 - 2000.




Figure 3. Relative abundance of ovigerous female American lobster in percent total females and catch per effort, Massachusetts coastal waters, 1981 - 2000.

Of all females sampled during 2000, 15.4% were ovigerous compared to 17.4% in 1999 (Appendix Table 4). Trends in statewide CPUE of ovigerous females (Appendix Tables 5-6) either decreased or remained unchanged and are depicted in Figure 3.

Approximately 92% of the legal catch in our inshore regions (Cape Ann south through Cape Cod Bay and Buzzards Bay) was comprised of new recruits (83 mm-94 mm CL), i.e., lobster which recruited to the legal size range during their most recent molt (Appendix Table 7). This index of the effect of fishing pressure on the size frequency was similar to 1999. The index fluctuated from 46% in 1999 to 51% in 2000 for the primarily offshore migrant lobster sampled east of Cape Cod. Estimates of total mortality (Z) for inshore Gulf of Maine regions ($Z = 1.33$ - 3.14 , $A = 74\%$ - 96%) and Buzzards Bay ($Z = 2.38$ - 2.70 , $A = 91\%$ - 93%) depict a heavily exploited resource while those for the outer Cape Cod region ($Z = 0.66$ - 0.82 , $A = 44\%$ - 45%) indicate that a lower level of fishing pressure was exerted on this lobster group (Appendix Tables 8A and 8B).

Estimates of instantaneous fishing mortality (F), the proportion of all deaths which are attributed to fishing, ranged from 0.63 off outer Cape Cod to 2.21 in Boston Harbor (Appendix Table 9A). Estimates of F using DeltaT values in place of von Bertalanffy growth parameters were lower (Table 9B), ranging from 0.39 off outer Cape Cod to 1.07 in Buzzards Bay. Exploitation rates (u), i.e. the fraction of the population that is removed by fishing, was relatively unchanged from 1999 data, fluctuating from 0.67 to 0.68 in 2000 (Appendix Table 10).

The relationship between fishing mortality, rate of exploitation, and mean lobster size is depicted in Figure 4. Carapace length exhibited a downward trend as fishing mortality and exploitation rates increased through 1987. Thereafter, increases in mean carapace length of 0.7 mm occurred in 1988 (mean size = 88.2 mm) and 1989 (mean size = 88.9 mm, Appendix Table 11) which reflected the similar numerical change in the minimum legal size during those years.

Carapace length then fluctuated downward until 1994, and increased during 1995 – 1997, reaching a time series high in 1999. The 2000 mean was similar (89.4mm). Fishing mortality rates for all regions combined edged upward to a time-series high of 1.48 in 1993, declined in 1994-1996 along with exploitation rates, then fluctuated upward in 1997-1998, before declining to a 1982 level in 1999. The 2000 estimate increased significantly. The relative change in size frequency between 1999 and 2000 is depicted by the overlay in Figure 5.

Sublegal sized lobster averaged 77.1 mm carapace length during 2000 compared to 77.4 mm during 1999 (Appendix Table 12). The mean size of all ovigerous females was static between 1999 and 2000 at 88.1 mm.




Figure 4. Relationship between exploitation rate, fishing mortality, and mean carapace length of marketable American lobster, Massachusetts coastal water, 1981 - 2000.

The percentage of culls (lobster with one or both claws missing or regenerating) among all lobster sampled decreased approximately 2% between 1999 (20.8%) and 2000 (18.2%) (Appendix Table 14). The cull rates for legal, marketable, and sublegal size groups also decreased comparably between years (Appendix Tables 15-17).

The coastwide incidence of lobster found dead in traps was 0.18%. This was unchanged from that of the previous year (Appendix Table 18) and is acceptably low.




Figure 5. Length frequencies of trap-caught American lobster, Massachusetts coastal waters, 1999 - 2000.

Water Temperature Time Series

In 1985, a coastal bottom water temperature monitoring project was initiated. Temperature monitors (RYAN Tempmentor) have been deployed for various lengths of time at several sites in Cape Cod Bay, outside Boston Harbor, and Buzzards Bay (Figure 6). Some of these sites are located on ship wrecks.

The longest time series of bottom temperatures is from Cleveland Light in Buzzards Bay. The last monitor to be deployed was at Rocky Point, off Plymouth. The Rocky Point, Manomet Point, *Endicott*, and *Mars* sites represent the 0-30 ft., 30-60 ft., 61-90 ft., and 91-120 ft. depth strata, respectively, in Cape Cod Bay. The Martin's Ledge (formerly *Romance*, off Boston Harbor), and Buzzards Bay-South (Barge) sites are located at 70-80 ft. and provide data from the north-south extremes in our series. The Cleveland Light monitor is located in 30 feet of water.

Monitors are retrieved and replaced annually by divers. Although the time series contained data from seven monitors at one point, we have collected data from only six sites since 10/5/91 when the monitor at the *Endicott* site was lost and has not been replaced. Figures 7 and 8 present the bottom water temperature at sites in Buzzards Bay and Cape Cod Bay/Massachusetts Bay, respectively. Figure 9 provides a comparison between the annual mean bottom temperature at Cleveland Light, Buzzards Bay-south, Manomet, Rocky Point, *Mars*, and Martin's Ledge, and the annual mean surface temperature at Boston and Woods Hole provided by NOAA/NOS.

Three sites (<20 ') were added in summer, 2001 (Onset Stowaway XTI monitors). These are site #'s 2, 6, and 8 in Buzzards Bay, Cape Cod Bay, and Boston Harbor (Figure 6).




Figure 6. Map of Massachusetts with locations of nine bottom temperature monitors.




Figure 7. Mean monthly bottom water temperatures at two sites in Buzzards Bay, 1986-2001.




Figure 8. Mean monthly bottom water temperatures at five sites in the Gulf of Maine, 1988-2001.




Figure 9. Mean annual bottom water temperatures at four sites monitored by the Coastal Lobster Project and mean annual surface temperature at two sites monitored by NOAA/NOS, 1986 - 2000.

ACKNOWLEDGEMENTS

We are indebted to the many commercial lobstermen whose cooperative spirit and concern for the American lobster resource sustain our lobster monitoring program. Gratitude is also extended to Bill Hoffman, Brad Chase, Ross Kessler, Jeff Plough, and Brian Kelly for data collection, Ann Spires, for data entry, and James Fair who administered the project and reviewed the manuscript. Rick Crawford provided assistance with GIS mapping. We also thank Thomas Hoopes for his data entry software design and assistance in data quality control.

LITERATURE CITED

- Beverton, R. J. H. and S. J. Holt. 1956. A review of methods for estimating mortality rates in exploited populations, with special reference to sources of bias in catch sampling. Reports et Proces-Verbaux des Reunions 140:67-83.
- Caputi, N., and R. S. Brown. 1986. Relationship between indices of juvenile abundance and recruitment in the western rock lobster (*Panulirus cygnus*) fishery. Can. J. Fish. Aquat. Sci. 43:2131-2139.
- Colton, J. B. 1964. History of oceanography in the offshore waters of the Gulf of Maine. U.S. Fish and Wildlife Service, Special Scientific Report, Fisheries No. 496. 18p.
- Estrella, B. T. and D. J. McKiernan. 1989. Catch-per-unit-effort and biological parameters from the Massachusetts coastal lobster (*Homarus americanus*) resource: description and trends. NOAA Tech. Rep. NMFS 81. 21 p.
- Fair, J. J., Jr. 1977. Lobster investigations in management area I: Southern Gulf of Maine. NOAA, NMFS State-Federal Relationships Division, Mass. Lobster Report No. 8., April 21, 1975 - April 20, 1977, 8 p. and Appendix, 14 p.
- Gulland, J. A. 1969. Manual of methods for fish stock assessment. Pt. 1, fish population analysis. FAO Manuals in Fisheries Science 4, 154 p.
- Jones, R. 1974. Assessing the long term effects of changes in fishing effort and mesh size from length composition data. ICES CM 1974/F:33.
- Nie, N. H. 1983. SPSS: statistical package for the social sciences, McGraw-Hill, New York, 806 p.
- Pope, J. G. 1972. An investigation of the accuracy of virtual population analysis using Cohort analysis. Research Bulletin Int. Comm. NW Atlant. Fish. 9:65-74.
- Sokal, R.R. and F.J. Rohlf. 1969. Biometry. W.H. Freeman and Company, San Francisco. 776 p.

APPENDIX

Table 1. CTH'3, by state and region, for all marketable lobster sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-2000.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
State	0.767	0.785	0.803	0.696	0.825	0.816	0.737	0.820	0.751	0.826	0.795	0.716	0.671	0.966	0.893	0.792	0.776	0.662	0.902	0.885
Cape Ann	0.732	0.808	0.624	0.663	0.634	0.699	0.669	0.496	0.721	0.904	0.868	0.724	0.770	1.015	0.979	1.272	1.106	0.645	0.699	0.930
Beverly-Salem	0.934	0.898	0.881	0.835	0.663	0.496	0.611	0.661	0.639	0.827	0.586	0.390	0.509	0.898	0.840	0.788	0.419	0.767	1.005	0.945
Boston Harbor	---	---	---	1.108	1.254	1.096	1.058	1.057	1.123	1.224	1.160	0.734	0.750	0.725	0.626	0.534	0.638	0.462	0.787	0.629
Cape Cod Bay	0.710	0.776	0.680	0.479	0.716	0.822	0.533	0.752	0.539	0.630	0.693	0.567	0.494	1.052	0.906	0.662	0.760	0.633	1.021	0.888
Outer Cape Cod	0.808	0.824	0.765	0.598	0.856	0.811	0.937	0.861	0.923	1.219	1.148	1.339	1.021	1.105	1.117	1.027	0.837	0.785	0.893	1.042
Buzzards Bay	0.611	0.571	1.110	0.870	0.953	0.907	0.952	1.064	0.934	0.598	0.575	0.817	0.834	0.852	0.893	0.866	1.169	0.719	0.740	0.882

Table 2. CTHSOD, by state and region, for all sub-legal American lobster, sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-2000.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
State	0.580	0.672	0.718	0.521	0.647	0.700	0.578	0.509	0.695	0.716	0.665	0.465	0.542	0.402	0.410	0.343	0.363	0.295	0.215	0.255
Cape Ann	0.067	0.109	0.586	0.450	0.395	0.474	0.417	0.388	0.670	0.589	0.728	0.726	0.447	0.658	0.800	0.573	0.831	0.356	0.171	0.222
Beverly-Salem	0.708	0.711	1.263	0.948	0.833	0.801	0.863	0.353	0.780	0.408	0.324	0.411	0.406	0.314	0.334	0.277	0.231	0.364	0.233	0.254
Boston Harbor	---	---	---	0.901	1.162	1.138	1.156	0.639	0.966	1.103	0.924	0.839	0.800	0.690	0.782	0.688	0.651	0.754	0.508	0.620
Cape Cod Bay	0.710	1.013	0.639	0.322	0.594	0.551	0.371	0.438	0.595	0.727	0.716	0.298	0.436	0.313	0.307	0.217	0.256	0.204	0.162	0.256
Outer Cape Cod	0.037	0.024	0.038	0.033	0.035	0.027	0.088	0.064	0.066	0.078	0.077	0.088	0.075	0.045	0.060	0.051	0.053	0.043	0.025	0.026
Buzzards Bay	0.787	0.620	0.638	0.785	0.848	1.312	0.871	1.153	1.188	1.236	1.072	0.784	1.156	0.649	0.492	0.540	0.510	0.254	0.272	0.158

Table 3. CTHAUL, by state and region, for all sub-legal American lobster, sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-2000.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
State	1.473	1.401	1.624	1.389	1.705	1.899	1.873	1.736	2.297	2.216	1.996	1.460	1.720	1.389	1.457	1.332	1.417	1.259	0.967	1.071
Cape Ann	0.256	0.199	1.044	0.909	1.031	1.126	1.143	1.062	1.765	1.782	1.783	1.661	1.562	1.725	2.323	1.660	1.940	1.074	0.606	0.776
Beverly-Salem	1.855	1.713	2.526	2.504	2.567	2.435	3.482	1.862	3.477	1.867	1.563	1.502	1.540	1.717	1.920	1.654	1.988	2.038	1.577	1.759
Boston Harbor	---	---	---	2.773	3.038	3.314	3.334	1.959	3.104	3.382	2.451	2.069	2.284	2.189	2.390	2.511	2.258	2.864	1.660	2.286
Cape Cod Bay	1.544	1.680	1.345	0.825	1.337	1.512	1.031	1.442	1.742	1.921	2.086	1.065	1.334	1.033	1.102	0.873	0.926	0.701	0.680	0.857
Outer Cape Cod	0.233	0.145	0.210	0.189	0.160	0.161	0.324	0.353	0.306	0.453	0.452	0.490	0.474	0.288	0.359	0.372	0.354	0.301	0.172	0.163
Buzzards Bay	2.381	1.916	2.316	1.965	2.452	3.118	3.090	3.722	3.984	3.994	3.181	2.602	3.501	2.179	1.599	1.835	2.051	1.546	1.456	0.927

Table 4. Percent of females ovigerous, by state and region, for all American lobster sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-2000.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
State	5.9	7.7	10.9	9.1	8.6	9.1	9.2	8.8	10.0	10.9	9.8	11.9	14.3	10.7	14.1	15.4	14.7	16.4	17.4	15.4
Cape Ann	1.7	3.1	4.4	3.2	4.6	5.0	4.5	3.5	6.3	6.9	4.3	6.7	9.3	4.7	5.3	6.4	8.3	9.3	10.4	8.0
Beverly-Salem	1.7	2.8	1.2	0.4	1.9	1.1	1.8	1.5	1.6	1.8	3.2	3.9	5.4	2.3	6.3	6.9	8.3	6.1	5.9	4.0
Boston Harbor	---	---	---	1.4	1.2	2.0	1.7	2.0	2.1	2.7	2.8	3.0	4.4	4.7	5.0	6.7	6.7	8.0	7.4	10.4
Cape Cod Bay	3.9	3.1	3.7	3.1	3.2	2.1	3.9	2.9	3.0	3.3	5.4	6.8	6.8	7.4	10.2	13.6	11.9	13.1	15.5	11.9
Outer Cape Cod	11.1	23.0	30.3	26.8	22.3	28.9	16.9	21.4	27.4	24.5	18.3	27.7	26.8	27.3	34.4	34.6	32.9	30.3	34.6	33.4
Buzzards Bay	16.0	16.9	32.5	26.6	25.0	25.3	31.0	27.8	29.2	35.0	28.2	28.8	40.9	22.1	26.0	23.6	22.2	32.9	29.5	27.0

Table 5. CTHSOD, by state and region, for all ovigerous female American lobster sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-2000.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
State	0.024	0.027	0.050	0.038	0.044	0.057	0.049	0.054	0.057	0.073	0.032	0.054	0.092	0.036	0.045	0.047	0.042	0.039	0.039	0.039
Cape Ann	0.002	0.011	0.024	0.015	0.016	0.017	0.016	0.010	0.037	0.035	0.024	0.050	0.038	0.024	0.030	0.031	0.056	0.037	0.026	0.016
Beverly-Salem	0.011	0.009	0.008	0.003	0.011	0.004	0.010	0.004	0.009	0.005	0.008	0.014	0.020	0.008	0.017	0.015	0.013	0.016	0.012	0.080
Boston Harbor	---	---	---	0.009	0.007	0.015	0.012	0.012	0.010	0.028	0.017	0.017	0.026	0.024	0.028	0.032	0.035	0.046	0.032	0.049
Cape Cod Bay	0.020	0.025	0.016	0.009	0.015	0.010	0.012	0.009	0.014	0.017	0.028	0.016	0.023	0.022	0.034	0.035	0.027	0.032	0.043	0.038
Outer Cape Cod	0.012	0.028	0.040	0.030	0.038	0.032	0.034	0.030	0.043	0.055	0.038	0.076	0.053	0.046	0.081	0.085	0.062	0.045	0.049	0.050
Buzzards Bay	0.079	0.053	0.230	0.183	0.193	0.297	0.234	0.289	0.270	0.349	0.073	0.197	0.446	0.110	0.088	0.098	0.083	0.067	0.060	0.060

Table 6. CTHAUL, by state and region, for all ovigerous female American lobster sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-2000.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
State	0.073	0.078	0.179	0.116	0.133	0.167	0.183	0.189	0.211	0.282	0.140	0.227	0.319	0.143	0.200	0.226	0.219	0.205	0.230	0.190
Cape Ann	0.010	0.016	0.038	0.027	0.039	0.047	0.048	0.031	0.096	0.109	0.056	0.088	0.135	0.064	0.085	0.081	0.124	0.097	0.096	0.066
Beverly-Salem	0.025	0.033	0.016	0.006	0.033	0.018	0.036	0.021	0.039	0.023	0.049	0.047	0.067	0.048	0.106	0.103	0.136	0.101	0.094	0.055
Boston Harbor	---	---	---	0.030	0.025	0.050	0.037	0.038	0.043	0.075	0.064	0.046	0.081	0.088	0.100	0.124	0.132	0.181	0.110	0.190
Cape Cod Bay	0.048	0.048	0.040	0.024	0.040	0.031	0.038	0.034	0.039	0.055	0.091	0.056	0.078	0.075	0.128	0.147	0.130	0.110	0.210	0.140
Outer Cape Cod	0.081	0.178	0.242	0.170	0.176	0.225	0.157	0.198	0.258	0.342	0.251	0.453	0.317	0.306	0.514	0.561	0.408	0.307	0.330	0.340
Buzzards Bay	0.243	0.139	0.828	0.515	0.555	0.748	0.889	0.929	0.953	1.291	0.359	0.847	1.438	0.383	0.343	0.414	0.477	0.527	0.450	0.390

Table 7. Estimated fishing pressure index, by state and region, commercial lobster trap catch survey, Massachusetts coastal water, 1981-2000.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
State	86	87	86	86	88	88	89	90	88	87	89	89	90	88	86	86	86	88	88	86
Cape Ann	91	92	87	89	87	87	88	90	84	81	90	87	83	86	88	85	88	85	80	83
Beverly-Salem	89	92	94	88	96	96	97	98	96	95	97	98	96	96	95	95	96	94	97	95
Boston Harbor	---	---	---	93	94	96	96	96	96	95	96	95	96	95	94	95	94	96	97	98
Cape Cod Bay	90	93	92	94	93	94	92	94	94	93	91	92	94	90	86	88	87	90	87	88
Outer Cape Cod	46	43	42	38	48	46	54	57	47	50	54	57	60	60	55	54	57	58	46	51
Buzzards Bay	98	96	96	94	96	97	97	97	95	94	95	97	97	98	96	95	92	96	96	95

Table 8A. Total instantaneous (Z)* and total annual (A)** mortality estimates (Gulland, 1969) of American lobster by state and region, Massachusetts coastal waters, 1981-2000.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
State	1.58	* 1.72	1.66	1.66	1.76	1.80	1.90	1.86	1.80	1.76	1.90	1.92	1.86	2.02	1.83	1.79	1.87	1.81	1.81	1.92
	79%	** 82%	81%	81%	83%	84%	85%	84%	83%	83%	85%	85%	84%	87%	84%	83%	85%	84%	84%	85%
Cape Ann	1.65	2.18	1.72	1.92	1.94	2.03	1.85	1.75	1.55	1.39	1.97	1.87	1.51	1.81	1.95	1.90	2.00	1.65	1.48	1.95
	81%	89%	82%	85%	86%	87%	84%	83%	79%	75%	86%	85%	78%	84%	86%	85%	86%	81%	77%	85%
Beverly-Salem	1.97	2.15	2.41	2.71	3.64	3.60	3.49	3.31	3.59	2.81	3.49	3.12	2.62	3.34	3.10	2.90	2.46	2.81	3.14	2.64
	86%	88%	91%	93%	97%	97%	97%	96%	97%	94%	97%	96%	93%	96%	95%	94%	91%	94%	96%	93%
Boston Harbor	---	---	---	2.52	3.59	2.60	2.77	2.86	2.96	3.00	3.40	3.54	3.26	3.21	2.87	2.65	2.77	2.59	2.86	3.14
	---	---	---	92%	97%	93%	94%	94%	95%	95%	97%	97%	96%	96%	94%	93%	94%	92%	94%	96%
Cape Cod Bay	2.53	2.69	2.42	2.52	2.31	2.83	2.26	2.74	2.43	2.46	2.33	2.58	2.60	3.10	2.35	2.09	2.14	2.02	2.14	2.73
	92%	93%	91%	92%	90%	94%	90%	94%	91%	91%	90%	92%	93%	95%	90%	88%	88%	87%	88%	93%
Outer Cape Cod	0.43	0.46	0.42	0.33	0.52	0.51	0.80	0.71	0.62	0.63	0.77	0.78	0.87	0.92	0.74	0.73	0.89	0.86	0.58	0.82
	35%	37%	34%	28%	41%	40%	55%	51%	46%	47%	54%	54%	58%	60%	52%	52%	59%	58%	44%	56%
Buzzards Bay	3.02	3.00	8.64	3.14	3.55	3.71	3.48	3.18	3.13	2.60	3.50	3.81	3.03	3.58	3.34	2.84	2.74	3.27	3.81	2.70
	95%	95%	99%	96%	97%	98%	97%	96%	96%	93%	97%	98%	95%	97%	96%	94%	94%	96%	98%	93%

Table 8B. Total instantaneous (Z)* and total annual (A)** mortality estimates (Beverton and Holt, 1956) of American lobster by state and region, Massachusetts coastal waters, 1981-2000.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
State	1.35	*1.45	1.39	1.41	1.47	1.49	1.54	1.56	1.53	1.50	1.73	1.70	1.79	1.67	1.57	1.54	1.55	1.62	1.62	1.53
	74%	**77%	75%	76%	77%	78%	79%	79%	78%	78%	82%	82%	83%	81%	79%	79%	79%	80%	80%	78%
Cape Ann	1.32	1.39	1.35	1.52	1.33	1.32	1.39	1.51	1.27	1.66	1.77	1.57	1.38	1.48	1.62	1.49	1.67	1.52	1.14	1.33
	73%	75%	74%	78%	74%	73%	75%	78%	72%	81%	83%	79%	75%	77%	80%	77%	81%	78%	68%	74%
Beverly-Salem	1.59	1.70	1.85	1.78	1.96	1.99	2.16	1.98	2.01	1.83	2.29	2.50	2.23	2.18	2.09	2.11	2.28	2.18	2.08	2.18
	80%	82%	84%	83%	86%	86%	88%	86%	87%	84%	90%	92%	89%	89%	88%	88%	90%	89%	88%	89%
Boston Harbor	---	---	---	1.82	1.75	1.92	1.88	1.84	1.94	1.87	2.19	2.14	2.33	2.28	2.09	2.18	2.09	2.63	2.12	2.73
	---	---	---	84%	83%	85%	85%	84%	86%	85%	89%	88%	90%	90%	88%	89%	88%	93%	88%	94%
Cape Cod Bay	1.64	1.92	1.72	2.07	1.88	1.92	1.78	1.87	1.97	1.95	1.96	2.01	2.14	1.93	1.65	1.66	1.57	1.65	1.41	1.63
	81%	85%	82%	87%	85%	85%	83%	85%	86%	86%	86%	87%	88%	86%	81%	81%	79%	81%	76%	80%
Outer Cape Cod	0.54	0.55	0.53	0.52	0.57	0.55	0.66	0.66	0.62	0.63	0.71	0.72	0.78	0.79	0.72	0.68	0.76	0.76	0.59	0.66
	42%	42%	41%	41%	43%	42%	48%	48%	46%	47%	51%	51%	54%	55%	51%	49%	53%	53%	45%	48%
Buzzards Bay	2.97	2.53	2.26	2.21	2.36	2.41	2.36	2.35	2.14	2.27	3.08	2.70	3.11	2.85	2.44	2.37	2.14	2.57	2.42	2.38
	95%	92%	90%	89%	91%	91%	91%	94%	88%	90%	95%	93%	96%	94%	91%	91%	88%	92%	91%	91%

Table 9A. Instantaneous fishing mortality estimates (F), COHORT analysis (von Bertalanffy), by state and region, commercial lobster trap catch survey, Massachusetts coastal waters, 1981-2000.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
State	1.14	1.21	1.17	1.19	1.25	1.28	1.32	1.36	1.36	1.32	1.43	1.44	1.48	1.41	1.32	1.29	1.33	1.36	1.22	1.38
Cape Ann	1.33	1.47	1.11	1.33	1.28	1.22	1.30	1.37	1.12	1.04	1.50	1.32	1.14	1.25	1.36	1.28	1.41	1.26	1.07	1.36
Beverly-Salem	1.42	1.47	1.64	1.68	1.81	1.93	1.89	2.02	1.95	1.86	2.08	2.16	1.96	1.94	1.88	1.91	2.01	1.87	2.03	1.88
Boston Harbor	---	---	---	1.77	1.70	1.80	1.87	1.83	1.94	1.86	2.01	1.97	1.97	1.90	1.85	1.89	1.79	2.13	1.98	2.21
Cape Cod Bay	1.53	1.60	1.58	1.73	1.59	1.70	1.56	1.70	1.82	1.72	1.66	1.71	1.85	1.66	1.47	1.44	1.47	1.52	1.40	1.59
Outer Cape Cod	0.47	0.48	0.45	0.42	0.47	0.47	0.57	0.53	0.54	0.51	0.59	0.61	0.65	0.68	0.62	0.59	0.65	0.64	0.52	0.63
Buzzards Bay	2.32	2.13	1.94	1.80	2.04	2.11	2.08	2.06	1.95	1.97	2.34	2.26	2.39	2.31	2.05	1.98	1.75	2.06	2.14	1.98

Table 9B. Instantaneous fishing mortality estimates (F), COHORT analysis with Delta T's, by state and region, commercial lobster trap catch survey, Massachusetts coastal waters, 1981-2000.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
State	0.82	0.88	0.86	0.87	0.91	0.95	0.96	0.97	0.96	0.93	0.99	0.99	1.04	0.98	0.93	0.92	0.94	0.97	0.89	0.91
Cape Ann	0.82	0.85	0.90	0.92	0.93	0.92	0.92	0.96	0.88	0.76	1.03	0.97	0.86	0.88	0.97	0.89	0.99	0.92	0.75	0.86
Beverly-Salem	0.86	1.02	1.05	0.88	1.21	1.23	1.19	1.26	1.25	1.16	1.26	1.32	1.21	1.19	1.19	1.22	1.16	1.18	1.25	1.16
Boston Harbor	---	---	---	1.15	1.14	1.10	1.15	1.17	1.22	1.20	1.24	1.21	1.23	1.22	1.17	1.22	1.18	1.34	1.24	1.54
Cape Cod Bay	1.04	1.14	1.04	1.19	1.12	1.19	1.09	1.13	1.19	1.17	1.09	1.11	1.19	1.00	0.95	0.91	0.98	1.00	0.99	1.01
Outer Cape Cod	0.31	0.34	0.31	0.28	0.34	0.34	0.41	0.38	0.39	0.39	0.44	0.44	0.50	0.51	0.49	0.46	0.52	0.50	0.39	0.40
Buzzards Bay	1.10	1.02	1.00	0.94	1.05	1.07	1.04	1.06	1.00	1.00	1.15	1.07	1.14	1.11	0.98	0.99	0.87	1.03	1.07	0.97

Table 10. Estimated exploitation rate (u), by state and region, commercial lobster trap catch survey, Massachusetts coastal waters, 1981-2000.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
State	0.62	0.64	0.63	0.64	0.65	0.66	0.68	0.69	0.69	0.69	0.68	0.69	0.69	0.68	0.66	0.66	0.68	0.67	0.67	0.68
Cape Ann	0.74	0.80	0.61	0.68	0.71	0.67	0.70	0.71	0.63	0.51	0.70	0.67	0.62	0.65	0.67	0.66	0.68	0.65	0.64	0.74
Beverly-Salem	0.71	0.71	0.75	0.79	0.79	0.83	0.77	0.88	0.76	0.85	0.82	0.79	0.78	0.79	0.79	0.80	0.79	0.76	0.86	0.74
Boston Harbor	---	---	---	0.82	0.81	0.80	0.84	0.84	0.86	0.85	0.82	0.81	0.76	0.75	0.78	0.77	0.75	0.75	0.82	0.74
Cape Cod Bay	0.75	0.71	0.75	0.73	0.72	0.75	0.73	0.77	0.79	0.76	0.73	0.74	0.76	0.74	0.72	0.70	0.74	0.75	0.75	0.76
Outer Cape Cod	0.37	0.37	0.35	0.33	0.36	0.36	0.41	0.38	0.40	0.38	0.42	0.44	0.45	0.47	0.44	0.43	0.45	0.45	0.40	0.45
Buzzards Bay	0.74	0.78	0.77	0.72	0.79	0.80	0.80	0.82	0.80	0.78	0.72	0.78	0.74	0.76	0.76	0.76	0.72	0.74	0.80	0.73

Table 11. Mean carapace length (mm), by state and region, for all marketable American lobster sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-2000.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
State	88.5	87.9	88.1	88.2	87.8	87.6	87.5	88.2	88.9	89.0	88.7	88.8	88.4	88.8	89.2	89.2	89.2	88.9	89.6	89.4
Cape Ann	88.6	88.3	88.3	87.9	88.4	88.3	88.0	88.3	89.3	90.3	88.4	88.8	89.6	89.6	88.7	89.5	88.7	89.0	90.8	90.1
Beverly-Salem	87.6	87.0	86.6	86.9	86.2	86.2	85.8	87.1	87.7	88.3	87.5	87.2	87.5	87.8	88.0	87.9	87.5	87.7	87.6	87.8
Boston Harbor	---	---	---	86.8	86.9	86.4	86.6	87.5	88.0	88.1	87.8	87.9	87.5	87.5	88.0	87.7	87.8	87.0	87.5	86.8
Cape Cod Bay	87.2	86.4	86.9	86.1	86.4	86.3	86.7	87.3	87.7	87.7	88.1	88.2	87.7	88.3	89.2	89.0	89.4	89.0	89.4	89.1
Outer Cape Cod	98.2	97.5	97.4	99.7	97.0	96.3	94.6	95.2	96.5	96.1	95.3	95.2	93.8	94.2	94.2	94.9	93.9	94.1	96.6	96.1
Buzzards Bay	84.7	85.2	85.7	85.8	85.2	85.3	85.3	86.1	87.4	87.0	86.4	86.9	86.5	86.5	87.4	87.3	87.8	87.0	87.0	87.4

Table 12. Mean carapace length (mm), by state and region for all sub-legal American lobster, sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-2000.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
State	75.8	76.3	76.2	76.1	76.3	76.1	76.1	76.3	77.5	77.6	76.7	76.2	76.9	77.5	77.8	78.2	77.8	77.2	77.4	77.1
Cape Ann	78.0	77.7	77.5	77.3	77.6	77.1	75.9	77.0	78.3	78.8	78.7	77.9	78.3	78.0	77.2	77.3	77.3	77.8	76.8	75.1
Beverly-Salem	74.3	76.5	74.9	76.1	75.9	74.7	74.7	74.5	76.4	76.1	73.4	73.5	75.1	75.8	76.2	76.8	76.0	74.9	75.4	75.6
Boston Harbor	---	---	---	77.1	76.9	76.9	76.5	75.6	76.8	77.4	75.4	74.6	75.3	76.0	77.3	76.9	76.3	72.7	74.9	74.4
Cape Cod Bay	76.6	76.4	76.7	75.6	76.1	76.2	75.6	76.9	77.9	77.8	77.4	76.8	76.7	78.6	78.6	79.8	79.1	79.3	78.9	78.6
Outer Cape Cod	75.9	76.2	77.1	75.1	76.6	75.9	77.0	77.1	76.8	78.8	78.8	79.0	79.4	78.3	80.0	79.9	80.0	79.6	79.6	79.7
Buzzards Bay	75.8	75.5	76.8	76.4	76.1	76.0	76.6	76.3	77.7	77.4	76.6	77.1	78.3	77.6	77.7	77.5	78.2	78.9	78.7	79.6

Table 13. Mean carapace length (mm) of all ovigerous female American lobster, by state and region, sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-2000.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
State	88.5	87.6	88.6	87.4	87.9	88.1	87.1	87.2	88.5	88.0	86.0	85.5	85.3	86.3	86.6	86.7	86.7	86.4	88.1	88.1
Cape Ann	109.0	100.3	94.3	90.5	93.8	95.0	91.6	94.0	100.4	95.1	91.7	91.9	91.0	92.7	93.3	91.5	90.7	91.3	94.7	95.7
Beverly-Salem	80.5	84.5	85.8	83.5	85.9	83.5	81.8	83.0	85.2	85.5	83.8	81.6	82.6	83.1	83.3	83.0	82.5	83.0	83.0	83.1
Boston Harbor	---	---	---	82.1	84.0	81.3	82.3	83.7	83.0	83.8	82.0	82.0	80.8	80.9	81.5	82.4	82.2	79.8	81.2	81.3
Cape Cod Bay	86.4	83.8	85.5	84.4	85.2	86.8	87.0	84.7	86.1	85.0	83.9	84.1	83.0	84.8	85.2	85.7	85.8	86.1	87.3	86.5
Outer Cape Cod	109.8	106.1	108.0	107.1	106.9	107.3	102.5	105.2	105.4	104.6	101.9	99.2	100.7	100.0	100.6	99.6	98.9	98.3	104.7	103.7
Buzzards Bay	78.1	79.6	81.6	83.0	80.1	79.4	80.2	80.6	81.3	80.8	79.8	79.9	81.0	81.5	81.8	82.6	84.0	82.7	82.8	84.5

Table 14. Cull rate (percent), by state and region, for all American lobster sampled during commercial lobster trap catch survey, Massachusetts coastal water, 1981-2000.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
State	10.0	10.8	10.7	14.8	18.1	20.9	17.0	18.2	19.2	18.6	18.1	18.5	20.3	21.9	22.0	17.1	20.4	20.9	20.8	18.2
Cape Ann	10.0	9.8	10.5	11.5	23.9	25.3	20.2	21.2	16.7	16.7	19.7	18.2	19.2	17.1	19.6	18.0	18.9	17.6	16.7	18.8
Beverly-Salem	8.3	8.6	10.2	20.9	23.0	30.0	24.1	26.3	28.6	27.3	28.9	22.7	28.3	30.8	25.1	24.6	23.4	25.1	23.6	21.8
Boston Harbor	---	---	---	13.3	19.3	19.1	16.9	16.3	13.8	14.7	13.5	17.2	23.4	23.0	22.2	17.8	19.8	17.4	17.7	15.3
Cape Cod Bay	11.1	10.7	10.9	15.6	18.3	21.6	16.2	17.4	22.8	20.5	18.9	18.3	18.1	19.4	21.5	16.2	23.6	25.6	24.6	20.0
Outer Cape Cod	5.7	11.3	8.9	13.0	13.4	16.1	12.6	15.0	14.0	15.5	13.2	15.7	17.3	20.1	19.0	13.3	14.5	14.2	14.9	16.4
Buzzards Bay	13.5	14.7	12.4	12.4	13.4	14.6	15.1	15.6	12.6	13.6	13.9	19.3	20.5	24.0	24.4	15.0	15.6	16.2	18.5	14.5

Table 15. Cull rate (percent), by state and region, for all legal-sized American lobster, sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-2000.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
State	8.1	9.7	9.2	12.7	14.8	17.0	14.7	15.7	14.9	15.4	15.6	17.1	17.4	21.2	20.9	15.1	19.3	19.5	19.1	17.8
Cape Ann	10.7	9.6	7.5	10.4	19.4	20.3	18.0	19.3	13.9	13.7	16.8	18.3	16.3	16.5	16.2	14.7	15.7	16.3	13.7	15.8
Beverly-Salem	4.3	7.7	7.4	15.5	19.3	22.1	17.1	21.4	18.7	25.6	22.8	19.9	24.6	25.4	18.8	20.2	17.2	20.0	18.2	17.8
Boston Harbor	---	---	---	10.1	16.2	15.8	12.9	13.1	9.9	9.9	12.3	14.0	17.5	18.0	20.1	17.3	17.3	14.2	14.0	15.6
Cape Cod Bay	9.3	9.3	10.0	13.2	14.5	18.1	15.0	15.6	12.0	16.3	17.8	16.8	16.3	21.7	23.3	16.0	25.3	26.2	24.3	20.9
Outer Cape Cod	5.3	10.3	8.1	13.3	12.5	14.9	13.1	14.3	13.3	14.1	12.8	15.3	16.4	19.9	18.3	12.6	13.6	13.5	14.5	16.4
Buzzards Bay	16.1	13.2	12.7	12.3	13.8	13.6	15.2	14.1	12.6	12.6	11.5	22.2	18.9	23.5	22.6	12.9	15.3	14.4	17.8	14.6

Table 16. Cull rate (percent), by state and region, for marketable American lobster sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-2000.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
State	8.2	9.9	9.2	13.2	16.2	17.6	14.7	16.0	15.2	15.6	16.1	17.6	17.6	21.9	21.5	15.6	19.2	19.7	19.6	18.4
Cape Ann	10.8	9.8	7.3	10.5	20.9	20.7	18.4	19.9	14.0	14.2	16.8	18.1	13.7	16.7	16.0	14.5	15.3	16.3	14.1	18.9
Beverly-Salem	4.4	8.0	7.4	15.6	18.5	22.2	17.2	21.3	18.9	23.8	23.1	20.0	24.7	25.5	19.0	20.0	17.0	20.1	18.0	21.7
Boston Harbor	---	---	---	10.2	16.2	15.7	12.8	13.1	9.9	9.9	12.4	14.0	17.5	18.0	20.1	17.4	17.4	14.5	14.2	15.6
Cape Cod Bay	9.3	9.3	10.0	13.2	15.9	18.2	14.8	15.6	19.1	16.2	17.8	16.7	16.2	22.3	23.3	16.6	25.5	26.1	24.7	21.1
Outer Cape Cod	5.3	10.9	8.6	14.8	12.9	16.8	13.2	14.9	13.9	14.6	14.1	16.8	17.3	21.6	20.4	14.0	13.7	14.0	15.6	17.2
Buzzards Bay	16.9	13.1	12.3	12.6	15.4	14.1	15.4	14.7	13.0	12.4	11.7	22.5	19.4	23.7	23.0	13.1	14.5	15.1	17.7	14.6

Table 17. Cull rate (percent), by state and region, for sub-legal American lobster, sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-2000.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
State	11.2	11.5	11.6	16.1	20.2	23.2	18.2	19.6	21.1	20.2	19.2	19.3	21.8	22.5	22.9	18.5	21.3	21.9	22.7	18.6
Cape Ann	8.0	10.6	12.6	12.2	26.9	28.7	21.5	22.1	17.9	18.3	21.0	18.2	20.9	17.4	21.2	20.6	20.8	18.4	20.5	23.0
Beverly-Salem	10.0	9.0	11.2	22.3	24.0	31.8	25.3	28.6	30.8	29.2	31.6	23.5	29.5	33.8	28.2	26.7	25.5	27.2	27.5	24.2
Boston Harbor	---	---	---	14.5	20.5	20.0	18.0	18.0	15.2	16.4	13.9	18.3	25.3	24.7	22.8	17.9	20.6	18.0	19.6	15.1
Cape Cod Bay	11.9	11.3	11.4	17.0	20.2	23.4	16.8	18.3	24.0	21.8	19.2	19.0	18.8	17.8	20.1	16.3	22.2	25.2	25.1	19.0
Outer Cape Cod	7.8	17.9	13.5	11.7	18.6	22.8	11.0	16.9	17.1	20.7	14.3	17.1	20.2	21.3	21.8	15.7	17.9	16.6	18.3	16.1
Buzzards Bay	12.7	15.2	12.2	12.4	13.3	14.9	15.0	16.2	12.6	13.9	14.5	18.3	21.0	24.2	25.6	16.4	15.8	17.4	19.1	14.5

Table 18. Percent trap mortality by state and region for all American lobster sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-2000.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
State	0.15	0.04	0.22	0.15	0.18	0.20	0.10	0.15	0.12	0.17	0.37	0.08	0.13	0.22	0.21	0.18	0.10	0.15	0.18	0.18
Cape Ann	0.00	0.00	0.09	0.27	0.03	0.16	0.00	0.03	0.13	0.09	0.48	0.10	0.11	0.14	0.28	0.17	0.04	0.13	0.40	0.16
Beverly-Salem	0.00	0.00	0.00	0.00	0.04	0.22	0.03	0.19	0.14	0.29	0.41	0.13	0.19	0.13	0.74	0.49	0.37	0.31	0.39	0.41
Boston Harbor	---	---	---	0.00	0.03	0.03	0.23	0.09	0.03	0.04	0.01	0.03	0.06	0.04	0.01	0.04	0.08	0.08	0.19	0.11
Cape Cod Bay	0.00	0.02	0.03	0.00	0.00	0.02	0.15	0.00	0.02	0.05	0.02	0.02	0.02	0.00	0.03	0.03	0.03	0.08	0.04	0.07
Outer Cape Cod	0.46	0.22	0.23	0.48	0.40	0.85	0.27	0.66	0.47	0.62	0.35	0.24	0.30	0.58	0.38	0.43	0.21	0.25	0.25	0.15
Buzzards Bay	0.62	0.00	1.13	0.43	0.76	0.25	0.01	0.18	0.11	0.18	1.74	0.10	0.29	0.71	0.16	0.21	0.00	0.19	0.20	0.34