



## Natural Heritage & Endangered Species Program

[www.mass.gov/nhesp](http://www.mass.gov/nhesp)

Massachusetts Division of Fisheries & Wildlife

## Upland White Goldenrod *Oligoneuron album* (Nutt.) Nesom

State Status: **Endangered**


Federal Status: **None**

**DESCRIPTION:** Upland White Goldenrod is a relatively short plant (30–45 cm; 12–18 in.) of the composite family (Asteraceae); though it looks similar to what we commonly identify as an aster, it is actually a goldenrod. Robust plants may have a cluster of numerous slender, straight stems, referred to as a “tuft.” Flower heads may be few or many, but are never densely grouped or in one-sided arrangements.

**AIDS TO IDENTIFICATION:** Stems are mostly glabrous below, but gradually become densely short pubescent above and on the branches. Depending on availability of moisture, basal and lower-most stem leaves may be present during flowering. Its firm, ascending leaves are narrow, and taper to long points. Leaves gradually become shorter higher on stems and branches. Unlike most goldenrods, it forms a flat-topped inflorescence of white-rayed, aster-like flower heads, which are located towards the ends of short branches. This species flowers from late July into early September; peak flowering is during August. Fruit matures within three weeks of cessation of flowering.



*Upland White Goldenrod in its favored open rocky ledge habitat.*  
Photo: Arieh Tal.



Distribution in Massachusetts  
1985 - 2010  
Based on records in the  
Natural Heritage Database

**SIMILAR SPECIES:** Upland White Goldenrod is the only white-rayed goldenrod that forms a flat-topped inflorescence. By comparison, Silverrod (*Solidago bicolor*) is typically taller (up to 100 cm; >3 ft.), with an elongate, slender inflorescence bearing numerous heads that are bunched close together at nodes along stems and branches. Upland White Goldenrod is more likely to be confused with two aster species in the genus *Sericocarpus*, specifically Toothed White-topped Aster (*S. asteroides*, syn. *Aster paternus*) and Narrow-leaf White-topped Aster (*S. linifolius*, syn. *Aster solidagineus*). All three of these species have flat-topped inflorescences and heads with both white rays and white disc florets.

*A Species of Greatest Conservation Need in the Massachusetts State Wildlife Action Plan*

## Massachusetts Division of Fisheries & Wildlife

1 Rabbit Hill Rd., Westborough, MA; tel: 508-389-6300; fax: 508-389-7890; [www.mass.gov/dfw](http://www.mass.gov/dfw)

Please allow the Natural Heritage & Endangered Species Program to continue to conserve the biodiversity of Massachusetts with a contribution for ‘endangered wildlife conservation’ on your state income tax form, as these donations comprise a significant portion of our operating budget.

[www.mass.gov/nhesp](http://www.mass.gov/nhesp)

However, unlike Upland White Goldenrod, the tips of the phyllaries (involucral bracts) of the two *Sericocarpus* species are abruptly recurved. Moreover, the leaves of the two *Sericocarpus* species are oval- or egg-shaped and short-pointed, rather than narrow and long-pointed as are the leaves of Upland White Goldenrod. Lastly, the leaves of the *Sericocarpus* species are not ascending.

**HABITAT:** This species inhabits open or partially-shaded dry rocky outcrops of sandstone, shale, or limestone. It prefers calcareous or circumneutral substrate. It is often found growing in cracks or fissures in the outcrops; current Massachusetts populations are known from rocky shores of the Connecticut River. Associated species vary among sites, but include Harebell (*Campanula rotundifolia*), Little Bluestem (*Schizachyrium scoparium*), Big Bluestem (*Andropogon gerardii*), wild ryes (*Elymus* spp.), goldenrods (*Solidago* spp.), and various asters (*Symphotrichum* spp.).

**RANGE:** Upland White Goldenrod ranges from Quebec to Saskatchewan, south to Oklahoma, Tennessee, and Georgia. However, it is most common in the central and north-central midwestern states. It is at the extreme eastern edge of its range in New England, and is rare throughout the region.

**POPULATION STATUS IN MASSACHUSETTS:** Upland White Goldenrod is listed under the Massachusetts Endangered Species Act as Endangered. All listed species are protected from killing, collecting, possessing, or sale and from activities that would destroy habitat and thus directly or indirectly cause mortality or disrupt critical behaviors. The few documented Massachusetts records are from Hampden, Hampshire, and Franklin Counties.

**MANAGEMENT AND THREATS:** Upland White Goldenrod requires significant sunlight exposure and should be monitored for over-shading and competition from invasive and aggressive native species; if deemed a threat, pruning or other means of vegetation control may be warranted. As current Massachusetts populations are associated with the Connecticut River, hydrologic regime changes, shoreline development, and recreational use are potential threats. Development and recreation (e.g., trails, boat launch areas) should be directed away from Upland White Goldenrod habitat. All active management of rare plant populations (including invasive species removal) is subject to review under the Massachusetts Endangered Species Act, and should be planned in close consultation with the Massachusetts Natural Heritage & Endangered Species Program.

**Flowering time in Massachusetts**

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec

Updated 2019

**A Species of Greatest Conservation Need in the Massachusetts State Wildlife Action Plan**

Please allow the Natural Heritage & Endangered Species Program to continue to conserve the biodiversity of Massachusetts with a contribution for 'endangered wildlife conservation' on your state income tax form, as these donations comprise a significant portion of our operating budget.