

Habitat Classification Feasibility Study for Coastal and Marine Environments in Massachusetts



Katherine Lund and Anthony R. Wilbur
Massachusetts Office of Coastal Zone Management
Executive Office of Environmental Affairs
Commonwealth of Massachusetts

2007



Cover photographs credits: United States Geological Survey, Fred Short (University of New Hampshire), and winter flounder (*Pseudopleuronectes americanus*).

Massachusetts Office of Coastal Zone Management
251 Causeway Street, Suite 800
Boston, MA 02114

CZM Information Line: (617) 626-1212

CZM Web Site: www.mass.gov/czm

To cite this document:

Lund, K. and A.R. Wilbur. 2007. Habitat classification feasibility study for coastal and marine environments in Massachusetts. Massachusetts Office of Coastal Zone Management. Boston, MA. 31 pages + appendices.

Contact:

Anthony Wilbur
Massachusetts Office of Coastal Zone Management
Tony.Wilbur@state.ma.us

TABLE OF CONTENTS

EXECUTIVE SUMMARY	1
OBJECTIVES	2
BACKGROUND	3
Estuarine and Marine Habitats in Massachusetts	3
Habitat Management	3
Habitat Classification	5
ROLE OF HABITAT CLASSIFICATION IN RESOURCE MANAGEMENT	7
Coastal Zone Management Considerations	8
Geographic Information System Considerations	9
HABITAT CLASSIFICATION SCHEMES REVIEWED BUT NOT RECOMMENDED	10
Cowardin et al. (1979)	10
Dethier (1992)	
Brown (1993)	
Allee et al. (2000)	
Brown (2002)	
Madley (2002)	
European Classifications Studies – Connor et al. (2004) and EUNIS (online system)	
RECOMMENDED HABITAT CLASSIFICATION SCHEMES	13
Madden et al. (2005), “Coastal and Marine Systems of North America – Framework for an Ecological Classification Standard: Version II”	13
Kutcher et al. (2005), “A Recommendation for a Comprehensive Habitat and Land Use Classification System for the National Estuarine Research Reserve System”	16
Greene et al. (In Press), “Construction of Digital Potential Marine Benthic Habitat Maps Using a Coded Classification Scheme and Their Application”	19
Valentine et al. (2005), “Classification of Marine Sublittoral Habitats, with Application to the Northeastern North America Region”	21
Comparison of Greene et al. (1999; In Press) and Valentine et al. (2005)	24

RECOMMENDATIONS	25
Before Choosing a Classification Scheme	25
Evaluating a Pilot Area	26
After Evaluating Classification Schemes	26
CONCLUSION	28
ACKNOWLEDGEMENTS	29
LITERATURE CITED	30
PERSONAL COMMUNICATION	32
APPENDICES	33
1: Classification hierarchy / organization of four recommended habitat classification studies	
A. Madden et al. (2005), "Coastal and Marine Systems of North America – Framework for an Ecological Classification Standard: Version II"	35
B. Kutcher et al. 2005, "A recommendation for a comprehensive habitat and land use classification system for the National Estuarine Research Reserve System"	48
C. Greene et. al (In Press), "Construction of digital potential marine benthic habitat maps using a coded classification scheme and their application"	51
D. Valentine et al. (2005), "Classification of Marine Sublittoral Habitats, with Application to the Northeastern North America Region"	55
2: Contact information of habitat classification experts	58

EXECUTIVE SUMMARY

Habitat classification is the process of identifying habitat types based on a set of standard terms and descriptors. While several marine habitat classification schemes are available and used by various researchers and resource management agencies, none are universally accepted because each scheme has unique benefits and challenges. For example, the most widely used habitat classification system was developed for wetlands by Cowardin et al. (1979), which does not adequately address subtidal marine habitats. Therefore, other classification schemes were developed to meet local needs and incorporate a wider range of coastal and marine habitats.

Habitat classification frameworks are typically designed to organize geospatial data in a nested hierarchy. Nested hierarchical classification divides major groups or classes into subordinate groups or classes and allows the necessary flexibility required for decision making and/or use of available data. However, existing classification schemes were developed with different goals, data organization structure and intended application. The differences in schemes result in particular frameworks suitable for defining broad scale (>km) systems or zones, such as estuary or deepwater basin, compared to other approaches more effective at delineating and describing specific fine-scale (cm to mm) habitat attributes (e.g., eelgrass bed and shellfish reef).

The Massachusetts Office of Coastal Zone Management (CZM) initiated this study to better understand published approaches to classify coastal and ocean habitats to facilitate mapping the state-wide distribution of habitats. CZM, in cooperation with the United States Geological Survey (USGS), initiated a seafloor mapping program in 2003 to produce high-resolution maps and geospatial data of seafloor topography and surficial sediments. These maps provide the foundation for a better understanding of oceanic benthic habitats. This study is the beginning of a process to identify appropriate habitat classification approaches for Massachusetts and evaluate the feasibility of applying the habitat classification approaches to the coastal and marine environments in Massachusetts.

An important resource for this study was the Coastal and Marine Ecological Classification Standard (CMECS; Madden et al. 2005), since CMECS included reviews and summaries of many habitat classification studies. CMECS was recently developed by NatureServe, National Oceanic and Atmospheric Administration (NOAA), and other partners to provide a national framework to classify estuarine, coastal and ocean environments of North America. CMECS and additional studies were reviewed during this study to identify and describe habitat classification schemes most pertinent for coastal and marine environments in Massachusetts. The following four habitat classification schemes are recommended for further consideration in Massachusetts:

- Greene, H.G., J.J. Bizzarro, V. O'Connell and C.K. Brylinsky. In press. Construction of digital potential marine benthic habitat maps using a coded classification scheme and their application. In H.G. Greene and B.J. Todd (eds.) , Mapping the Seafloor for Habitat Characterization. Geological Association of Canada, Special Paper 47.
- Kutcher, T.E., N.H. Garfield and K.B. Raposa. 2005 (draft). A recommendation for a comprehensive habitat and land use classification system for the National Estuarine Research Reserve System. National Estuarine Research Reserve, Estuarine Reserves Division. Draft report to NOAA/NOS/OCRM. Silver Spring, MD. 26 pp.

- Madden, C.J., D.H. Grossman and K.L. Goodin. 2005. Coastal and Marine Systems of North America: Framework for an Ecological Classification Standard: Version II. NatureServe, Arlington, VA.
- Valentine, P.C., B.J. Todd and V.E. Kostylev. 2005. Classification of marine sublittoral habitats, with application to the northeastern North America region. American Fisheries Society Symposium 41:183-200.

This report summarizes the goal, geographic focus, hierarchical or classification organization, classification coding system and data sources and scale for the four classification studies. The advantages and disadvantages of each classification were also evaluated in relation to determining the most suitable framework or combination of frameworks for classifying coastal and marine habitats in Massachusetts.

In addition to summarizing the four recommended classification frameworks, this study outlines the role of habitat classification in resource management, considerations for application to GIS (geographic information systems) and other related habitat classification activities in the region. Staff at the Massachusetts Division of Marine Fisheries (DMF), USGS, National Estuary Programs, US Environmental Protection Agency (USEPA) and the National Estuarine Research Reserve System not only informed this project, but could also be considered as future sources of technical advice. It is important to identify potential collaborations and overlapping efforts of these organizations and agencies.

The study also raises questions and recommends steps to consider before choosing a classification scheme, determining a pilot area to test classification schemes and after evaluating classification schemes. The application and examination of the four recommended schemes in the coastal and marine environments of Massachusetts is a particularly important step. The results of the pilot study will guide the process to ultimately identify an appropriate framework or hybrid of frameworks to create a singular and flexible classification system to classify coastal and marine habitats in Massachusetts.

OBJECTIVES

The goals of this feasibility study are to (1) identify and describe habitat classification schemes most pertinent for coastal, estuarine and marine environments in Massachusetts and (2) assess benefits, limitations, similarities and differences of select classification schemes recommended for consideration in Massachusetts. A thorough literature search and communication with contacts at federal and state agencies as well as academic researchers informed this study, which also provides an update on activities for the schemes most likely to be used by CZM. This report will also help inform the ocean management planning efforts in Massachusetts by identifying recommendations and next steps to ultimately produce a single classification framework for Massachusetts. While habitat classification is of interest to a wide variety of users, the audience for this feasibility study is coastal and marine resource managers to help them understand habitat mapping and classification issues and the referenced research and decide which classification approach(es) best meet resource management needs.

BACKGROUND

The coastal and marine environments in Massachusetts contain a relatively diverse mosaic of habitats. The types and functions of habitats are largely influenced by the position of Massachusetts at the intersection of northern waters of the Gulf of Maine and southern waters of the mid-Atlantic Bight. Managing coastal and marine resources requires a common understanding of the types of habitats within Massachusetts and a comprehension of terminology used to study and describe habitats. Habitat classification is a forthcoming and evolving tool to standardize technical jargon and organize habitat information to facilitate habitat management.

Estuarine and Marine Habitats in Massachusetts

Two biogeographic regions encompass Massachusetts, with Cape Cod marking the boundary between the Acadian and Virginian provinces (Figure 1). The provinces are distinguished by substantial differences in physical characteristics, weather patterns and biological communities. This variation exerts a strong influence on habitat abundance and function. Managing, mapping and classifying resources and habitats across biogeographic zones requires recognition of the variability in ecological features and human influences of each region.




Figure 1: Location of Massachusetts, including the Acadian and Virginian Provinces.

During the last major glacial period in New England, 10,000 to 20,000 years ago, glaciers advanced to Cape Cod. Glaciers scoured landforms and deposited sediments, leaving markedly distinct environments in northern and southern Massachusetts. The coastline in northern Massachusetts is recognized for rocky shores, while southern Massachusetts is dominated by sandy beaches. These coastal features, with a mosaic of other bottom types such as mud, are largely mirrored in the underwater environment. While the ocean environment in northern and southern Massachusetts share many characteristics and habitats, the differences in geologic framework and species assemblages influence the type and function of marine habitat supported in each region. Tyrrell (2005) and Massachusetts Office of Coastal Zone Management (2004-2005) provide a summary of physical and biogenic habitats and ecological relationships found in Massachusetts.

Habitat Management

Coastal and fishery resource managers are frequently tasked with making decisions about uses of the coastal zone and ocean environment without sufficient knowledge of the habitat types that may be impacted by proposed projects. Massachusetts faces strong development pressures in the coastal zone from existing and proposed projects such as aquaculture, wind farms, pipeline and cable installations, deepwater ports, construction of docks and piers and sewage outfalls. Pollutants, such as excessive nutrients and heavy metals, also enter coastal and ocean waters

from a variety of sources (e.g., atmospheric deposition, groundwater, stormwater, etc.). The combination of anthropogenic influences and natural disturbances potentially disrupt and degrade habitat functions and values (see Wilbur and Pentony 1998 for review of human-induced impacts to marine habitat). Without comprehensive maps of the ocean environment and standardized classification of habitats, resource management is hindered and the assessment of the relative abundance of various habitat types is impossible.

The advent of technology to map seafloor topography and surficial geology is, however, providing a fundamental information base for the ocean environment in Massachusetts. Acoustic – or sonar – mapping (e.g., sidescan, multibeam and seismic reflection) in combination with underwater imagery and benthic sampling provides accurate views of the seafloor landscape and supports efforts to classify benthic environments. By acquiring these types of data and being better able to identify and classify habitats in a standardized way, resource managers can enhance their ability to protect particularly sensitive or productive habitat types (Tyrrell 2004).

Many steps go into mapping a habitat, including collecting and analyzing physical, chemical and biological data, applying a classification system to identify and classify habitat types and analyzing data to depict habitat boundaries. Terminology to describe habitat management activities is often confusing. The following key terms are defined to provide a basis for the remainder of this study (Valentine et al. 2005):

- **Habitat characterization** produces narrative and illustrative descriptions of habitats based on geological, biological, chemical and oceanographic observations and sampling.
- **Habitat classification** is the process of identifying habitat types based on a set of standard terms.
- **Habitat mapping** is the spatial representation of described and classified habitat units.

Many different data types are used to characterize, classify and map ocean habitats. Data applied to habitat studies range from previously published seafloor geologic, biologic and bathymetric maps to newly collected remote sensing images (e.g., multibeam bathymetry, sidescan sonar and LIDAR – Light Detection And Ranging) and in situ observation such as video, still photographs, benthic grabs and trawls (Greene et al. 2005). Figure 2 shows imagery and a map illustrating habitat classes in southeast Alaska (Greene et al. In Press). In general, there are various approaches to study the seafloor, ranging from broad geologic framework research to species behavior and habitat requirements observations. The result of the variety of seafloor-related studies is a more comprehensive understanding of the types, distribution, abundance and ecological function of habitats in the estuarine and ocean environment. Furthermore, increasing the scientific knowledge of the marine environment guides the development of appropriate habitat classification frameworks.

According to the “Strategic Plan for Mapping Massachusetts’ Benthic Marine Habitats,” the first phase for a mapping program in Massachusetts should involve high-resolution acoustic or spectral sensor mapping of surficial geology (Tyrrell 2004). The second phase should integrate biological data gathered by groundtruth sampling or other biological sampling (e.g., stock assessment and benthic community monitoring) into the geological maps or targeted studies to produce seafloor habitat maps. However, “one of the biggest obstacles to conducting large-

scale benthic habitat mapping may be the lack of a commonly accepted marine habitat classification system” (Tyrrell 2004). Therefore, closely coupled with the second phase is the need to identify and choose a habitat classification scheme so that mapping efforts begin using standardized terminology and methods to classify habitat types.




Figure 2. Example of multibeam bathymetry (a.) and seafloor habitat interpretation (b.) from southeast Alaska. Habitat codes follow the benthic habitat classification scheme of Greene et al. (1999; In Press). Images reproduced from Greene et al. (In Press).

Habitat Classification

Habitat classification schemes generally assess similar abiotic and biotic variables and assign names to a combination of habitat characteristics. For example, seafloor topography and surficial sediments are fundamental to appropriately classify a seafloor habitat in all classification schemes. The number of habitats that will be recognized and classified in a region

depends on the physical and biological heterogeneity of the seabed, data availability and scale (Valentine et al. 2005).

An effective classification scheme is helpful to resource managers and scientists because it standardizes terminology, organizes data in a logical manner, allows habitat types to be coded for data management and facilitates communication among users. For example, a national agency seeking to identify and catalog all large estuaries in North America can restrict their analysis to the upper, coarse-scale levels of a classification hierarchy, while a local agency working within an estuary will use the lower, more detailed levels of a classification scheme. Using classification codes as a common standard, both agencies will be able to organize and compare results using a unified vocabulary within a common data framework. The differences in classification schemes reflect how designers choose to organize, understand and rank the structures and functions of natural systems (Valentine et al. 2005).

The most effective classification schemes are organized in a nested hierarchy, which allows data to be organized at the level of detail desired by the user and aids data analysis and mapping. A nested hierarchy is structured in such a way that the top level class is subdivided into second level classes and each second level class into third level classes and so on (Kurtz et al. 2006; Figure 3). Each lower level describes the habitat at a higher degree of detail and spatial precision.




Figure 3. Schematic showing structure of nested hierarchy and generic difference in size of features among classes.

When classifying habitat types, the level of classification achieved depends on the scale of data gathered for a particular area. For example, a map showing classified data developed from remotely sensed images without ground validation will often only be valid for less-detailed, higher (broad) classification levels. This highlights the need for groundtruthing remote sensing data and gathering more detailed information (e.g., grain size and composition, biota and water column properties) to classify habitat to the lowest and most-detailed levels in a classification hierarchy (Madden et al. 2004b). Thus, a habitat can be mapped as narrowly or as broadly as the data and purpose permit, and this flexibility of scale influences the development of habitat classification schemes (Valentine et al. 2005).

Classified habitats require consistent names and coding systems to organize data to facilitate communication among the management, science and stakeholder communities. The coding or

naming conventions typically include numeric or letter sequences (or both). Although habitat name codes can be cumbersome and require effort to decipher, they are necessary shorthand for communication in situations where long text descriptions are inappropriate and where the information needs to be incorporated into a database (Kutcher et al. 2005; Valentine et al. 2005). In geographic information system (GIS) and spreadsheet formats, a coding system also facilitates data organization and queries (Kutcher et al. 2005).

Of particular interest in Massachusetts is classifying benthic habitat. A standard approach to classifying marine benthic habitats is not established and endorsed by the scientific and management communities, even though several schemes exist or are evolving (e.g., Valentine et al. 2005 and Greene et al. In Press). The lack of a generally accepted habitat classification scheme, encompassing benthic, water column and coastal environments, leads resource managers to either modify existing approaches for their study site or develop new schemes.

ROLE OF HABITAT CLASSIFICATION IN RESOURCE MANAGEMENT

Habitat management is emerging as an important consideration for regulating coastal and fisheries resources. While aspects of habitat management, such as wetlands protection and restoration, are well-founded in resource management, the improved understanding of the ocean environment coupled with continued anthropogenic impacts highlights the need for a comprehensive strategy to manage coastal and ocean resources. New approaches to manage ocean resources (e.g., ecosystem-based management) require a solid information base, which includes identification and delineation of estuarine and marine habitat.

Maps showing habitat characteristics, such as seafloor topography and surficial sediments, are expected to improve the efficiency of managing the ocean environment throughout Massachusetts (Tierney et al. 2004). Examples of processes that will improve as a result of habitat mapping include: resource managers referencing the maps when siting projects (e.g., docks and piers, mooring fields, dredged material disposal, cable and pipeline routes, aquaculture sites, no wake zones and other activities); scientists identifying monitoring locations based on specific habitat features; and GIS analysts quantifying the amount of different habitat types protected by various management options (Tyrrell 2004). In addition, habitat maps and classifications are important to the identification of essential fish habitat (EFH) and the facilitation of habitat-based fishery management. For example, a classification code created in the contiguous western U.S. for NOAA's National Marine Fisheries Service was adopted and modified to produce EFH maps (Greene et al. In Press).

Overall, coastal and fishery managers can benefit from adopting a classification scheme because it will: (1) provide a mechanism for identifying and mapping habitats in the ocean with standardized descriptions; (2) facilitate communication among scientists, managers and stakeholders; (3) help habitat mapping efforts by combining spatial information with a standardized coding system in GIS; and (4) better organize and communicate habitat information to help address a variety of management issues.

Coastal Zone Management Considerations

Since every classification approach is a tool to accomplish a given set of objectives (Kutcher 2006), CZM needs to identify objectives for classifying habitats. The following questions are important to consider while studying classification approaches:

- *What are the specific management questions CZM hopes to address with a classification approach?*

Will a classification scheme be used to facilitate project review, organize habitat map products, help monitor habitat change through time, design restoration strategies and/or plan areas for zoning and conservation designation? The answer is most likely yes to all of these questions. Management needs should be clearly articulated to help define the type and level of classification required to achieve management objectives. Furthermore, identifying management objectives will assist in determining the resolution of data required to support a particular scheme and management strategy.

Data requirements to support classification efforts intended to address management issues need *a-priori* decision-making. For example, preliminary planning efforts to site a submerged pipeline may use coarse resolution surveys (remote sensing technologies) to understand general sedimentary characteristics and distribution of seagrass, while fine-scale resolution of biological and physical characteristics are needed to assess potential environmental impacts and determine a final pipeline route.

CZM would benefit from a scheme (or combination of schemes) that includes both detailed and broad habitat features. Nested hierarchal structures, beginning at the landscape level and succeeding into ecosystem function and habitats, lend utility to different needs of scale, data availability and management issues. Identifying key management questions will help ensure that the development and selection of a classification scheme will appropriately support management issues.

- *Does CZM want to adopt a system that has local, regional and/or national relevance?*

Madden et al. (2005) was developed at the national level and is in the process of being sanctioned by NOAA as a national approach. Future funding and resources may become available from national funding sources for projects using this scheme. In addition, the value of using a classification system may be enhanced if the scheme were congruent with a nationally accepted marine and estuarine habitat classification approach so that eventually a regional synthesis can be made between states and perhaps even Canadian provinces. It may be useful to link habitat classification and mapping efforts in Massachusetts to the wider region so that environmental problems at local scales can benefit from the ability to link to regional scale data.

Classification schemes with a particular geographic focus, such as northeastern North America (Valentine et al. 2005) or zone (e.g., deep-sea; Greene et al. 1999), are seemingly effective in classify habitats in the intended focus areas. These types of studies are adaptable (see Greene et al. In Press) to other geographic areas and habitats and may provide a more-effective means of describing fine-scale elements of habitat.

- *Does CZM want to include nearshore and even terrestrial habitats in coastal watersheds (coastal wetlands, tidal rivers, etc) in a classification approach?*

Several schemes solely focus on the offshore, ocean environment, such as the Greene et al. (1999 and In Press), Brown (2002) and Valentine et al. (2005). For example, Brown (2002) includes a supratidal zone, but falls short of capturing the upland habitats or even habitats on the coastal plain influenced by the spray zone. Both Valentine et al. (2005) and Greene et al. (1999 and In Press) focus on seafloor habitats and do not detail the nearshore component.

The National Estuarine Research Reserve System (NERRS; Kutcher et al. 2005) and CMECS (Madden et al. 2005) approaches incorporate nearshore and upland habitat components within their classifications. However, the NERRS approach may not include the level of detail CZM needs for offshore areas that Valentine et al. (2005) and Greene et al. (In Press) achieve.

If classification of wetland, estuarine and intertidal resources is desired, then a combination of the deepwater approaches with nearshore and upland approaches may be appropriate. Or otherwise, Madden et al. (2005), which includes a general overview of upland to offshore habitats, may be sufficient. These types of decisions cannot be made until CZM defines the spatial area where the classification scheme will be applied (and this may be related to management questions).

These management questions are critical to answer before choosing a habitat classification scheme. This report better equips CZM to choose among classification approaches and will ultimately help future mapping and habitat-based management projects.

Geographic Information System Considerations

Classification studies typically use unique numerical and/or letter codes to organize data and classify habitats, which facilitate the use of a geographic information system (GIS). GIS also provides a consistent organizational tool for managing, analyzing and monitoring habitat mapping and classification products. For example, GIS is the primary tool used by the scientific community for the compilation, analysis and display of seafloor data and the creation of seafloor habitat maps (Greene et al. In Press).

GIS eases the incorporation of maps and geospatial data from a variety of sources into a database and facilitates the use of historical and contemporary data sources in unison. GIS is also effective for updating maps and can be used to quantify and track changes in habitat types. However, without standard definitions for classifying and quantifying habitats, informative comparisons between areas cannot be easily made (Kutcher et al. 2005). Marine benthic habitat maps created with a GIS, for example, must be properly attributed (Greene et al. 2005) to accurately analyze and describe spatial and trends in seafloor conditions.

Although GIS has increased the availability, and in certain circumstances the quality and quantity, of data, users are often unaware of data limitations. Data collection and interpretation protocols must be thoroughly described within robust metadata to facilitate appropriate use of data. Seafloor mapping technology is still in relatively early stages of development, so protocols must be established to clearly identify data type, quality and interpretive processes (Greene et al. 2005). Greene et al. (2005) describes how habitat classification map products have occasionally been misinterpreted and incorrectly used because the process and quality of

the source data was not fully understood. See Greene et al. (2005) for a more thorough explanation of the advantages and disadvantages encountered and possible solutions in using GIS for mapping and classifying marine benthic habitats.

GIS is a valuable tool in science, data management and resource management. A habitat classification scheme will ultimately result in the creation of spatial data. GIS is an effective means to analyze and display spatial data and facilitates effective use of data by multiple users. Metadata is critical to facilitate appropriate application of GIS data to management strategies. GIS will be used to facilitate the identification of an appropriate method to classify habitats.

HABITAT CLASSIFICATION SCHEMES – REVIEWED BUT NOT RECOMMENDED

Many classification schemes have a particular geographic focus, may not include habitat types common in Massachusetts or may be outdated. The following classification schemes were developed with a national or international focus, or to meet local needs, and are often cited and reviewed in habitat classification literature. These schemes are briefly described, highlighting limitations for CZM's use.

Cowardin et al. (1979)

The national benchmark, "Classification of Wetlands and Deepwater Habitats of the United States," provides a geographically comprehensive, nested-hierarchical classification scheme for wetland and deepwater habitats. Cowardin et al. (1979) provides an intuitive, useful and logical classification that is based on ecological parameters. This scheme was designed by the U.S. Fish and Wildlife Service (USFWS) in the late 1970s with the goal of inventorying wetland and deepwater habitats at a national scale. It became a very successful tool and provided the basis for many subsequent studies (e.g., Dethier 1992; Greene et al. 1999; Allee et al. 2000; Kutcher et al. 2005; and Greene et al. In Press). Each of these later classifications modified the original approach to address shortcomings in marine and estuarine descriptions, deepwater habitats, high-energy environments and upland and cultural habitats. Although inclusive of deepwater habitats, the Cowardin et al. (1979) system is mostly derived from a wetland-centric perspective and heavily emphasizes wetland habitat types, particularly coastal marshes. While the concept of a universally applicable coastal habitat classification could be construed as being initially tested and validated by the Cowardin system, CZM should focus on later-derived schemes to address its shortcomings (e.g., lack of detail for subtidal environments).

Dethier (1992)

"The Marine and Estuarine Habitat Classification System for Washington State" is based on Cowardin et al. (1979). Dethier (1992) adds energy as a level, significantly enhancing the utility of the 1979 classification not only for Washington State but for use in high-energy coasts. The energy element incorporates concepts such as exposed, moderately exposed and sheltered designations for rocky substrates. This modification is an example of adapting a regional approach to reflect local features. Another example of modifying an existing scheme was the narrowing of the Dethier (1992) classification focus to marine and estuarine systems, eliminating rivers and lakes. Dethier (personal communication) indicated that the 1992 scheme would not be as appropriate for Massachusetts because of its focus on Pacific Northwest environments.

Brown (1993)

“A Classification System of Marine and Estuarine Habitats in Maine: An Ecosystem Approach to Habitats” describes a preliminary hierarchical classification system organized by substratum, depth, energy level and salinity. This study included a review of classification systems (however, since this report is relatively old, the study did not include recent classification schemes that were reviewed as part of the CMECS approach [Madden et al. 2005]). Ultimately, the scheme for Maine builds primarily on those developed by Cowardin et al. (1979) and Dethier (1992). Hierarchical levels of the classification include: system (biome types), subsystem (tidal regimes), class (substrata), subclass (energy levels), modifiers (as needed), diagnostic species and common species.

This study contains good descriptions of habitat types and associated species and provides examples of applications of the classification approach. Habitat types identified for Maine closely relate to those found in northern Massachusetts, making this study a potential model for presenting information for specifically classified areas. However, the document does not include classification for pelagic areas and lacks a numeric or letter coding system that is important for developing a GIS.

Allee et al. (2000)

The “Marine and Estuarine Ecosystem and Habitat Classification” scheme was developed to be comprehensive for coastal and marine systems, encompassing the entire U.S. coastal margin from the landward extent of tidal influence to the outer edge of the continental shelf. Allee et al. (2000) was the first national classification since Cowardin et al. (1979) and presents several enhancements, including (1) focus on universal applicability across a variety of target regions; (2) stronger focus on marine and estuarine landscapes; and (3) linkages between geologic framework, energy and biology.

The NERRS reviewed Allee et al. (2000) and found the scheme difficult to use because of departure from traditional language in Cowardin et al. (1979), its complex format and the detail necessary to classify at lower levels (NERRS 2000). In addition, NERRS reviewers characterized the scheme at its lowest levels of habitat descriptions to be “open-ended” with choices being subjective and dependent on user knowledge and discretion rather than having standardized categories, which made it harder to compare across sites. In addition, marine and estuarine categories are classified together, which resulted in NERRS sites with the same classification for freshwater and brackish habitats, despite having completely different species composition due to the salinity gradient. These categories should be distinct and split early in the hierarchy (NERRS 2000). Another concern identified in the NERRS review was that Allee et al. (2000) does not follow a hierarchy, which makes it less user-friendly.

Brown (2002)

“Our Living Oceans Benthic Habitat Classification System” was developed to describe and define critical habitats for federally managed fishery species and consists of five major habitat types (Freshwater, Estuarine, Nearshore, Offshore and Oceanic Islands and Banks). This hierarchical scheme was also reviewed by the NERRS (NERRS 2000). The scheme does not provide protocol for dealing with habitat disturbances or habitat gradients. Brown (personal communication) recommended using Madden et al. (2005), which, “although it's still in the developmental phase, is a much more mature product. The framework I had been working on has pretty much dropped by the wayside.”

Madley (2002)

The “Florida System for Classification of Habitats in Estuarine and Marine Environments (SCHEME)” focuses on estuarine and marine environments that are relevant to coastal resource managers in Florida. Habitats of particular focus include nearshore and neritic areas inhabited by corals, hardbottom and seagrass communities. The boundary of the classification scheme is from the high tide line to the edge of the continental shelf. This approach lacks provision for descriptions of geologic structure, coastal complexity or hydrodynamic features – apparently more reflective of the calm coastal environment of the Florida Gulf coast. The scheme is data driven with Light Detection and Ranging (LIDAR), single beam sonar and aerial photography. While the hierarchy itself may not be useful to CZM for reasons above, the mapping methodology and the approach to determine how to use certain data sources to create the maps for each kind of habitat may still be relevant.

European Classification Studies

Since the mid-1990s, a European movement to map and classify coastal and marine habitats produced several notable projects, including BIOMAR (Marine Habitat Classification for Britain and Ireland) and EUNIS (European Nature Information System). These international efforts are described below because they contain overlapping ideas and useful approaches:

- The “Marine Habitat Classification for Britain and Ireland” developed in the late 1990s and revised in 2004 (Connor et al. 2004) is often referred to as the BIOMAR project. While the BIOMAR system has several useful components that CZM may want to consider (e.g., web-based), it only covers marine habitats from high tide seaward (excluding salt marshes) and does not extend to the deep ocean beyond 80m. Connor et al. (2004) focused on marine areas closely associated with the coastline, with little attention to estuarine systems and their connection to upstream watersheds. Important components of the BIOMAR scheme are incorporated in the CMECS classification (i.e., Madden et al. 2005).

It is worth noting that the BIOMAR system is designed to be easy to use and is largely web-based (<http://www.jncc.gov.uk/MarineHabitatClassification>). Five levels of classification are identified on the website including: Level 1 – Marine Environment; Level 2 – Broad Habitats; Level 3 – Main Habitats; Level 4 – Biotope Complexes; Level 5 – Biotopes. The website contains a full listing of habitat types within these levels, a detailed description of each type, distribution maps and color photographs along with a glossary of terms. The web-based approach should be considered as a model for future data presentation.

- The EUNIS (European Union Nature Information System) classification incorporates major themes of BIOMAR. EUNIS is a broader study that was developed for aquatic and terrestrial habitats for most of Europe. In addition to having a terrestrial focus, the EUNIS classification is also different from the BIOMAR because it includes deeper water column habitats. It also has a web application <http://eunis.eea.europa.eu/index.jsp>, which contains the full classification hierarchy with keys for identification of all habitat types along with a glossary of terms and background information on the classification’s rationale. While the online EUNIS classification leads users through a useful series of logical questions to help identify classification levels (i.e., permanently water-covered? shelf? substrate type? characterized by macroalgae?), the marine and coastal component is only a small focus of the overall classification scheme that has eight other main levels devoted to terrestrial habitats. In addition, the habitat types represented range from Northeast Atlantic (England

and Ireland) to the Mediterranean and Black Sea marine habitats (Davies et al. 2004). This approach is more complex and covers a much wider range of habitats than CZM needs.

RECOMMENDED HABITAT CLASSIFICATION SCHEMES

Four habitat classification schemes are recommended for CZM's consideration. Ultimately, resource management questions previously identified in this report will help CZM determine the most suitable classification scheme(s). It is important to note that the following descriptions of the four recommended classifications may be confusing, especially examples of the coding approaches, without referencing Appendix 1. Appendix 1 shows details of each hierarchy and classification code.

Madden et al. (2005), "Coastal and Marine Systems of North America – Framework for an Ecological Classification Standard: Version II"

The Coastal and Marine Ecological Classification Standard (CMECS), developed by NatureServe, NOAA and other national partners, is a classification approach for estuaries, coasts and oceans of North America (Madden et al. 2005). The study intended to create a national standard for classifying habitats to ensure data collected by different parties is consistent with regional, national and international standards. Development of standards for a U.S. marine and coastal classification system also supports federally mandated marine protected area and essential fish habitat efforts. Finally, NOAA viewed development of this scheme as a way to assist integration of federal, local, state and regional efforts to develop ecosystem-oriented resource management plans (Madden et al. 2005).

NOAA Fisheries (National Marine Fisheries Service) and NOAA's Coastal Services Center (CSC) established a partnership with NatureServe, previously the scientific arm of The Nature Conservancy, in 2002 to build upon previous NOAA studies (e.g., Allee et al. 2000) and several existing national and international frameworks. Madden et al. (2004a and 2005) conducted a thorough review of existing classification studies (e.g., Cowardin et al 1979; Dethier 1992; Greene et al. 1999; Allee et al. 2000; Madley et al. 2002; Zacharias et al. 2000; and Connor 1997 and 2004) and described elements of each study and approach to incorporate into the CMECS. Madden et al. (2005) attempted to combine the best elements of previous schemes to establish a single national marine and estuarine classification standard with broad, regional and local applicability.

Geographic Focus

The CMECS extends from the head-of-tides in the coastal zone to the deep ocean. This area encompasses estuaries, wetlands, rivers, shorelines, islands, intertidal and benthic zones and the entire water column from the shore to the deep ocean at scales of <1m to >10⁶m (Madden et al. 2005).

Hierarchical Organization

The CMECS is organized into a nested hierarchy with the following six levels (Madden et al. 2005):

- **Regime** (Level 1) – is differentiated by a combination of salinity, geomorphology and depth organized in five categories: estuarine, fresh water-influenced marine, nearshore marine, neritic and oceanic (10km² to > 1000km²).
- **Formation** (Level 2) – is relatively large physical structures formed by water (currents) or substrate (islands) (10,000m² to 100km²).
- **Zone** (Level 3) – distinguishes between water column, littoral or sea bottom (100m² to 10,000km²).
- **Macrohabitat** (Level 4) – is large physically complex structures that typically contain several habitats, such as a rocky shore (100m² to 1,000m²).
- **Habitat** (Level 5) – is a specific combination of physical (i.e., grain size) and energy characteristics that create a suitable place for colonization or use by biota (1 m² to 100m²).
- **Biotope** (Level 6) – is identified by characteristic biology associated with a specific habitat (1 m² to 100m²).

A key feature of the CMECS is the capacity to handle water column data in three dimensions with the use of the category *Zone* (water column, littoral, bottom). While users may see this complexity as a drawback if a simpler approach for mapping purposes is desired (Kutcher 2006), others see this classification level as a decision point on the focus area (i.e., water column, littoral, bottom) before continuing to lower levels of the hierarchy (Wright personal communication 2006).

Classification Coding System

Madden et al. (2005) developed a coding system based on a sequence of numbers and letters. Regime is represented by a capitol letter (**X**); Formation follows to two decimal points (**X.00**); Zone is represented by bottom (B), littoral (L) or water column (W) – **X.00.B/L/W**; Macrohabitat is distinguished by a lower case letter (**X.00.B/L/Z.x**); and Habitat is a series of numbers to two decimal points (**X.00.B/L/Z.x.00**). It is uncertain how Biotope is coded in this classification system. This classification scheme is evolving, and the coding system may change (Allee personal communication).

Example:

A	Estuarine Regime
A.02	Estuarine embayment Formation
A.02.B	Estuarine embayment bottom Zone
A.02.B.c	Estuarine embayment softbottom Macrohabitat
A.02.B.c.09	Estuarine embayment subtidal vegetated mud and mixed-fine sediments Habitat

See also <http://www.natureserve.org/getData/CMECS/app/classification/tree/pivot/browse> and <http://www.natureserve.org/getData/CMECS/> for more examples.

Data Sources and Scale

The CMECS sets uniform classification rules that function at multiple scales across a diversity of environments. The hierarchy is conceptually divided into two parts based on data required for applying the classification. The upper levels (Regime through Zone) can be identified from maps, bathymetry, remote imagery and historical data. Lower levels (Macrohabitat through Biotope) are distinguished at local spatial scales and data are collected through direct observation and measurement. The lower, more-detailed four levels require the most work and are valuable for applying the classification for purposes of habitat conservation and management (Madden et al. 2004a).

Data availability will – in most cases – prevent classification to the full (most-detailed) extent. Allee (personal communication), a primary contributor to CMECS, indicated that the Macrohabitat or Habitat levels are the most realistic levels to classify habitats because of data availability (not enough information exists to get at many Habitat or Biotope levels for regional assessments). The CMECS was, however, designed to partially identify classes at lower levels even if comprehensive data are not available. In other words, a user can skip levels where data are lacking, and as additional data in the area are gathered, the entire hierarchy will continue to grow and strengthen the understanding of the whole system (Madden et al. 2004a).

Comments and Updates

The CMECS is a collaborative and ongoing process, with NOAA, USFWS, Census of Marine Life, The Nature Conservancy, several universities and NatureServe contributing to the development of the classification framework. Development of the scheme included a national workshop in 2003, numerous interviews with scientists and managers that map and study habitat and pilot testing (Madden et al. 2005). The most recent 2005 version is now available through the NatureServe website www.natureserve.org/getData/CMECS. This product is a revision of Madden et al. (2004a). NOAA's Coastal Service Center and partners are updating CMECS and actively advocating its use as a national standard.

Numerous habitats are included in the CMECS, encompassing intertidal environments through oceanic areas. The online catalog, however, does not include a comprehensive list of all coastal and marine types. CMECS is not well developed for certain areas (e.g., nearshore non-estuarine benthic habitats). The upper three levels (Regime, System, and Zone) are fairly complete, while lower three levels are less complete. CZM should review the CMECS to make sure the scheme includes habitats in Massachusetts.

An important stage in the development of the CMECS was piloting the classification. Pilot projects were undertaken in Florida, the south Atlantic Bight and the Columbia River to help make the classification more robust and populated with information from different ecological systems. The purpose of the Florida Keys project was to map shallow water seagrass and coral reef habitat; the purpose of the south Atlantic Bight project was to locate and map marine hardbottom habitat; a large-scale test of the classification in the Columbia River estuary was to examine suitability for mapping salmon habitat (Madden et al. 2004b).

The pilot studies included examining data sources, scale, procedure, analysis and lessons learned along with tables and figures. The Columbia River study particularly provided insight to the strengths and challenges of CMECS as a national marine data structure. This 24 month pilot project implemented the scheme from Level 1 (Regime) through Level 3 (Zone) with limited use information at Level 4 (Macrohabitat). A manuscript is in development that outlines

challenges, benefits and considerations of CMECS (Wright In Press). Nan Wright is the author of this study and expressed interest in providing training on the use of CMECS for those interested in applying it to their local waters. CZM should consider this opportunity since she has experience from piloting the CMECS.

NatureServe is the primary developer of CMECS, with Chris Madden as principal investigator and major guidance and funding from NOAA. CMECS continues to evolve and NOAA's CSC is interested in testing CMECS in Massachusetts (Allee personal communication). NOAA intends to go on the road with the latest scheme for introduction to coastal managers and explain how it was developed and how it can be used (Wright personal communication). This local training could be an opportunity for CZM to gain an understanding of CMECS and applications in Massachusetts. As CZM proceeds with examining, developing and adopting a habitat classification framework for Massachusetts, NOAA's CSC should be actively engaged to ensure mutual objectives of CZM and CSC can be achieved while maintaining local research and management initiatives.

Kutcher et al. (2005), "A Recommendation for a Comprehensive Habitat and Land Use Classification System for the National Estuarine Research Reserve System"

The National Estuarine Research Reserve System (NERRS) is investigating the development of a habitat classification scheme to facilitate measuring the magnitude and extent of habitat change in estuarine systems and linking observed changes to watershed land-use practices (Kutcher et al. 2005). The NERRS habitat classification scheme is an ecologically based, nested hierarchy intended to encompass ongoing classification efforts within the NERRS, allow effective use of existing data and comprehensively inventory and classify all land-cover types. The classification scheme is GIS-based to allow integration of varying scales of data and facilitate use of data with scientists and non-scientists (Kutcher 2006).

The NERRS scheme is largely based on Cowardin et al. (1979), with classes directly adopted or slightly modified from Cowardin et al. (1979) and expanded to include upland and cultural habitats (Kutcher et al. 2005). Cowardin et al. (1979) provided the foundation for the collection and classification of wetland data through the National Wetlands Inventory (NWI), and NWI data were available for most sites. Incorporating NWI, with expanded classes and descriptions for site-specific features, allowed historical analyses within the NERRS (Kutcher personal communication). NOAA's CSC Coastal Change Analysis Program (C-CAP; Dobson et al. 1995) approach also informed the NERRS scheme. The C-CAP was referenced for coarse level mapping of watershed habitats and land uses. Anderson et al. (1976) informs cultural land use classes.

Geographic Focus

Boundaries of NERRS typically include coastal watersheds and estuaries without an offshore, ocean environment component. This classification scheme was designed to facilitate objectives related to geospatial inventory and land cover change analyses of coastal uplands, wetlands and nearshore habitats.

Hierarchical Organization

The NERRS scheme is organized in a five-level, nested hierarchy. The following hierarchy begins with a broad level and progresses to finer classes, designed to guide the user through progressively more detailed attributes:

- **System** (Level 1) – is based on predominant water source (e.g., marine, estuarine).
- **Sub-System** (Level 2) – is based on hydrologic influence such as intertidal.
- **Class** (Level 3) – is based on vegetation or physical attributes (surficial geology).
- **Subclass** (Level 4) – is based on attributes such as grain size.
- **Descriptors** (Level 5) – are based on published scientific literature and defines habitat based on features such as single plant species (eelgrass) or substrate traits (mudflat).
- **Modifiers** – are used to describe habitats with additional natural features (e.g., salinity regime) and anthropogenic influences such as diked or excavated.

Although recommended scale/units were not provided for each of the top four classification levels, Walker et al. (2005) suggests a mapping scale between 1:24,000 and 1:3,000 if the source data are based on aerial photography and using on-screen digitizing techniques, a consistent on-screen viewing scale (e.g., 1:3,000 to 1:5,000) and minimum mapping unit (e.g., 0.25 acre) are needed to ensure integrity.

Descriptor and Modifiers are not within the hierarchical structure and are meant to be added as fields (columns) in the GIS attribute table. Descriptors and Modifiers are used to further describe habitats with qualitative information. For example, Modifiers describe habitat polygons like “sandy ocean bottom” with information such as circulation and tidal regime, water chemistry (salinity), anthropogenic pressures affecting habitats or percent cover of invasive species. Descriptors are common names and definitions chosen to standardize a particular habitat type such as “New England Saltmarsh.” These common names are important in helping local managers use the same terminology for a habitat type (Kutcher et al. 2005).

More background on descriptors, modifiers and how the Cowardin system was adapted to meet the needs of the NERRS is described in Kutcher et al. (2005).

Classification Coding System

Kutcher et al. (2005) use a numeric code that follows Anderson et al. (1976) and Cowardin et al. (1979), with the addition of Level 5 (Descriptors) and the option for a narrative description of detailed features (Modifiers). The coding system is arranged by eight Systems (1,000-8,000), followed by progressively more detailed classes (Sub-System, Class and Sub-Class). Descriptors (Level 5) are added to Sub-Class and represented by two decimal places (Kutcher et al. 2005).

Example:

NERR Code:	2261.01	
System:	Estuarine	(2000)
Subsystem:	Intertidal Haline	(2200)
Class:	Emergent Wetland	(2260)
Subclass:	Persistent	(2261)
Descriptor:	New England Low Salt Marsh	(2261.01)

Data Sources and Scale

The NERRS classification scheme requires data to be collected at two geographic scales: 1) broad-scale, low resolution data for characterizing entire watersheds; and 2) detailed, fine-scale, high resolution data for characterizing specific properties (Walker et al. 2005). Watershed-level data are used to measure habitat and land-use change in the area of interest, assess the sensitivity of habitats to changes in adjacent uplands and analyze the relationship between land use and water quality over time. Broad-scale data are primarily derived from 30m Landsat TM imagery and classification is automated. Reserve-level, finer scale data identify species-habitat associations and landscape ecology features and are mostly interpreted from high resolution imagery (e.g., aerial photography) or field observations.

The NERRS classification scheme provides the foundation to create a comprehensive GIS. The GIS data are capable of characterizing, organizing and displaying strictly defined land cover units (and associated attributes) that allow direct spatial analysis of all habitat and land-cover types. Each level or class represents a single column in an attribute data set, and each polygon is a single row in a GIS database. Point data (e.g., grab samples) can be assigned as Modifiers or Descriptors and organized in data columns, further characterizing each polygon with more information. Modifier or Descriptor data can also be organized as discrete GIS layers (Kutcher personal communication).

Comments and Updates

A modified version of the NERRS classification scheme was recently approved, and a Habitat Mapping and Change Technical Advisory Panel is determining how best to implement the approach (Kutcher personal communication). The NERRS classification approach will continue to adapt and improve.

The original CMECS scheme (derived from Allee et al. 2000) was piloted by several NERRS in 2000 and 2006. The major issue with the CMECS approach was that coastal uplands were not incorporated, which are crucial for analyzing processes that NERRS scientists wanted to consider (Kutcher personal communication). Kutcher et al. (2005) was developed with particular focus on estuarine and upland habitats and does not cover deeper subtidal benthic areas. The NERRS scheme contains codes for offshore areas and altered habitats but are largely untested, as most reserves do not contain deepwater (Kutcher personal communication). The use of descriptors and modifiers are not clear and should be explained in more detail, if CZM decides to use Kutcher et al. (2005).

Emergent (and most submerged) habitats were recently mapped to a species assemblage level in the Waquoit Bay National Estuarine Research Reserve (WBNERR; Annett personal communication). This mapping is more detailed than the NERRS classification system, but the NERRS approach allows flexibility in detail. WBNERR has not yet agreed to appropriate names to classify species assemblage polygons and has not populated the database for the mapped features. CZM could coordinate with WBNERR on these tasks and gain experience with the NERRS approach (Kutcher et al. 2005).

Greene et al. (In Press), “Construction of Digital Potential Marine Benthic Habitat Maps Using a Coded Classification Scheme and Their Application”

Efforts to classify seafloor habitats in California began in the late 1990s (Greene et al. 1999) and have continued to evolve, broaden and improve (Greene et al. In Press). Greene’s original classification in 1999 was developed for coastal California deep seafloor habitats, with the goal of understanding and predicting spatial distribution of fish species. The Greene et al. (1999) approach developed during more than 15 years of mapping seafloor habitats and was adopted by several organizations, agencies and countries. Greene et al. (In Press) refined the 1999 study, with the goal of establishing a standard technique for making habitat designations and comparisons of deepwater marine benthic habitats worldwide. This scheme is GIS compatible and generally based on geomorphology, substrate type and textures produced by physical processes, and sessile biology (Greene et al. In Press).

Geographic Focus

Greene et al. (1999) was developed primarily for deepwater (greater than 30m), however the approach is easily applied to nearshore shallow coastal areas and estuaries (Greene personal communication). The classification scheme applies to seafloor habitat types throughout marine regions, from high (sub-arctic) to low (tropical) latitudes, and shallow, intertidal regions and estuaries to abyssal plains (Greene et al. In Press). While the scheme is flexible and can be modified to match habitat mapping objectives, the approach should not be altered to the extent that reproducibility of delineating habitat classes is lost (Greene personal communication).

Hierarchical Organization

Physiography, depth, seafloor induration (hardness of substrate), geomorphology, texture and biology are used to classify seafloor habitats (Greene et al. In Press). Seafloor mapping datasets (e.g., multibeam bathymetry, backscatter intensity and groundtruth imagery) and analogous optical data, such as LIDAR, provide the foundation for applying this scheme. The hierarchy is largely based on spatial scale – in terms of mapping scale. Examples of the large geologic features mapped at small-scale (e.g., 1:1,000,000) are the continental shelf, abyssal plain and submarine canyons. Small characteristics such as biogenic structure (fauna forming observable features on the seafloor) and grain size require large-scale of 1:50,000 or less. The following classes progress from small-scale features to the larger-scale attributes (Greene et al. 1999):

- **Megahabitat** – is a large feature (>kilometer) defined at a small scale (e.g., 1:1,000,000), such as abyssal plain.
- **Mesohabitat** – is a relatively large feature, such as submerged canyons and extensive bedrock outcrops, defined at scales of 1:250,000 or less.
- **Macrohabitat** – ranges in size from one to 10s of meters, consisting of attributes delineated at scales of 1:50,000 or less (e.g., large boulders, sediment waves, algal mats and kelp beds).
- **Microhabitat** – is a small feature (<m) such as grain size and biogenic structure (individual habitat forming fauna such as anemones).

Classification Coding System

The attribute code associated with this scheme is adapted from Greene et al. (1999) and consists of seven primary letter and number characters (Greene et al. In Press). Codes represent (1) physiography and depth (i.e., Megahabitat – one of nine) – capital letter; (2) substrate induration – lower case letter; (3) geomorphology (i.e., Meso- and Macrohabitat) – lower case letter; (4) modifiers for texture, lithology, bedform and biology – subscript letter preceded by an underline; (5) seafloor slope or inclination – represented by five slope categories (numbers 1 to 5), (6) seafloor complexity (e.g., rugosity, vertical relief) – represented by capital A to E as measure of rugosity, and (7) geologic units represented by standard geologic symbols (Greene et al. In Press). Codes for classes one to five are typically derived from high-resolution seafloor mapping data. Characters 5-7 are optional and included only when slope and complexity can be calculated and when the geology is known.

Additional attribute codes are available to distinguish habitat types, when underwater imagery (e.g., video and photographic data) exist, consisting of geologic or substrate attributes and biological features. Underwater imagery usually allows classification to Macrohabitat or Microhabitat features, with Macrohabitat and Microhabitat preceded by an asterisk (*) in the coding system. Geologic attributes are in parentheses and biological features are in brackets. When information is inferred, a question mark (?) denotes the area of uncertainty.

The code was written to easily distinguish each habitat type and to facilitate ease of use and queries in a GIS (e.g., ArcGIS). The code is intuitive with the use of unique (non-repeatable) letters and numbers for each category (Greene et al. In Press). See Greene et al. (In Press) for detailed codes for each category.

Example: Shpd1S(Q/R)*(m)[w]1C

Large-scale habitat type:

Continental shelf megahabitat (S); flat, highly complex hard (h) seafloor with pinnacles (p) differentially eroded.

Geologic unit:

Quaternary/Recent (Q/R).

*Small-scale habitat type:

Flat or nearly flat mud (100%) bottom (m) with worm tubes [w]; flat (1) and moderate rugosity (C).

Data Sources and Scale

This scheme was developed to use acoustic (e.g., multibeam bathymetry and sidescan sonar) and/or optical (e.g., LIDAR and hyperspectral) mapping data to classify the seafloor environment. The classification approach pays particular attention to scale and emphasizes the need to groundtruth interpretations made from remotely sensed data (Greene et al. In Press). Greene et al. (1999; In Press) refer to scale in terms of mapping scale. Small-scale refers to mapping large or coarse-resolution features (e.g., continental shelf) and large-scale indicates smaller or fine resolution attributes, such as grain size, species assemblages and habitat associations.

Megahabitat can often be delineated from existing, coarse-scale data (1:1,000,000 or greater), such as maps and satellite images. Classifying Mesohabitat (1:250,000 or less) and Macrohabitat (1:50,000 or less) require high frequency (e.g., 200 kHz) seafloor mapping systems. Imaging and classification of Microhabitat is best accomplished from in-situ

observations, such as benthic grabs or underwater video (Greene et al. 2005). Classifying Microhabitats is generally more difficult and time consuming compared to the other habitat classes.

While this approach is tested in relatively deeper waters using seafloor mapping data, bathymetric LIDAR, hyperspectral systems and other technologies (e.g., digital photography) are being used in shallow-water environments and provide data to apply Green et al. (1999; In Press).

Comments and Updates

Although this scheme was based on earlier habitat classification studies by Cowardin et al. (1979) and Dethier (1992), it represents a departure from coastal habitat and moves into the relatively deep ocean. Greene et al. (1999; In Press) focus on geologic structure of the bottom, leaving the classification of estuarine and freshwater systems to existing classifications. Classifying habitats in the offshore environment is an important aspect of habitat classification in Massachusetts. However, estuarine and coastal habitats are also important to include in a singular classification framework for Massachusetts.

Greene et al. (1999; In Press) is peer reviewed and is adopted by many organizations. Pilot mapping and classification is complete in Alaska, Washington and California, and new efforts are starting in the Pacific island areas, the Baltic and elsewhere using this scheme (Greene personal communication). Thus, the approach is becoming established, is field tested and can be applied in areas with seafloor mapping data.

Valentine et al. (2005), “Classification of Marine Sublittoral Habitats, with Application to the Northeastern North America Region”

The goal of the Valentine et al. (2005) scheme is to develop a practical method to classify marine sublittoral (subtidal continental-shelf and shelf-basin) habitats in northeastern North America. The classification approach is based on topographical, geological, biological and oceanographical attributes and natural and anthropogenic processes. Valentine et al. (2005) reviews several other classifications (e.g., Cowardin et al. 1979; EUNIS 1999; Greene et al. 1999; Allee et al. 2000; and Conner et al. 2004), and uses concepts from these approaches, tailoring the classification to the region under study and basing classes on extensive seabed observations (Valentine et al. 2005).

Geographic Focus

The focus is on marine sublittoral environments, with particular focus on the seafloor, in the northwest Atlantic Ocean from the Gulf of Maine to the continental shelf off New Jersey. The classification includes subtidal areas of the continental shelf basins of the Gulf of Maine that reach depths of approximately 400m and submarine canyon heads that incise the continental shelf in depths of up to 800m (Valentine et al. 2005).

Classification Organization

The classification published by Valentine et al. (2005) is slightly different than the other schemes by being only partly hierarchical. Given this structure, classification levels do not always fit neatly into broad classes and fine classes. Each class has broad geographic application, except in instances where floral and faunal features are regional. Valentine et al.

(2005) emphasizes seabed substrate type, substrate dynamics and complexity of physical and biological features on the seabed in classifying seafloor habitats.

There are eight **Themes**, with an associated **Class**, **Subclass**, **Category** and **Attribute**. Themes are the highest level of classification and include: **One** (Topographical Setting), **Two** (Seabed Dynamics and Currents), **Three** (Seabed Texture, Hardness and Layering), **Four** (Grain Size Analysis), **Five** (Seabed Roughness), **Six** (Fauna and Flora features), **Seven** (Faunal Association, Human Use and State of Disturbance), and **Eight** (understanding of Habitat Recovery). Classes are identified by seabed features (e.g., grain size, texture and roughness), flora and fauna, and natural and anthropogenic processes that affect habitat characteristics. Subclass, Category and Attribute include more-detailed habitat characteristics that are fundamental for recognizing and analyzing habitats (Valentine et al. 2005).

Examples of Classes include seabed dynamics and currents, texture and hardness and predominant grain size; Subclass includes features such as percent composition of grain size, mobility of sediments and bedforms; Category includes type of current (tidal or storm-induced) and biogenic features (burrows); and Attributes provide a semi-quantitative assessment of an area (e.g., percent cover of biogenic structure).

Classification Coding System

Valentine et al. (2005) developed descriptive habitat codes (and names) that incorporate information on seabed substrate type, seabed substrate dynamics and the degree of physical and biological complexity on the seabed. The least descriptive habitat name is based on seabed substrate type only (e.g., mud; M). The most descriptive habitat name has a seabed dynamics term, substrate type term and structural complexity term (e.g., Immobile - I; mud - M; physical structural complexity low - ps5-10L; biological structural complexity very, very low - bs<1VVL; I_M_ps5-10L_bs<1VVL). The following are examples from Valentine et al. (2005) of the application of this coding convention:

Example 1:

Habitat code based on seabed type: Mud (M)

Habitat code based on dynamic seabed type: Immobile; mud (I_M)

Habitat code based on dynamic seabed type with physical structure (ps) and biological structure (bs): Immobile; mud; physical structural complexity low; biological structural complexity very, very low (I_M_ps5-10L_bs<1VVL)

Explanation: Habitat name is based on seabed dynamics, seabed substrate type, and physical and biological structural complexity. The code is created as follows: Immobile (I); mud (M); physical structures cover 5-10% of the seabed (ps5-10), physical complexity is low (L); biological structures cover less than 1% of the seabed (bs<1), and biological complexity is very very low (VVL)

Example 2:

Habitat code based on seabed type: Sandy gravel (sG)

Habitat code based on dynamic seabed type: Intermixed mobile/immobile; sandy gravel (IMI_sG)

Habitat code based on dynamic seabed type with physical structure (ps) and biological structure (bs): Intermixed mobile/immobile; sandy gravel; physical structural complexity very high; biological structural complexity low (IMI_sG_ps>90VH_bs5-10L)

Explanation: Habitat name is based on seabed dynamics, on seabed substrate type, and on physical and biological structural complexity. Intermixed mobile/immobile (IMI); sandy gravel (sG); physical structures cover more than 90% of the seabed (ps>90), physical structure complexity is very high (VH); biological structures cover 5-10% of the seabed (bs5-10), and biological structural complexity is low (L).

Valentine et al. (2005) describes the coding and naming process in more detail.

Data Sources and Scale

Valentine et al. (2005) is based on seafloor observations using multibeam and sidescan sonar, video and photographic transects and sediment and biological sampling (i.e., benthic grabs). Geophysical surveys, such as multibeam and sidescan sonar, depict broad geological environments and provide a framework for identifying and classifying habitats. Although modern seafloor imagery is a solid foundation for the study of habitats, classification often requires additional biological information such as underwater imagery and benthic sampling.

Themes, Classes and Subclasses can generally be interpreted from seafloor mapping data. Identifying Categories and Attributes require underwater observations (e.g., video transects and/or benthic grabs). Since it is not always feasible to quantify the abundance of individual structures and organisms, this scheme allows for the inclusion of visual semi-quantitative estimates of habitat characteristics (Valentine et al. 2005).

Data and imagery availability dictate mapping scale of classified habitat. Observations needed to classify habitats at the higher levels (i.e., Theme and Class) range from tens of meters to >1 km and can be defined using maps and seafloor images at a scale of 1:250,000-1:1,000,000. Lower classification levels require maps in the range of 1:25,000-1:100,000 (Valentine et al. 2005). While the level of mapping and classification depends in a large part on the data and imagery available, the Valentine et al. (2005) approach is flexible, allowing classification of large geologic features and fine-resolution attributes such as grain size (mm-cm).

Comments and Updates

Valentine et al. (2005) focuses on subtidal seafloor habitats and does not cover near-coastal and estuarine benthic areas. Valentine (personal communication) believes most existing classification studies are weak on using geological attributes for characterizing habitats. This assertion is founded in the geologic-centric belief that habitats are “for the most part composites of geological and oceanographic attributes and process” (Valentine personal communication). Biological resources are also fundamental to habitat, since the attraction to an area (area described by physical or biological features) is based on the dispersal, behavior and/or life history requirements of a species.

Valentine described how much of the Massachusetts sublittoral environment can be classified to the level of Subclass vs. Category vs. Attributes using his scheme:

“The goal should be to classify habitats to the “attribute” level, and this can be achieved with video imagery and sampling. The question is how much sampling (groundtruthing) is required, and that depends on an analysis of the sidescan and multibeam imagery and ultimately on the complexity of the seabed. Some areas will require more sampling than others. The attribute level quantifies a characteristic of the habitat (usually the

percentage of some characteristic), and this kind of data can be estimated from video imagery.”

Therefore, it seems the amount of groundtruthing (video imagery and sediment sampling) will dictate the ability of classifying habitat to the Attribute level. However, locating groundtruth samples during a seafloor mapping cruise is determined by geological features and not biological resources. Groundtruth data are not randomly collected and using groundtruth data to classify large areas of the seafloor without sufficient sampling coverage should be approached cautiously. Groundtruthing data certainly allow classification and characterization of fine-detailed seafloor habitats in areas with data, but because biological resources, such as the relative abundance of flora or fauna, are often patchy and unpredictable, sampling requires a statistically random design to describe biological patterns through space or time. Regardless of this uncertainty, groundtruthing data are often the best available data on benthic resources and should be used to classify seafloor habitat to fine-detail, as appropriate. Additional sampling of the seafloor environment may, however, be required to extrapolate between samples and classify larger areas of the seafloor.

The USGS is currently classifying habitats in the Stellwagen Bank region using Valentine et al. (2005). While this classification seems more complex with multiple choices in the Classes, Subclasses and Categories, it gets to a higher level of detail for sublittoral marine environments in New England than other schemes such as the NERRS or CMECS. Recent seafloor mapping and sampling in Massachusetts is generating detailed data and therefore, the more complex Valentine et al. (2005) approach may be appropriate. Valentine et al. (2005) is best described by reviewing the information and examples presented in Appendix 1-5 (in Valentine et al. 2005). Furthermore, results from Stellwagen Bank will help determine the detail and required data to implement the scheme. CZM should communicate with Valentine to ask about lessons learned.

Comparison of Greene et al. (1999; In Press) and Valentine et al. (2005)

Greene et al. (1999; In Press) and Valentine et al. (2005) approach habitat classification from a geologic-centric perspective and describe the use of seafloor mapping data to classify benthic habitat. Neither approach encompasses water column or intertidal environments. These studies are compared to better distinguish between the two similar approaches .

Valentine et al. (2005) compares closely to Greene et al. (1999; In Press) in that it was developed to characterize marine sublittoral (chiefly subtidal continental shelf and shelf basin) habitats predominately using seafloor mapping data. Valentine et al. (2005) is based on recent observations in northeast North America. Greene states that the Valentine et al. (2005) approach was concurrently developed with Greene et al. (1999; In Press), but Valentine et al. (2005) was specific to the East Coast and to glacial deposits where Greene et al. (1999; In Press) is more encompassing (e.g., can be used in the arctic as well as the tropics). Valentine (personal communication) describes the Greene et al. (1999; In Press) studies as using habitat as a major criterion, which the Valentine et al. (2005) does not. The Valentine et al. (2005) approach is based on describing habitats in terms of geology, oceanography, biology and related processes (e.g. sediment movement; current types; flora/fauna modifications of the seabed; etc.), designed with the idea that video imagery would be the basic data source for describing the seabed and flora/fauna (Valentine personal communication).

While this commentary helps elucidate differences between the two approaches, more clarity is needed to fully describe differences between the two. This clarity can be gained by either inviting both authors to a meeting at CZM (an action identified in the recommendations) to

discuss and provide examples of their approach in more detail, or by CZM independently, evaluating the two schemes by choosing a pilot area and implementing these two classification approaches with existing data.

RECOMMENDATIONS

The review of existing classification studies in this report is a valuable first step in determining which approach or aspects of approaches are appropriate for classifying habitats in coastal and marine environments of Massachusetts. This report also raises questions to consider when testing or adopting classification studies and offers suggestions to consider as next steps. As with any technical project plan, certain resources must be secured before work can begin. In particular, personnel, technical knowledge and supporting infrastructure need to be appropriately allocated before applying and evaluating habitat classification frameworks.

The recommendations given below are organized by the stage of the project, as follows (1) before choosing a classification scheme, (2) identifying a pilot area and (3) after evaluating classification schemes. The following recommendations are based on this study and communications with habitat classification experts and are intended to inform next steps at CZM.

Before Choosing a Classification Scheme

- CZM should assemble a technical advisory group to provide input on the choice and implementation of a classification scheme. The advisory group should be composed of scientists and managers to ensure that the approach is valid and useful in a management context. Membership in this group should include a balance of expertise with knowledge of habitat ecology, fishery resources, natural resource management, mapping technology and GIS, including staff from DMF, USGS and USEPA, among others.
- CZM and potential collaborators should become familiar with the four classification schemes recommended in this study and determine which scheme or combination of schemes include habitats that are most important to classify. This step will include identifying habitat types needed in a classification framework.
- The classification scheme chosen should support a variety of data sources, since geospatial habitat data take many forms such as grab samples and sonar images.
- The geographic coverage of a classification framework needs to be determined to facilitate the identification of appropriate studies or combination of studies needed to achieve objectives. That is – should a classification system encompass the coastal watershed and ocean environment? For example, Greene et al. (1999; In Press) and Valentine et al. (2005) do not adequately cover nearshore or coastal habitats and would need to be combined with a classification that includes these habitats (e.g., Kutcher et al. 2005 or Madden et al. 2005), if such coverage is necessary.
- Experts involved in the creation and/or implementation of the four recommended classifications should be contacted and/or gathered to ask questions, discuss applicability, solicit advice on the selection of an appropriate scheme and possibly

receive training on implementing an approach. Experts include: Nan Wright (University of Idaho), Becky Allee (NOAA's CSC), Mark Finkbeiner (NOAA's CSC) and Chris Madden (NatureServe) for CMECS (Madden et al. 2005); Tom Kutcher (Narragansett Bay NERR) and Brendan Annett (Waquoit Bay NERR) for NERRS (Kutcher et al. 2005); Gary Greene (Moss Landing Marine Laboratories) for Green et al. (1999; In Press); and Page Valentine (USGS, Woods Hole) and Kathryn Ford (DMF) for Valentine et al. (2005). This meeting should have clear objectives to facilitate the identification of next steps.

NOAA's CSC is intimately involved with CMECS (Madden et al. 2005), and CSC staff could provide assistance. Specifically, CSC can offer lessons learned from the Columbia River and other pilot projects, determine scales and data needed to populate different levels of the hierarchy, demonstrate the use of benthic data in the CMECS and train CZM and other staff on the use of the CMECS.

Evaluating a Pilot Area

- CZM should consider organizing a workshop with key partners to discuss existing habitat classification schemes, apply pre-selected classification scheme(s) to a pilot area and examine advantages and disadvantages of using one scheme over another in Massachusetts waters. Goals of this meeting could be to: 1) review existing classification schemes highlighted in this paper; 2) identify which scheme (or parts that need to be combined) best meets Massachusetts needs; 3) propose additional pilot areas to implement the classification scheme and discuss data requirements.
- The pilot area should have sufficient existing data to evaluate classification schemes. Data such as bathymetry and sediment characteristics are primary habitat attributes and important for determining the distribution and abundance of sessile invertebrates as well as motile species such as crustaceans and fish (Auster et al. 1998; Peter Auster, University of Connecticut, could advise classification efforts by CZM).
- The evaluation of habitat classification in Massachusetts should include a data gap analysis to determine data requirements to achieve specific levels of classification within each habitat classification framework.

After Evaluating Classification Schemes

- CZM should review "Recommended Guidelines for Adoption and Implementation of the NERRS Habitat and Land Use Classification System" (Walker et al. 2005) to provide a brief, practical set of guidelines for successful adoption and use of the chosen classification system. For example, Walker et al. (2005) describes the following steps: (1) identify and compile existing map resources and/or habitat description materials; (2) select appropriate scale for the habitat classification pilot; (3) design and document the GIS database structure; (4) develop a flow diagram for the implementation process; (5) document key "areas of interest" and habitats; (6) apply classification codes accordingly; and (7) generate metadata for the classification database.
- Resource management considerations should be identified (e.g., application of a classification system to address project review) and explicitly considered in choosing a classification scheme or modifying existing schemes.

- Incorporating biological and ecological characteristics into detailed, finer resolution habitat classes is not a strength in any of the four recommended classification schemes. Biotic characteristics such as the presence and relative abundance of epifaunal communities and macroalgae beds are temporally variable, as well as patchily distributed. Distributions of biological features are also greatly altered by human impacts, are a substantial part of the societal and resource value of habitats and are frequently the basis of regulatory strategies. More work is required to integrate biological data into a habitat classification framework, if resource management and/or scientific efforts require detailed, finer resolution habitat classification. The EUNIS approach to classifying biotopes (biological habitat class) may provide further guidance on incorporating biological resources into a habitat classification framework and could be reexamined while developing a classification system for Massachusetts.
- If a hybrid or modification of existing studies is selected, nomenclature must be exact and clearly constrain the meaning of terms. An official glossary of terms must be agreed upon and implemented by all users to maintain consistency in the database.
- CZM should designate one or more technical staff members to undertake future habitat classification exercises. Staff should familiarize themselves with classification schemes, mapping techniques, habitat structure and digital data management. Personnel consistency during database development is particularly important to develop and document approaches to classify habitat.
- Habitat classification scheme results should be incorporated into an existing, searchable data management system to store and disseminate data and habitat maps. CZM developed the Massachusetts Ocean Resources Information System (MORIS; a GIS database containing coastal and marine data layers), which could be used to store, view and disseminate habitat mapping products.
- CZM should consider hosting a workshop to evaluate and discuss the application of the recommended habitat classification frameworks to Massachusetts. Environmental managers, researchers, policy makers and other stakeholders involved with habitat characterization, classification and mapping should be invited to the workshop to guide identifying next steps needed to create a single habitat classification framework for Massachusetts. CZM should be mindful that revisions will most likely need to be made to the chosen classification system.
- Since habitat classification is an evolving research and management topic, a list of emerging pilot projects and studies should be maintained and coordination with overlapping activities and partners should be an ongoing process. USGS, DMF, US EPA, Massachusetts Bay National Estuary Program, Waquoit Bay and Narragansett Bay NERR and NOAA's CSC are potential partners that could be involved in discussions about the choice, modification and implementation of a classification scheme in Massachusetts.

CONCLUSION

There are many benefits of incorporating a standardized habitat classification scheme into mapping and characterization projects. This report describes 12 approaches to classify habitat, with four studies (i.e., Madden et al. 2005; Kutcher et al. 2005; Greene et al. In Press; and Valentine et al. 2005) highlighted for consideration by CZM. No one, standard classification approach is currently endorsed by the scientific and management community. In order for an approach to be successful and useful on local, regional or national scales, users need to accept a disciplined, standardized terminology to classify and describe habitats at various scales.

The selection of a singular and flexible habitat classification framework for coastal and marine environments requires a clear identification of objectives for using a scheme and rigorous testing of chosen classification(s) with existing data in a pilot area. None of the four recommended schemes sufficiently classify habitats ranging from coastal areas to oceanic environments. Evaluating the application of the four recommended schemes in coastal and marine environments in Massachusetts will facilitate a process to augment or hybridize existing schemes to achieve management objectives in Massachusetts.

Managers will ultimately benefit from a classification scheme that provides geographic representation of habitats on a map, standardized terminology for different habitat types, guidelines for organizing data, facilitation of mapping efforts, incorporation of the classification results and associated ancillary data into a well-designed GIS and improved communication. To ensure the use of a habitat classification approach, CZM must engage potential partners, including government agencies – particularly NOAA's CSC (a major partner in CMECS), academic institutions and other stakeholders.

ACKNOWLEDGMENTS

This project was the result of many discussions within the Massachusetts Office of Coastal Zone Management. Susan Snow-Cotter and Kate Killerlain Morrison provided particularly useful feedback while developing the scope of this study. The Massachusetts Division of Marine Fisheries, specifically Kathryn Ford, also helped scope the study. Authors appreciate the support from Susan Snow-Cotter and Bruce Carlisle (CZM). Anne Donovan (CZM), Dan Sampson (CZM), Rachel Franks Taylor (The Nature Conservancy) and Giancarlo Cicchetti (USEPA) provided valuable feedback on an earlier version of this report.

LITERATURE CITED

- Allee, R.J., M. Detheir, D. Brown, L. Deegan, R.G. Ford, T.F. Hourigan, J. Maragos, C. Schoch, K. Sealey, R. Twilley, M.P. Weinstein and M. Yoklavich. 2000. Marine and estuarine ecosystem and habitat classification. NOAA Technical Memorandum NMFS-F/SPO-43.
- Anderson, J.R., E.E. Hardy, J.T. Roach and R.E. Witmer. 1976. A land use and land cover classification system for use with remote sensor data. U.S. Geological Survey circular 671. United States Government Printing Office, Washington, D.C.
Available at: <http://www.ncrs.fs.fed.us/4153/deltawest/landcover/LLCoverPub.html>.
- Auster, P.J., C. Michalopoulos, P.C. Valentine and R.J. Malatesta. 1998. Delineating and monitoring habitat management units in a temperate deep-water marine protected area. Pages 169-185 in N.W.P. Munro and J.H.M. Willison, editors. Linking Protected Areas with Working Landscapes, Conserving Biodiversity. Science and Management of Protected Areas Association, Wolfville, Nova Scotia.
- Auster, P.J. 2006. Linking Seafloor Habitat Mapping Protocols to Management and Policy Needs. Funding application to US Environmental Protection Agency.
- Brown, B. 1993. A Classification System of Marine and Estuarine Habitats in Maine: An Ecosystem Approach to Habitats. Maine Natural Areas Program. Department of Economic and Community Development. 51 pages + appendices. Augusta, ME.
- Brown, S.K. 2002. Our living Oceans Benthic Habitat Classification System. NOAA NMFS Office of Science and Technology.
- Connor, D.W., J. Allen, N. Golding, K. Howell, L. Lieberknecht, K. Northen and J. Reker. 2004. The national marine habitat classification for Britain and Ireland, version 04.05. Joint Nature Conservation Committee. Available at: http://www.jncc.gov.uk/pdf/04_05_introduction.pdf .
- Cowardin, L.M., V. Carter, F.C. Golet and E.T. LaRoe. 1979. Classification of wetlands and deepwater habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31. Available at: http://www.fws.gov/nwi/Pubs_Reports/Class_Manual/class_titlepg.htm.
- Davies, C., D. Moss and M.O. Hill. 2004. EUNIS Habitat Classification. European Environment Agency.
- Dethier, M.N. 1992. Classifying marine and estuarine natural communities: an alternative to the Cowardin system. *Natural Areas Journal* 12(2):90-100.
- Dobson, J.E., E.A. Bright, R.L. Ferguson, D.W. Field, L.L. Wood, K.D. Haddad, H. Iredale, J.R. Jenson, V.V. Klemas, R.J. Orth and J.P. Thomas. 1995. NOAA Coastal Change Analysis Program (C-CAP): Guidance for Regional Implementation. NOAA Technical Report NMFS 123. Available at: <http://www.csc.noaa.gov/crs/lca/pdf/protocol.pdf>.
- European Environment Agency. 1999. European Topic Centre on Nature Conservation. Work Programme: Task 4.3 EUNIS Habitat Classification, Draft Final Report. 209 pp. Available at: <http://www.searchmesh.net/>.
- Greene, H.G., M.M. Yoklavich, R.M. Starr, V.M. O'Connell, W.W. Wakefield, D.E. Sullivan, J.E. McRea Jr. and G.M. Caillet. 1999. A classification scheme for deep seafloor habitats. *Oceanologica Acta* 22:663-678.
- Greene, H.G., J.J. Bizzarro, J.E. Tilden, H.L. Lopez and M.D. Erdey. 2005. The benefits and pitfalls of geographic information systems in marine benthic habitat mapping. Pages 34-46, in Wright, D.J. and Scholz, A.J., eds. *Place Matters*. Oregon State University Press, Portland, OR.

- Greene, H.G., J.J. Bizzarro, V. O'Connell and C.K. Brylinsky. In Press. Construction of digital potential marine benthic habitat maps using a coded classification scheme and their application. In H.G. Greene and B.J. Todd (eds.) , Mapping the Seafloor for Habitat Characterization. Geological Association of Canada, Special Paper 47.
- Kurtz, J.C., N.D. Detenbeck, V.D. Engle, K. Ho, L.M. Smith, S.J. Jordan and D. Cambell. 2006. Classifying coastal waters: current necessity and historical perspective. *Estuaries and Coasts* 29(1):107-123.
- Kutcher, T.E., N.H. Garfield and K.B. Raposa. 2005 (draft). A recommendation for a comprehensive habitat and land use classification system for the National Estuarine Research Reserve System. National Estuarine Research Reserve, Estuarine Reserves Division. Draft report to NOAA/NOS/OCRM. Silver Spring, MD. 26 pp.
- Kutcher, T.E. 2006. A comparison of functionality between two coastal classification schemes developed within the National Oceanic and Atmospheric Administration. NERRS/ERD Proposal (report to NOAA/NOS/OCRM), Silver Spring, MD, 5 pp.
- Madden, C.J. and D.H. Grossman. 2004a. A framework for a coastal/marine ecological classification standard. NatureServe, Arlington, VA. Prepared for the National Oceanic and Atmospheric Administration, Under Contract EA-133C-03-SE-0275.
- Madden, C.J., and D.H. Grossman. 2004b. National coastal/marine classification pilot projects. Draft final report. NatureServe, Arlington, VA.
- Madden, C.J., D.H. Grossman and K.L. Goodin. 2005. Coastal and Marine Systems of North America: Framework for an Ecological Classification Standard: Version II. NatureServe, Arlington, VA. Available at: http://www.natureserve.org/getData/CMECS/cm_pub.pdf
- Madley, K. 2002. Florida system for classification of habitats in estuarine and marine environments (SCHEME). Report to US Environmental Protection Agency. Florida Fish and Wildlife Conservation Commissions. Florida Marine Research Institute. FMRI File Cod 2277-00-02-F. 43 pp.
- Massachusetts Office of Coastal Zone Management (CZM). 2004-2005. Coastlines – Winter 2004-2005. Annual publication of the Massachusetts Office of Coastal Zone Management. Boston, MA. Available at: <http://www.mass.gov/czm/coastlines/2004-2005/index04-05.htm>.
- National Estuarine Research Reserve System (NERRS). 2000. Evaluations of marine and estuarine ecosystem and habitat classification. NOAA Technical Memorandum NMFS-F/SPO-43.
- Tierney, S. (Chair of Massachusetts Ocean Management Task Force). 2004. Waves of Change: The Massachusetts Ocean Management Task Force Report and Recommendations. Report submitted to Ellen Roy Herzfelder, The Commonwealth of Massachusetts Secretary of Environmental Affairs. Boston, MA.
- Tyrrell, M.C. 2004. Strategic plan for mapping Massachusetts' benthic marine habitats. Massachusetts Office of Coastal Zone Management, Boston, MA.
- Tyrrell, M.C. 2005. Gulf of Maine Marine Habitat Primer. Gulf of Maine Council on the Marine Environment. 54pp. Available at : <http://www.gulfofmaine.org/habitatprimer>.
- Valentine, P.C., B.J. Todd and V.E. Kostylev. 2005. Classification of marine sublittoral habitats, with application to the northeastern North America region. *American Fisheries Society Symposium* 41:183-200. Available at: <http://woodshole.er.usgs.gov/project-pages/stellwagen/index.html>.

Walker, S.P., N.H. Garfield, T.E. Kutcher and T. Reed. 2005. Recommended guidelines for adoption and implementation of the NERRS comprehensive habitat and land use classification system. Technical document produced for pilot National Estuarine Research Reserves.

Wilbur, A.R. and M.W. Pentony. 1999. Human-induced nonfishing threats to Essential Fish Habitat in the New England region. In, L. Benaka, editor. Fish Habitat: Essential Fish Habitat and Rehabilitation. American Fisheries Society Symposium Proceedings 22:299-321.

PERSONAL COMMUNICATION

Allee, R.J. 2006. Personal communication. National Oceanic and Atmospheric Administration. Silver Springs, MD.

Annett, B. 2006. Personal communication. Waquoit Bay National Estuarine Research Reserve. Waquoit, MA.

Dethier, M. 2006. Personal communication. University of Washington and Friday Harbors Labs. Friday Harbors, WA.

Greene, H.G. 2006. Personal communication. United States Geological Survey. Moss Landing Marine Laboratories. Moss Landing, CA.

Kutcher, T.E. 2006. Personal communication. Narragansett Bay National Estuarine Research Reserve. Prudence Island, RI.

Wright, N. 2006. Personal communication. Department of Fish and Wildlife Resources, University of Idaho. Moscow, ID.

APPENDICES

APPENDIX 1: Classification hierarchy / organization of four recommended habitat classification studies.

- A.** Madden et al. (2005), "Coastal and Marine Systems of North America – Framework for an Ecological Classification Standard: Version II"
- B.** Kutcher et al. 2005, "A recommendation for a comprehensive habitat and land use classification system for the National Estuarine Research Reserve System"
- C.** Greene et. al (In Press), "Construction of digital potential marine benthic habitat maps using a coded classification scheme and their application"
- D.** Valentine et al. (2005), "Classification of Marine Sublittoral Habitats, with Application to the Northeastern North America Region"

A. Madden et al. (2005), “Coastal and Marine Systems of North America – Framework for an Ecological Classification Standard: Version II”

Contact: Chris Madden, cmadden@sfwmd.gov, NatureServe

Website: <http://www.natureserve.org/getData/CMECS/>

Level 1 Level 2 Level 3 Level 4 Level 5 (Habitat)

Regime	Formation	Zone_Suffix	Macrohabitat_Suffix	Macrohab	Unit_id	Unit_Name_Suffix	Unit_Name	
A	Estuarine	1 Lagoon	WU Water Column upper layer	a	estuarine water column	A.01.WU.a	501 estuarine turbidity maximum	Estuarine Lagoon Upper water column Turbidity Maximum
A	Estuarine	1 Lagoon	WU Water Column upper layer	a	estuarine water column	A.01.WU.a	502 pycnocline	Estuarine Lagoon Pycnocline
A	Estuarine	1 Lagoon	WU Water Column upper layer	a	estuarine water column	A.01.WU.a	503 upper water column	Estuarine Lagoon surface foam
A	Estuarine	1 Lagoon	WU Water Column upper layer	a	estuarine water column	A.01.WU.a	504 small fresh water lens	Estuarine Lagoon Freshwater Lens
A	Estuarine	1 Lagoon	WU Water Column upper layer	a	estuarine water column	A.01.WU.a	505 hyperhaline estuarine water column	Estuarine Lagoon Hypersaline Water
A	Estuarine	1 Lagoon	WU Water Column upper layer	a	estuarine water column	A.01.WU.a	506 phytoplankton bloom	Estuarine Lagoon Phytoplankton Bloom
A	Estuarine	1 Lagoon	WU Water Column upper layer	a	estuarine water column	A.01.WU.a	507 epipelagic zone	Estuarine Lagoon Epipelagic Layer
A	Estuarine	1 Lagoon	WU Water Column upper layer	a	estuarine water column	A.01.WU.a	508 floating vegetation mat	Estuarine Lagoon Floating Vegetation Mat
A	Estuarine	1 Lagoon	WU Water Column upper layer	a	estuarine water column	A.01.WU.a	509 tributary discharge zone	Estuarine Lagoon Tributary Discharge
A	Estuarine	1 Lagoon	WU Water Column upper layer	a	estuarine water column	A.01.WU.a	510 counter current	Estuarine Lagoon Counter Current Zone
A	Estuarine	1 Lagoon	WB Water Column bottom layer	a	estuarine water column	A.01.WB.a	511 anoxic bottom water	Estuarine Lagoon Anoxic Bottom Water
A	Estuarine	1 Lagoon	WB Water Column bottom layer	a	estuarine water column	A.01.WB.a	512 oxix bottom water	Estuarine Lagoon Oxix Bottom Water
A	Estuarine	1 Lagoon	WB Water Column bottom layer	a	estuarine water column	A.01.WB.a	513 salt wedge	Estuarine Lagoon Salt Wedge
A	Estuarine	1 Lagoon	WB Water Column bottom layer	a	estuarine water column	A.01.WB.a	514 groundwater seep	Estuarine Lagoon Groundwater Seep
A	Estuarine	1 Lagoon	WB Water Column bottom layer	a	estuarine water column	A.01.WB.a	515 benthic boundary layer	Estuarine Lagoon Benthic Boundary Layer
A	Estuarine	1 Lagoon	LS Littoral Supratidal	a	unconsolidated sediments	A.01.LS.a	516 bare organic mud	Estuarine Lagoon Bare Organic Sediment
A	Estuarine	1 Lagoon	LS Littoral Supratidal	a	unconsolidated sediments	A.01.LS.a	517 bare carbonate sediment	Estuarine Lagoon Bare Carbonate Sediment
A	Estuarine	1 Lagoon	LS Littoral Supratidal	a	unconsolidated sediments	A.01.LS.a	518 bare carbonate mud	Estuarine Lagoon Bare Carbonate Mud
A	Estuarine	1 Lagoon	LS Littoral Supratidal	a	unconsolidated sediments	A.01.LS.a	519 bare carbonate sand	Estuarine Lagoon Bare Carbonate Sands
A	Estuarine	1 Lagoon	LS Littoral Supratidal	a	unconsolidated sediments	A.01.LS.a	520 bare mixed-coarse sediment softbottom	Estuarine Lagoon Bare Mixed-Coarse Sediment
A	Estuarine	1 Lagoon	LS Littoral Supratidal	a	unconsolidated sediments	A.01.LS.a	521 vegetated mineral sediments	Estuarine Lagoon Vegetated Mineral Sediments
A	Estuarine	1 Lagoon	LS Littoral Supratidal	a	unconsolidated sediments	A.01.LS.a	522 vegetated organic sediments	Estuarine Lagoon Vegetated Organic Sediments
A	Estuarine	1 Lagoon	LS Littoral Supratidal	b	rocky shore	A.01.LS.b	523 bedrock shore	Estuarine Lagoon Bedrock Shore
A	Estuarine	1 Lagoon	LS Littoral Supratidal	b	rocky shore	A.01.LS.b	524 hardpan shore	Estuarine Lagoon Hardpan Shore
A	Estuarine	1 Lagoon	LS Littoral Supratidal	b	rocky shore	A.01.LS.b	525 boulder shore	Estuarine Lagoon Boulder Shore
A	Estuarine	1 Lagoon	LS Littoral Supratidal	b	rocky shore	A.01.LS.b	526 cobble shore	Estuarine Lagoon Cobble Shore
A	Estuarine	1 Lagoon	LS Littoral Supratidal	b	rocky shore	A.01.LS.b	527 sand-bedrock shore	Estuarine Lagoon Sand-Bedrock Shore
A	Estuarine	1 Lagoon	LS Littoral Supratidal	c	dune system	A.01.LS.c	528 foredune	Estuarine Lagoon Foredune
A	Estuarine	1 Lagoon	LS Littoral Supratidal	c	dune system	A.01.LS.c	529 dune crest	Estuarine Lagoon Dune Crest

A	Estuarine	1 Lagoon	LS	Littoral Supratidal	c	dune system	A.01.LS.c	530	backdune	Estuarine Lagoon Backdune
A	Estuarine	1 Lagoon	LS	Littoral Supratidal	d	cliff	A.01.LS.d	531	cliff notch	Estuarine Lagoon Cliff Notch
A	Estuarine	1 Lagoon	LS	Littoral Supratidal	d	cliff	A.01.LS.d	532	cliff bioerosion notch	Estuarine Lagoon Bioeroded Cliff Notch
A	Estuarine	1 Lagoon	LS	Littoral Supratidal	d	cliff	A.01.LS.d	533	cliff cave	Estuarine Lagoon Cliff Cave
A	Estuarine	1 Lagoon	LS	Littoral Supratidal	d	cliff	A.01.LS.d	534	cliff fracture	Estuarine Lagoon Cliff Fracture
A	Estuarine	1 Lagoon	LS	Littoral Supratidal	d	cliff	A.01.LS.d	535	cliff rubble zone	Estuarine Lagoon Cliff Rubble Zone
A	Estuarine	1 Lagoon	LS	Littoral Supratidal	e	mangrove swamp	A.01.LS.e	536	prop root zone	Estuarine Lagoon Mangrove Prop Root Zone
A	Estuarine	1 Lagoon	LS	Littoral Supratidal	e	mangrove swamp	A.01.LS.e	537	basin swamp	Estuarine Lagoon Mangrove Basin
A	Estuarine	1 Lagoon	LS	Littoral Supratidal	e	mangrove swamp	A.01.LS.e	538	buttonwood ridge	Estuarine Lagoon Mangrove Buttonwood Ridge
A	Estuarine	1 Lagoon	LS	Littoral Supratidal	e	mangrove swamp	A.01.LS.e	539	pneumatophore zone	Estuarine Lagoon Mangrove Pneumatophore Zone
A	Estuarine	1 Lagoon	LS	Littoral Supratidal	e	mangrove swamp	A.01.LS.e	540	swamp creek	Estuarine Lagoon Mangrove Swamp Creek
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	a	unconsolidated sediments	A.01.LI.a	541	bare organic mud softbottom	Estuarine Lagoon Intertidal Bare Organic Mud sediments
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	a	unconsolidated sediments	A.01.LI.a	542	bare carbonate sediment softbottom	Estuarine Lagoon Intertidal Bare Carbonate Sediment sediments
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	a	unconsolidated sediments	A.01.LI.a	543	bare carbonate mud softbottom	Estuarine Lagoon Intertidal Bare Carbonate Mud sediments
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	a	unconsolidated sediments	A.01.LI.a	544	bare carbonate sand softbottom	Estuarine Lagoon Intertidal Bare Carbonate Sand sediments
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	a	unconsolidated sediments	A.01.LI.a	545	bare mixed-coarse sediment softbottom	Estuarine Lagoon Intertidal Bare Mixed-coarse Sediment sediments
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	a	unconsolidated sediments	A.01.LI.a	546	vegetated softbottom- holdfast (macroalgae be	Estuarine Lagoon Intertidal Vegetated Softbottom- non-rooted
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	a	unconsolidated sediments	A.01.LI.a	547	vegetated softbottom- rooted (seagrass bed)	Estuarine Lagoon Intertidal Vegetated Softbottom- rooted
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	a	unconsolidated sediments	A.01.LI.a	548	vegetated organic softbottom	Estuarine Lagoon Intertidal Vegetated Organic Softbottom
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	b	mudflat	A.01.LI.b	549	fringing vegetated brackish mudflat	Estuarine Lagoon Intertidal Fringing Vegetated Brackish Mudflat
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	b	mudflat	A.01.LI.b	550	fringing vegetated saline mudflat	Estuarine Lagoon Intertidal Fringing Vegetated saline Mudflat
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	b	mudflat	A.01.LI.b	551	bare brackish mudflat	Estuarine Lagoon Intertidal Bare Brackish Mudflat
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	b	mudflat	A.01.LI.b	552	bare saline mudflat	Estuarine Lagoon Intertidal Bare Saline Mudflat
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	c	saline fringing wetland (marsh)	A.01.LI.c	553	saline fringing wetland tidal pass	Estuarine Lagoon Intertidal saline marsh tidal pass
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	c	saline fringing wetland (marsh)	A.01.LI.c	554	saline fringing wetland tidal creek	Estuarine Lagoon Intertidal saline marsh tidal creek
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	c	saline fringing wetland (marsh)	A.01.LI.c	555	saline fringing wetland tidal creek bank	Estuarine Lagoon Intertidal saline marsh tidal creek bank
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	c	saline fringing wetland (marsh)	A.01.LI.c	556	saline wetland hammock	Estuarine Lagoon Intertidal saline marsh hammock
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	c	saline fringing wetland (marsh)	A.01.LI.c	557	saline wetland inland basin marsh	Estuarine Lagoon Intertidal saline marsh basin
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	c	saline fringing wetland (marsh)	A.01.LI.c	558	saline wetland streamside marsh	Estuarine Lagoon Intertidal saline streamside marsh
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	d	brackish fringing wetland (marsh)	A.01.LI.d	559	brackish fringing wetland tidal pass	Estuarine Lagoon Intertidal brackish marsh tidal pass
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	d	brackish fringing wetland (marsh)	A.01.LI.d	560	brackish fringing wetland tidal creek	Estuarine Lagoon Intertidal brackish marsh tidal creek
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	d	brackish fringing wetland (marsh)	A.01.LI.d	561	brackish fringing wetland tidal creek bank	Estuarine Lagoon Intertidal brackish marsh tidal creek bank
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	d	brackish fringing wetland (marsh)	A.01.LI.d	562	brackish wetland hammock	Estuarine Lagoon Intertidal brackish marsh hammock
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	d	brackish fringing wetland (marsh)	A.01.LI.d	563	brackish wetland inland basin marsh	Estuarine Lagoon Intertidal brackish marsh basin
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	d	brackish fringing wetland (marsh)	A.01.LI.d	564	brackish wetland streamside marsh	Estuarine Lagoon Intertidal brackish streamside marsh
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	e	hardbottom	A.01.LI.e	565	bare gravel hardbottom	Estuarine Lagoon Intertidal bare gravel
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	e	hardbottom	A.01.LI.e	566	bare limestone pavement hardbottom	Estuarine Lagoon Intertidal bare limestone pavement
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	e	hardbottom	A.01.LI.e	567	bare bedrock hardbottom	Estuarine Lagoon Intertidal bare bedrock

A	Estuarine	1 Lagoon	LI	Littoral Intertidal	f cliff	A.01.LI.f	568 notch	Estuarine Lagoon Intertidal cliff notch
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	f cliff	A.01.LI.f	569 bioerosion notch	Estuarine Lagoon Intertidal Bioeroded cliff notch
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	f cliff	A.01.LI.f	570 cave	Estuarine Lagoon Intertidal cliff cave
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	f cliff	A.01.LI.f	571 fracture	Estuarine Lagoon Intertidal cliff fracture
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	f cliff	A.01.LI.f	572 rubble zone	Estuarine Lagoon Intertidal cliff rubble zone
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	g mangrove swamp	A.01.LI.g	573 prop root zone	Estuarine Lagoon Intertidal mangrove prop root zone
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	g mangrove swamp	A.01.LI.g	574 basin swamp	Estuarine Lagoon Intertidal mangrove basin swamp
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	g mangrove swamp	A.01.LI.g	575 buttonwood ridge	Estuarine Lagoon Intertidal mangrove buttonwood ridge
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	g mangrove swamp	A.01.LI.g	576 pneumatophore zone	Estuarine Lagoon Intertidal mangrove pneumatophore zone
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	g mangrove swamp	A.01.LI.g	577 swamp creek	Estuarine Lagoon Intertidal mangrove creek
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	g mangrove swamp	A.01.LI.g	578 basin pond	Estuarine Lagoon Intertidal mangrove basin pond
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	h coastal beach	A.01.LI.h	579 sand beach	Estuarine Lagoon Intertidal sand beach
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	h coastal beach	A.01.LI.h	580 mixed-fine sediment beach	Estuarine Lagoon Intertidal mixed-fine sediment beach
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	h coastal beach	A.01.LI.h	581 mixed-fine sand and mud beach	Estuarine Lagoon Intertidal mixed-fine sand and mud beach
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	h coastal beach	A.01.LI.h	582 mud beach	Estuarine Lagoon Intertidal mud beach
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	h coastal beach	A.01.LI.h	583 rock and boulder beach	Estuarine Lagoon Intertidal rock and boulder beach
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	h coastal beach	A.01.LI.h	584 cobble beach	Estuarine Lagoon Intertidal cobble beach
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	h coastal beach	A.01.LI.h	585 mixed-coarse sediment beach	Estuarine Lagoon Intertidal mixed-coarse sediment beach
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	h coastal beach	A.01.LI.h	586 gravel beach	Estuarine Lagoon Intertidal gravel beach
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	i rocky shore	A.01.LI.i	587 bedrock shore	Estuarine Lagoon Intertidal bedrock shore
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	i rocky shore	A.01.LI.i	588 hardpan shore	Estuarine Lagoon Intertidal hardpan shore
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	i rocky shore	A.01.LI.i	589 boulder shore	Estuarine Lagoon Intertidal boulder shore
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	i rocky shore	A.01.LI.i	590 cobble shore	Estuarine Lagoon Intertidal cobble shore
A	Estuarine	1 Lagoon	LI	Littoral Intertidal	i rocky shore	A.01.LI.i	591 sand-bedrock shore	Estuarine Lagoon Intertidal sand-bedrock shore
A	Estuarine	1 Lagoon	B	Bottom	a oyster reef	A.01.B.a	592 live oyster reef	Estuarine Lagoon Subtidal oyster reef
A	Estuarine	1 Lagoon	B	Bottom	a oyster reef	A.01.B.a	593 relict oyster reef	Estuarine Lagoon Subtidal relict oyster reef
A	Estuarine	1 Lagoon	B	Bottom	a oyster reef	A.01.B.a	595 periodically emergent oyster reef	Estuarine Lagoon Intertidal oyster reef
A	Estuarine	1 Lagoon	B	Bottom	a oyster reef	A.01.B.a	596 oyster shell midden	Estuarine Lagoon Subtidal oyster shell midden
A	Estuarine	1 Lagoon	B	Bottom	b worm reef	A.01.B.b	597 live worm reef	Estuarine Lagoon Subtidal worm reef
A	Estuarine	1 Lagoon	B	Bottom	b worm reef	A.01.B.b	598 relict worm reef	Estuarine Lagoon Subtidal relict worm reef
A	Estuarine	1 Lagoon	B	Bottom	b worm reef	A.01.B.b	599 live mussel reef	Estuarine Lagoon Subtidal mussel reef
A	Estuarine	1 Lagoon	B	Bottom	b worm reef	A.01.B.b	600 relict mussel reef	Estuarine Lagoon Subtidal relict mussel reef
A	Estuarine	1 Lagoon	B	Bottom	c unconsolidated sediments	A.01.B.c	601 bare sandy softbottom	Estuarine Lagoon Subtidal bare sandy sediments
A	Estuarine	1 Lagoon	B	Bottom	c unconsolidated sediments	A.01.B.c	602 bare mineral mud softbottom	Estuarine Lagoon Subtidal bare mineral mud sediments
A	Estuarine	1 Lagoon	B	Bottom	c unconsolidated sediments	A.01.B.c	603 bare organic mud softbottom	Estuarine Lagoon Subtidal bare organic mud sediments
A	Estuarine	1 Lagoon	B	Bottom	c unconsolidated sediments	A.01.B.c	604 bare carbonate mud-shell hash softbottom	Estuarine Lagoon Subtidal bare carbonate mud-shell hash sediments
A	Estuarine	1 Lagoon	B	Bottom	c unconsolidated sediments	A.01.B.c	605 bare carbonate mud softbottom	Estuarine Lagoon Subtidal bare carbonate mud sediments
A	Estuarine	1 Lagoon	B	Bottom	c unconsolidated sediments	A.01.B.c	606 bare carbonate sand softbottom	Estuarine Lagoon Subtidal bare carbonate sand sediments
A	Estuarine	1 Lagoon	B	Bottom	c unconsolidated sediments	A.01.B.c	607 bare mixed-coarse sediment softbottom	Estuarine Lagoon Subtidal bare mixed-coarse sediment sediments

A	Estuarine	1 Lagoon	B	Bottom	c	unconsolidated sediments	A.01.B.c	608	colonized mixed-fine softbottom	Estuarine Lagoon Subtidal colonized mixed-fine sediments
A	Estuarine	1 Lagoon	B	Bottom	c	unconsolidated sediments	A.01.B.c	609	vegetated mud and mixed-fine softbottom	Estuarine Lagoon Subtidal vegetated mud and mixed-fine sediments
A	Estuarine	1 Lagoon	B	Bottom	c	unconsolidated sediments	A.01.B.c	610	bare gravel softbottom	Estuarine Lagoon Subtidal bare gravel
A	Estuarine	1 Lagoon	B	Bottom	c	unconsolidated sediments	A.01.B.c	611	vegetated softbottom	Estuarine Lagoon Subtidal vegetated softbottom
A	Estuarine	1 Lagoon	B	Bottom	c	unconsolidated sediments	A.01.B.c	612	colonized softbottom	Estuarine Lagoon Subtidal colonized softbottom
A	Estuarine	1 Lagoon	B	Bottom	c	unconsolidated sediments	A.01.B.c	613	vegetated colonized softbottom	Estuarine Lagoon Subtidal vegetated colonized softbottom
A	Estuarine	1 Lagoon	B	Bottom	d	hardbottom	A.01.B.d	614	bare limestone pavement hardbottom	Estuarine Lagoon Subtidal bare limestone pavement
A	Estuarine	1 Lagoon	B	Bottom	d	hardbottom	A.01.B.d	615	vegetated rock and boulder hardbottom	Estuarine Lagoon Subtidal vegetated rock and boulder
A	Estuarine	1 Lagoon	B	Bottom	d	hardbottom	A.01.B.d	616	cobble hardbottom	Estuarine Lagoon Subtidal cobble
A	Estuarine	1 Lagoon	B	Bottom	d	hardbottom	A.01.B.d	617	gravel hardbottom	Estuarine Lagoon Subtidal gravel
A	Estuarine	1 Lagoon	B	Bottom	d	hardbottom	A.01.B.d	618	boulder hardbottom	Estuarine Lagoon Subtidal boulder
A	Estuarine	1 Lagoon	B	Bottom	d	hardbottom	A.01.B.d	619	bedrock hardbottom	Estuarine Lagoon Subtidal bedrock
A	Estuarine	1 Lagoon	B	Bottom	d	hardbottom	A.01.B.d	620	colonized tidal creek bottom	Estuarine Lagoon Subtidal colonized tidal creek bottom
A	Estuarine	2 Embayment	WU	Water Column (upper layer	a	estuarine water column	A.02.WU.a	621	estuarine turbidity maximum	Estuarine Embayment Turbidity Maximum
A	Estuarine	2 Embayment	WU	Water Column (upper layer	a	estuarine water column	A.02.WU.a	622	pycnocline	Estuarine Embayment Pycnocline
A	Estuarine	2 Embayment	WU	Water Column (upper layer	a	estuarine water column	A.02.WU.a	623	upper water column	Estuarine Embayment surface foam
A	Estuarine	2 Embayment	WU	Water Column (upper layer	a	estuarine water column	A.02.WU.a	624	small fresh water lens	Estuarine Embayment Freshwater Lens
A	Estuarine	2 Embayment	WU	Water Column (upper layer	a	estuarine water column	A.02.WU.a	625	hyperhaline estuarine water column	Estuarine Embayment Hypersaline Water
A	Estuarine	2 Embayment	WU	Water Column (upper layer	a	estuarine water column	A.02.WU.a	626	phytoplankton bloom	Estuarine Embayment Phytoplankton Bloom
A	Estuarine	2 Embayment	WU	Water Column (upper layer	a	estuarine water column	A.02.WU.a	627	epipelagic zone	Estuarine Embayment Epipelagic Layer
A	Estuarine	2 Embayment	WU	Water Column (upper layer	a	estuarine water column	A.02.WU.a	628	floating vegetation mat	Estuarine Embayment Floating Vegetation Mat
A	Estuarine	2 Embayment	WU	Water Column (upper layer	a	estuarine water column	A.02.WU.a	629	tributary discharge zone	Estuarine Embayment Tributary Discharge
A	Estuarine	2 Embayment	WB	Water Column (bottom laye	a	estuarine water column	A.02.WB.a	630	counter current	Estuarine Embayment Counter Current Zone
A	Estuarine	2 Embayment	WB	Water Column (bottom laye	a	estuarine water column	A.02.WB.a	631	anoxic bottom water	Estuarine Embayment Anoxic Bottom Water
A	Estuarine	2 Embayment	WB	Water Column (bottom laye	a	estuarine water column	A.02.WB.a	632	oxic bottom water	Estuarine Embayment Oxic Bottom Water
A	Estuarine	2 Embayment	WB	Water Column (bottom laye	a	estuarine water column	A.02.WB.a	633	salt wedge	Estuarine Embayment Salt Wedge
A	Estuarine	2 Embayment	WB	Water Column (bottom laye	a	estuarine water column	A.02.WB.a	634	groundwater seep	Estuarine Embayment Groundwater Seep
A	Estuarine	2 Embayment	WB	Water Column (bottom laye	a	estuarine water column	A.02.WB.a	635	benthic boundary layer	Estuarine Embayment Benthic Boundary Layer
A	Estuarine	2 Embayment	LS	Littoral Supratidal	a	unconsolidated sediments	A.02.LS.a	636	bare organic mud	Estuarine Embayment Bare Organic Sediment
A	Estuarine	2 Embayment	LS	Littoral Supratidal	a	unconsolidated sediments	A.02.LS.a	637	bare carbonate sediment	Estuarine Embayment Bare Carbonate Sediment
A	Estuarine	2 Embayment	LS	Littoral Supratidal	a	unconsolidated sediments	A.02.LS.a	638	bare carbonate mud	Estuarine Embayment Bare Carbonate Mud
A	Estuarine	2 Embayment	LS	Littoral Supratidal	a	unconsolidated sediments	A.02.LS.a	639	bare carbonate sand	Estuarine Embayment Bare Carbonate Sand
A	Estuarine	2 Embayment	LS	Littoral Supratidal	a	unconsolidated sediments	A.02.LS.a	640	bare mixed-coarse sediment softbottom	Estuarine Embayment Bare Mixed-Coarse Sediment
A	Estuarine	2 Embayment	LS	Littoral Supratidal	a	unconsolidated sediments	A.02.LS.a	641	vegetated mineral sediments	Estuarine Embayment Vegetated Mineral Sediments
A	Estuarine	2 Embayment	LS	Littoral Supratidal	a	unconsolidated sediments	A.02.LS.a	642	vegetated organic sediments	Estuarine Embayment Vegetated Organic Sediments
A	Estuarine	2 Embayment	LS	Littoral Supratidal	b	rocky shore	A.02.LS.b	643	bedrock shore	Estuarine Embayment Bedrock Shore
A	Estuarine	2 Embayment	LS	Littoral Supratidal	b	rocky shore	A.02.LS.b	644	hardpan shore	Estuarine Embayment Hardpan Shore
A	Estuarine	2 Embayment	LS	Littoral Supratidal	b	rocky shore	A.02.LS.b	645	boulder shore	Estuarine Embayment Boulder Shore

A	Estuarine	2 Embayment	LS	Littoral Supratidal	b	rocky shore	A.02.LS.b	646	cobble shore	Estuarine Embayment Cobble Shore
A	Estuarine	2 Embayment	LS	Littoral Supratidal	b	rocky shore	A.02.LS.b	647	sand-bedrock shore	Estuarine Embayment Sand-Bedrock Shore
A	Estuarine	2 Embayment	LS	Littoral Supratidal	c	dune system	A.02.LS.c	648	foredune	Estuarine Embayment Foredune
A	Estuarine	2 Embayment	LS	Littoral Supratidal	c	dune system	A.02.LS.c	649	dune crest	Estuarine Embayment Dune Crest
A	Estuarine	2 Embayment	LS	Littoral Supratidal	c	dune system	A.02.LS.c	650	backdune	Estuarine Embayment Backdune
A	Estuarine	2 Embayment	LS	Littoral Supratidal	d	cliff	A.02.LS.d	651	cliff notch	Estuarine Embayment Cliff Notch
A	Estuarine	2 Embayment	LS	Littoral Supratidal	d	cliff	A.02.LS.d	652	cliff bioerosion notch	Estuarine Embayment Bioeroded Cliff Notch
A	Estuarine	2 Embayment	LS	Littoral Supratidal	d	cliff	A.02.LS.d	653	cliff cave	Estuarine Embayment Cliff Cave
A	Estuarine	2 Embayment	LS	Littoral Supratidal	d	cliff	A.02.LS.d	654	cliff fracture	Estuarine Embayment Cliff Fracture
A	Estuarine	2 Embayment	LS	Littoral Supratidal	d	cliff	A.02.LS.d	655	cliff rubble zone	Estuarine Embayment Cliff Rubble Zone
A	Estuarine	2 Embayment	LS	Littoral Supratidal	e	mangrove swamp	A.02.LS.e	656	prop root zone	Estuarine Embayment Mangrove Prop Root Zone
A	Estuarine	2 Embayment	LS	Littoral Supratidal	e	mangrove swamp	A.02.LS.e	657	basin swamp	Estuarine Embayment Mangrove Basin
A	Estuarine	2 Embayment	LS	Littoral Supratidal	e	mangrove swamp	A.02.LS.e	658	buttonwood ridge	Estuarine Embayment Mangrove Buttonwood Ridge
A	Estuarine	2 Embayment	LS	Littoral Supratidal	e	mangrove swamp	A.02.LS.e	659	pneumatophore zone	Estuarine Embayment Mangrove Pneumatophore Zone
A	Estuarine	2 Embayment	LS	Littoral Supratidal	e	mangrove swamp	A.02.LS.e	660	swamp creek	Estuarine Embayment Mangrove Swamp Creek
A	Estuarine	2 Embayment	LI	Littoral Intertidal	a	unconsolidated sediments	A.02.LI.a	661	bare organic mud softbottom	Estuarine Embayment Intertidal Bare Organic Mud Sediments
A	Estuarine	2 Embayment	LI	Littoral Intertidal	a	unconsolidated sediments	A.02.LI.a	662	bare carbonate sediment softbottom	Estuarine Embayment Intertidal Bare Carbonate Sediments
A	Estuarine	2 Embayment	LI	Littoral Intertidal	a	unconsolidated sediments	A.02.LI.a	663	bare carbonate mud softbottom	Estuarine Embayment Intertidal Bare Carbonate Mud Sediments
A	Estuarine	2 Embayment	LI	Littoral Intertidal	a	unconsolidated sediments	A.02.LI.a	664	bare carbonate sand softbottom	Estuarine Embayment Intertidal Bare Carbonate Sand Sediments
A	Estuarine	2 Embayment	LI	Littoral Intertidal	a	unconsolidated sediments	A.02.LI.a	665	bare mixed-coarse sediment softbottom	Estuarine Embayment Intertidal Bare Mixed-coarse Sediments
A	Estuarine	2 Embayment	LI	Littoral Intertidal	a	unconsolidated sediments	A.02.LI.a	666	vegetated softbottom- holdfast (macroalgae be	Estuarine Embayment Intertidal Vegetated Softbottom- non-rooted
A	Estuarine	2 Embayment	LI	Littoral Intertidal	a	unconsolidated sediments	A.02.LI.a	667	vegetated softbottom- rooted (seagrass bed)	Estuarine Embayment Intertidal Vegetated Softbottom- rooted
A	Estuarine	2 Embayment	LI	Littoral Intertidal	a	unconsolidated sediments	A.02.LI.a	668	vegetated organic softbottom	Estuarine Embayment Intertidal Vegetated Organic Softbottom
A	Estuarine	2 Embayment	LI	Littoral Intertidal	b	mudflat	A.02.LI.b	669	fringing vegetated brackish mudflat	Estuarine Embayment Intertidal Fringing Vegetated Brackish Mudflat
A	Estuarine	2 Embayment	LI	Littoral Intertidal	b	mudflat	A.02.LI.b	670	fringing vegetated saline mudflat	Estuarine Embayment Intertidal Fringing Vegetated Saline Mudflat
A	Estuarine	2 Embayment	LI	Littoral Intertidal	b	mudflat	A.02.LI.b	671	bare brackish mudflat	Estuarine Embayment Intertidal Bare Brackish Mudflat
A	Estuarine	2 Embayment	LI	Littoral Intertidal	b	mudflat	A.02.LI.b	672	bare saline mudflat	Estuarine Embayment Intertidal Bare Saline Mudflat
A	Estuarine	2 Embayment	LI	Littoral Intertidal	c	saline fringing wetland (marsh)	A.02.LI.c	673	saline fringing wetland tidal pass	Estuarine Embayment Intertidal saline marsh tidal pass
A	Estuarine	2 Embayment	LI	Littoral Intertidal	c	saline fringing wetland (marsh)	A.02.LI.c	674	saline fringing wetland tidal creek	Estuarine Embayment Intertidal saline marsh tidal creek
A	Estuarine	2 Embayment	LI	Littoral Intertidal	c	saline fringing wetland (marsh)	A.02.LI.c	675	saline fringing wetland tidal creek bank	Estuarine Embayment Intertidal saline marsh tidal creek bank
A	Estuarine	2 Embayment	LI	Littoral Intertidal	c	saline fringing wetland (marsh)	A.02.LI.c	676	saline wetland hammock	Estuarine Embayment Intertidal saline marsh hammock
A	Estuarine	2 Embayment	LI	Littoral Intertidal	c	saline fringing wetland (marsh)	A.02.LI.c	677	saline wetland inland basin marsh	Estuarine Embayment Intertidal saline marsh basin
A	Estuarine	2 Embayment	LI	Littoral Intertidal	c	saline fringing wetland (marsh)	A.02.LI.c	678	saline wetland streamside marsh	Estuarine Embayment Intertidal saline streamside marsh
A	Estuarine	2 Embayment	LI	Littoral Intertidal	d	brackish fringing wetland (marsh)	A.02.LI.d	679	brackish fringing wetland tidal pass	Estuarine Embayment Intertidal brackish marsh tidal pass
A	Estuarine	2 Embayment	LI	Littoral Intertidal	d	brackish fringing wetland (marsh)	A.02.LI.d	680	brackish fringing wetland tidal creek	Estuarine Embayment Intertidal brackish marsh tidal creek
A	Estuarine	2 Embayment	LI	Littoral Intertidal	d	brackish fringing wetland (marsh)	A.02.LI.d	681	brackish fringing wetland tidal creek bank	Estuarine Embayment Intertidal brackish marsh tidal creek bank
A	Estuarine	2 Embayment	LI	Littoral Intertidal	d	brackish fringing wetland (marsh)	A.02.LI.d	682	brackish wetland hammock	Estuarine Embayment Intertidal brackish marsh hammock
A	Estuarine	2 Embayment	LI	Littoral Intertidal	d	brackish fringing wetland (marsh)	A.02.LI.d	683	brackish wetland inland basin marsh	Estuarine Embayment Intertidal brackish marsh basin

A	Estuarine	2 Embayment	LI	Littoral Intertidal	d	brackish fringing wetland (marsh	A.02.LI.d	684	brackish wetland streamside marsh	Estuarine Embayment Intertidal brackish streamside marsh
A	Estuarine	2 Embayment	LI	Littoral Intertidal	e	hardbottom	A.02.LI.e	685	bare gravel hardbottom	Estuarine Embayment Intertidal bare gravel
A	Estuarine	2 Embayment	LI	Littoral Intertidal	e	hardbottom	A.02.LI.e	686	bare limestone pavement hardbottom	Estuarine Embayment Intertidal bare limestone pavement
A	Estuarine	2 Embayment	LI	Littoral Intertidal	e	hardbottom	A.02.LI.e	687	bare bedrock hardbottom	Estuarine Embayment Intertidal bare bedrock
A	Estuarine	2 Embayment	LI	Littoral Intertidal	f	cliff	A.02.LI.f	688	cliff notch	Estuarine Embayment Intertidal cliff notch
A	Estuarine	2 Embayment	LI	Littoral Intertidal	f	cliff	A.02.LI.f	689	cliff bioerosion notch	Estuarine Embayment Intertidal Bioeroded cliff notch
A	Estuarine	2 Embayment	LI	Littoral Intertidal	f	cliff	A.02.LI.f	690	cliff cave	Estuarine Embayment Intertidal cliff cave
A	Estuarine	2 Embayment	LI	Littoral Intertidal	f	cliff	A.02.LI.f	691	cliff fracture	Estuarine Embayment Intertidal cliff fracture
A	Estuarine	2 Embayment	LI	Littoral Intertidal	f	cliff	A.02.LI.f	692	cliff rubble zone	Estuarine Embayment Intertidal cliff rubble zone
A	Estuarine	2 Embayment	LI	Littoral Intertidal	g	mangrove swamp	A.02.LI.g	693	prop root zone	Estuarine Embayment Intertidal mangrove prop root zone
A	Estuarine	2 Embayment	LI	Littoral Intertidal	g	mangrove swamp	A.02.LI.g	694	basin swamp	Estuarine Embayment Intertidal mangrove basin swamp
A	Estuarine	2 Embayment	LI	Littoral Intertidal	g	mangrove swamp	A.02.LI.g	695	buttonwood ridge	Estuarine Embayment Intertidal mangrove buttonwood ridge
A	Estuarine	2 Embayment	LI	Littoral Intertidal	g	mangrove swamp	A.02.LI.g	696	pneumatophore zone	Estuarine Embayment Intertidal mangrove pneumatophore zone
A	Estuarine	2 Embayment	LI	Littoral Intertidal	g	mangrove swamp	A.02.LI.g	697	swamp creek	Estuarine Embayment Intertidal mangrove creek
A	Estuarine	2 Embayment	LI	Littoral Intertidal	g	mangrove swamp	A.02.LI.g	698	basin pond	Estuarine Embayment Intertidal mangrove basin pond
A	Estuarine	2 Embayment	LI	Littoral Intertidal	h	beach	A.02.LI.h	699	sand beach	Estuarine Embayment Intertidal sand beach
A	Estuarine	2 Embayment	LI	Littoral Intertidal	h	beach	A.02.LI.h	700	mixed-fine sediment beach	Estuarine Embayment Intertidal mixed-fine sediment beach
A	Estuarine	2 Embayment	LI	Littoral Intertidal	h	beach	A.02.LI.h	701	mixed-fine sand and mud beach	Estuarine Embayment Intertidal mixed-fine sand and mud beach
A	Estuarine	2 Embayment	LI	Littoral Intertidal	h	beach	A.02.LI.h	702	mud beach	Estuarine Embayment Intertidal mud beach
A	Estuarine	2 Embayment	LI	Littoral Intertidal	h	beach	A.02.LI.h	703	rock and boulder beach	Estuarine Embayment Intertidal rock and boulder beach
A	Estuarine	2 Embayment	LI	Littoral Intertidal	h	beach	A.02.LI.h	704	cobble beach	Estuarine Embayment Intertidal cobble beach
A	Estuarine	2 Embayment	LI	Littoral Intertidal	h	beach	A.02.LI.h	705	mixed-coarse sediment beach	Estuarine Embayment Intertidal mixed-coarse sediment beach
A	Estuarine	2 Embayment	LI	Littoral Intertidal	h	beach	A.02.LI.h	706	gravel beach	Estuarine Embayment Intertidal gravel beach
A	Estuarine	2 Embayment	LI	Littoral Intertidal	i	rocky shore	A.02.LI.i	707	bedrock shore	Estuarine Embayment Intertidal bedrock shore
A	Estuarine	2 Embayment	LI	Littoral Intertidal	i	rocky shore	A.02.LI.i	708	hardpan shore	Estuarine Embayment Intertidal hardpan shore
A	Estuarine	2 Embayment	LI	Littoral Intertidal	i	rocky shore	A.02.LI.i	709	boulder shore	Estuarine Embayment Intertidal boulder shore
A	Estuarine	2 Embayment	LI	Littoral Intertidal	i	rocky shore	A.02.LI.i	710	cobble shore	Estuarine Embayment Intertidal cobble shore
A	Estuarine	2 Embayment	LI	Littoral Intertidal	i	rocky shore	A.02.LI.i	711	sand-bedrock shore	Estuarine Embayment Intertidal sand-bedrock shore
A	Estuarine	2 Embayment	LI	Littoral Intertidal	j	dune system	A.02.LI.j	712	foredune	Estuarine Embayment Intertidal dune system foredune
A	Estuarine	2 Embayment	LI	Littoral Intertidal	j	dune system	A.02.LI.j	713	dune crest	Estuarine Embayment Intertidal dune system dune crest
A	Estuarine	2 Embayment	LI	Littoral Intertidal	j	dune system	A.02.LI.j	714	backdune	Estuarine Embayment Intertidal dune system backdune
A	Estuarine	2 Embayment	B	Bottom	a	oyster reef	A.02.B.a	715	live oyster reef	Estuarine Embayment Subtidal oyster reef
A	Estuarine	2 Embayment	B	Bottom	a	oyster reef	A.02.B.a	716	relict oyster reef	Estuarine Embayment Subtidal relict oyster reef
A	Estuarine	2 Embayment	B	Bottom	a	oyster reef	A.02.B.a	718	periodically emergent oyster reef	Estuarine Embayment Intertidal oyster reef
A	Estuarine	2 Embayment	B	Bottom	a	oyster reef	A.02.B.a	719	oyster shell midden	Estuarine Embayment Subtidal oyster shell midden
A	Estuarine	2 Embayment	B	Bottom	b	worm reef	A.02.B.b	720	live worm reef	Estuarine Embayment Subtidal worm reef
A	Estuarine	2 Embayment	B	Bottom	b	worm reef	A.02.B.b	721	relict worm reef	Estuarine Embayment Subtidal relict worm reef
A	Estuarine	2 Embayment	B	Bottom	b	worm reef	A.02.B.b	722	live mussel reef	Estuarine Embayment Subtidal mussel reef
A	Estuarine	2 Embayment	B	Bottom	b	worm reef	A.02.B.b	723	relict mussel reef	Estuarine Embayment Subtidal relict mussel reef

A	Estuarine	2 Embayment	B	Bottom	c	unconsolidated sediments	A.02.B.c	724	bare sandy softbottom	Estuarine Embayment Subtidal bare sandy sediments
A	Estuarine	2 Embayment	B	Bottom	c	unconsolidated sediments	A.02.B.c	725	bare mineral mud softbottom	Estuarine Embayment Subtidal bare mineral mud sediments
A	Estuarine	2 Embayment	B	Bottom	c	unconsolidated sediments	A.02.B.c	726	bare organic mud softbottom	Estuarine Embayment Subtidal bare organic mud sediments
A	Estuarine	2 Embayment	B	Bottom	c	unconsolidated sediments	A.02.B.c	727	bare carbonate mud-shell hash softbottom	Estuarine Embayment Subtidal bare carbonate mud-shell hash sediments
A	Estuarine	2 Embayment	B	Bottom	c	unconsolidated sediments	A.02.B.c	728	bare carbonate mud softbottom	Estuarine Embayment Subtidal bare carbonate mud sediments
A	Estuarine	2 Embayment	B	Bottom	c	unconsolidated sediments	A.02.B.c	729	bare carbonate sand softbottom	Estuarine Embayment Subtidal bare carbonate sand sediments
A	Estuarine	2 Embayment	B	Bottom	c	unconsolidated sediments	A.02.B.c	730	bare mixed-coarse sediment softbottom	Estuarine Embayment Subtidal bare mixed-coarse sediment sediments
A	Estuarine	2 Embayment	B	Bottom	c	unconsolidated sediments	A.02.B.c	731	colonized mixed-fine softbottom	Estuarine Embayment Subtidal colonized mixed-fine sediments
A	Estuarine	2 Embayment	B	Bottom	c	unconsolidated sediments	A.02.B.c	732	vegetated mud and mixed-fine softbottom	Estuarine Embayment Subtidal vegetated mud and mixed-fine sediments
A	Estuarine	2 Embayment	B	Bottom	c	unconsolidated sediments	A.02.B.c	733	bare gravel softbottom	Estuarine Embayment Subtidal bare gravel
A	Estuarine	2 Embayment	B	Bottom	c	unconsolidated sediments	A.02.B.c	734	vegetated softbottom	Estuarine Embayment Subtidal vegetated softbottom
A	Estuarine	2 Embayment	B	Bottom	c	unconsolidated sediments	A.02.B.c	735	colonized softbottom	Estuarine Embayment Subtidal colonized softbottom
A	Estuarine	2 Embayment	B	Bottom	c	unconsolidated sediments	A.02.B.c	736	vegetated colonized softbottom	Estuarine Embayment Subtidal vegetated colonized softbottom
A	Estuarine	2 Embayment	B	Bottom	d	hardbottom	A.02.B.d	737	bare limestone pavement hardbottom	Estuarine Embayment Subtidal bare limestone pavement
A	Estuarine	2 Embayment	B	Bottom	d	hardbottom	A.02.B.d	738	vegetated rock and boulder hardbottom	Estuarine Embayment Subtidal vegetated rock and boulder hardbottom
A	Estuarine	2 Embayment	B	Bottom	d	hardbottom	A.02.B.d	739	cobble hardbottom	Estuarine Embayment Subtidal cobble hardbottom
A	Estuarine	2 Embayment	B	Bottom	d	hardbottom	A.02.B.d	740	gravel hardbottom	Estuarine Embayment Subtidal gravel hardbottom
A	Estuarine	2 Embayment	B	Bottom	d	hardbottom	A.02.B.d	741	boulder hardbottom	Estuarine Embayment Subtidal boulder hardbottom
A	Estuarine	2 Embayment	B	Bottom	d	hardbottom	A.02.B.d	742	bedrock hardbottom	Estuarine Embayment Subtidal bedrock hardbottom
A	Estuarine	2 Embayment	B	Bottom	d	hardbottom	A.02.B.d	743	colonized tidal creek bottom	Estuarine Embayment Subtidal colonized tidal creek bottom
A	Estuarine	3 Open Shor	W	Water Column	a	estuarine water column	A.03.W.a	744	sea surface	Estuarine Open Shoreline sea surface
A	Estuarine	3 Open Shor	W	Water Column	a	estuarine water column	A.03.W.a	745	surf zone	Estuarine Open Shoreline surf zone
A	Estuarine	3 Open Shor	B	Bottom	c	reef	A.03.B.c	746	worm reef	
A	Estuarine	3 Open Shor	B	Bottom	c	reef	A.03.B.c	747	mussel reef	
A	Estuarine	4 Surface ch	W	Water Column	a	river current	A.04.W.a	748	turbid upper water column	
A	Estuarine	4 Surface ch	W	Water Column	a	river current	A.04.W.a	749	clear upper water column	
A	Estuarine	4 Surface ch	L	Littoral	a	riverbank	A.04.L.a	750	vegetated	
A	Estuarine	4 Surface ch	L	Littoral	a	riverbank	A.04.L.a	751	mud	
A	Estuarine	4 Surface ch	L	Littoral	a	riverbank	A.04.L.a	752	rocky	
A	Estuarine	4 Surface ch	B	Bottom	a	riverbed	A.04.B.a	753	softbottom	
A	Estuarine	4 Surface ch	B	Bottom	a	riverbed	A.04.B.a	754	hardbottom	
A	Estuarine	4 Surface ch	B	Bottom	a	riverbed	A.04.B.a	755	scour hole	
A	Estuarine	4 Surface ch	B	Bottom	a	riverbed	A.04.B.a	756	organic deposition zone	
A	Estuarine	5 Island	L	Littoral	c	reef	A.05.L.c	757	worm reef bed	
A	Estuarine	5 Island	L	Littoral	c	reef	A.05.L.c	758	mussel bed	
A	Estuarine	5 Island	L	Littoral	c	reef	A.05.L.c	759	coral head	
A	Estuarine	9 Wetland	W	Water Column	a	deeply flooded wetland water c	A.09.W.a	760	deeply flooded wetland water column	
A	Estuarine	11 Reef	B	Bottom	c	coral reef	A.11.B.c	765	relict coral	

A	Estuarine	19 River plum	W	Water Column	a	estuarine water column	A.19.W.a	766	phytoplankton bloom	
B	Freshwat	1 Estuarine p	WU	Water Column (upper layer	a	turbid estuarine water column	B.01.WU.a	761	phytoplankton bloom	
B	Freshwat	1 Estuarine p	WU	Water Column (upper layer	a	turbid estuarine water column	B.01.WU.a	762	floating mat	Freshwater plume upper water column floating mat
B	Freshwat	2 Fresh wat	WU	Water Column (upper layer	u	undefined	B.02.WU.u	763	phytoplankton bloom	Freshwater-influenced nearshore marine phytoplankton bloom
B	Freshwat	2 Fresh wat	WU	Water Column (upper layer	u	undefined	B.02.WU.u	764	floating mat	
C	Nearshore	1 Lagoon	WU	Water Column (upper layer	a	marine water column	C.01.WU.a	767	upper water column	Marine Lagoon upper surface foam
C	Nearshore	1 Lagoon	WU	Water Column (upper layer	a	marine water column	C.01.WU.a	768	pycnocline	Nearshore marine Lagoon pycnocline
C	Nearshore	1 Lagoon	WU	Water Column (upper layer	a	marine water column	C.01.WU.a	769	hyperhaline water column	Nearshore marine Lagoon hyperhaline water column
C	Nearshore	1 Lagoon	WU	Water Column (upper layer	a	marine water column	C.01.WU.a	770	phytoplankton bloom	Nearshore marine Lagoon phytoplankton bloom
C	Nearshore	1 Lagoon	WU	Water Column (upper layer	a	marine water column	C.01.WU.a	771	epipelagic zone	Nearshore marine Lagoon epipelagic zone
C	Nearshore	1 Lagoon	WU	Water Column (upper layer	a	marine water column	C.01.WU.a	772	floating vegetation mat	Nearshore marine Lagoon floating vegetation mat
C	Nearshore	1 Lagoon	WU	Water Column (upper layer	a	marine water column	C.01.WU.a	773	counter current	Nearshore marine Lagoon counter current
C	Nearshore	1 Lagoon	WU	Water Column (upper layer	a	marine water column	C.01.WU.a	774	surf zone	Nearshore marine Lagoon surf zone
C	Nearshore	1 Lagoon	WU	Water Column (upper layer	a	marine water column	C.01.WU.a	775	sea surface	Nearshore marine Lagoon sea surface
C	Nearshore	1 Lagoon	WB	Water Column (bottom laye	u	undefined	C.01.WB.u	776	anoxic bottom water	Nearshore marine Lagoon anoxic bottom water
C	Nearshore	1 Lagoon	WB	Water Column (bottom laye	u	undefined	C.01.WB.u	777	oxic bottom water	Nearshore marine Lagoon oxic bottom water
C	Nearshore	1 Lagoon	WB	Water Column (bottom laye	u	undefined	C.01.WB.u	778	salt wedge	Nearshore marine Lagoon salt wedge
C	Nearshore	1 Lagoon	WB	Water Column (bottom laye	u	undefined	C.01.WB.u	779	groundwater seep	Nearshore marine Lagoon groundwater seep
C	Nearshore	1 Lagoon	WB	Water Column (bottom laye	u	undefined	C.01.WB.u	780	benthic boundary layer	Nearshore marine Lagoon benthic boundary layer
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	a	unconsolidated sediments	C.01.LS.a	781	bare organic mud	Nearshore marine Lagoon bare organic mud
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	a	unconsolidated sediments	C.01.LS.a	782	bare carbonate sediment	Nearshore marine Lagoon bare carbonate sediment
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	a	unconsolidated sediments	C.01.LS.a	783	bare carbonate mud	Nearshore marine Lagoon bare carbonate mud
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	a	unconsolidated sediments	C.01.LS.a	784	bare carbonate sand	Nearshore marine Lagoon bare carbonate sand
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	a	unconsolidated sediments	C.01.LS.a	785	bare mixed-coarse sediment softbottom	Nearshore marine Lagoon bare mixed-coarse sediment softbottom
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	a	unconsolidated sediments	C.01.LS.a	786	vegetated mineral sediments	Nearshore marine Lagoon vegetated mineral sediments
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	a	unconsolidated sediments	C.01.LS.a	787	vegetated organic sediments	Nearshore marine Lagoon vegetated organic sediments
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	b	rocky shore	C.01.LS.b	788	bedrock shore	Nearshore marine Lagoon bedrock shore
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	b	rocky shore	C.01.LS.b	789	hardpan shore	Nearshore marine Lagoon hardpan shore
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	b	rocky shore	C.01.LS.b	790	boulder shore	Nearshore marine Lagoon boulder shore
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	b	rocky shore	C.01.LS.b	791	cobble shore	Nearshore marine Lagoon cobble shore
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	b	rocky shore	C.01.LS.b	792	sand-bedrock shore	Nearshore marine Lagoon sand-bedrock shore
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	c	dune system	C.01.LS.c	793	foredune	Nearshore marine Lagoon dune system foredune
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	c	dune system	C.01.LS.c	794	dune crest	Nearshore marine Lagoon dune system dune crest
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	c	dune system	C.01.LS.c	795	backdune	Nearshore marine Lagoon dune system backdune
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	d	cliff	C.01.LS.d	796	cliff notch	Nearshore marine Lagoon cliff cliff notch
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	d	cliff	C.01.LS.d	797	cliff bioerosion notch	Nearshore marine Lagoon cliff cliff bioerosion notch

C	Nearshore	1 Lagoon	LS	Littoral Supratidal	d	cliff	C.01.LS.d	798	cliff cave	Nearshore marine Lagoon cliff cliff cave
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	d	cliff	C.01.LS.d	799	cliff fracture	Nearshore marine Lagoon cliff cliff fracture
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	d	cliff	C.01.LS.d	800	cliff rubble zone	Nearshore marine Lagoon cliff cliff rubble zone
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	e	mangrove swamp	C.01.LS.e	801	prop root zone	Nearshore marine Lagoon mangrove swamp prop root zone
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	e	mangrove swamp	C.01.LS.e	802	basin swamp	Nearshore marine Lagoon mangrove swamp basin swamp
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	e	mangrove swamp	C.01.LS.e	803	buttonwood ridge	Nearshore marine Lagoon mangrove swamp buttonwood ridge
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	e	mangrove swamp	C.01.LS.e	804	pneumatophore zone	Nearshore marine Lagoon mangrove swamp pneumatophore zone
C	Nearshore	1 Lagoon	LS	Littoral Supratidal	e	mangrove swamp	C.01.LS.e	805	swamp creek	Nearshore marine Lagoon mangrove swamp swamp creek
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	a	unconsolidated sediments	C.01.LI.a	806	bare organic mud softbottom	Nearshore marine Lagoon Intertidal bare organic mud softbottom
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	a	unconsolidated sediments	C.01.LI.a	807	bare carbonate sediment softbottom	Nearshore marine Lagoon Intertidal bare carbonate sediment softbottom
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	a	unconsolidated sediments	C.01.LI.a	808	bare carbonate mud softbottom	Nearshore marine Lagoon Intertidal bare carbonate mud softbottom
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	a	unconsolidated sediments	C.01.LI.a	809	bare carbonate sand softbottom	Nearshore marine Lagoon Intertidal bare carbonate sand softbottom
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	a	unconsolidated sediments	C.01.LI.a	810	bare mixed-coarse sediment softbottom	Nearshore marine Lagoon Intertidal bare mixed-coarse sediment softbottom
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	a	unconsolidated sediments	C.01.LI.a	811	vegetated softbottom- holdfast (macroalgae be	Nearshore marine Lagoon Intertidal vegetated softbottom- non-rooted
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	a	unconsolidated sediments	C.01.LI.a	812	vegetated softbottom- rooted (seagrass bed)	Nearshore marine Lagoon Intertidal vegetated softbottom- rooted
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	a	unconsolidated sediments	C.01.LI.a	813	vegetated organic softbottom	Nearshore marine Lagoon Intertidal vegetated organic softbottom
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	b	mudflat	C.01.LI.b	814	fringing vegetated brackish mudflat	Nearshore marine Lagoon Intertidal fringing vegetated brackish mudflat
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	b	mudflat	C.01.LI.b	815	fringing vegetated saline mudflat	Nearshore marine Lagoon Intertidal fringing vegetated saline mudflat
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	b	mudflat	C.01.LI.b	816	bare brackish mudflat	Nearshore marine Lagoon Intertidal bare brackish mudflat
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	b	mudflat	C.01.LI.b	817	bare saline mudflat	Nearshore marine Lagoon Intertidal bare saline mudflat
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	c	saline fringing wetland (marsh)	C.01.LI.c	818	saline fringing wetland tidal pass	Nearshore marine Lagoon Intertidal saline fringing wetland tidal pass
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	c	saline fringing wetland (marsh)	C.01.LI.c	819	saline fringing wetland tidal creek	Nearshore marine Lagoon Intertidal saline fringing wetland tidal creek
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	c	saline fringing wetland (marsh)	C.01.LI.c	820	saline fringing wetland tidal creek bank	Nearshore marine Lagoon Intertidal saline fringing wetland tidal creek bank
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	c	saline fringing wetland (marsh)	C.01.LI.c	821	saline wetland hammock	Nearshore marine Lagoon Intertidal saline wetland hammock
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	c	saline fringing wetland (marsh)	C.01.LI.c	822	saline wetland inland basin marsh	Nearshore marine Lagoon Intertidal saline wetland inland basin marsh
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	c	saline fringing wetland (marsh)	C.01.LI.c	823	saline wetland streamside marsh	Nearshore marine Lagoon Intertidal saline wetland streamside marsh
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	d	brackish fringing wetland (marsh)	C.01.LI.d	824	brackish fringing wetland tidal pass	Nearshore marine Lagoon Intertidal brackish fringing wetland tidal pass
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	d	brackish fringing wetland (marsh)	C.01.LI.d	825	brackish fringing wetland tidal creek	Nearshore marine Lagoon Intertidal brackish fringing wetland tidal creek
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	d	brackish fringing wetland (marsh)	C.01.LI.d	826	brackish fringing wetland tidal creek bank	Nearshore marine Lagoon Intertidal brackish fringing wetland tidal creek bank
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	d	brackish fringing wetland (marsh)	C.01.LI.d	827	brackish wetland hammock	Nearshore marine Lagoon Intertidal brackish wetland hammock
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	d	brackish fringing wetland (marsh)	C.01.LI.d	828	brackish wetland inland basin marsh	Nearshore marine Lagoon Intertidal brackish wetland inland basin marsh
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	d	brackish fringing wetland (marsh)	C.01.LI.d	829	brackish wetland streamside marsh	Nearshore marine Lagoon Intertidal brackish wetland streamside marsh
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	e	hardbottom	C.01.LI.e	830	bare gravel hardbottom	Nearshore marine Lagoon Intertidal hardbottom bare gravel hardbottom
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	e	hardbottom	C.01.LI.e	831	bare limestone pavement hardbottom	Nearshore marine Lagoon Intertidal hardbottom bare limestone pavement hardbottom
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	e	hardbottom	C.01.LI.e	832	bare bedrock hardbottom	Nearshore marine Lagoon Intertidal hardbottom bare bedrock hardbottom
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	f	cliff	C.01.LI.f	833	notch	Nearshore marine Lagoon Intertidal cliff notch
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	f	cliff	C.01.LI.f	834	bioerosion notch	Nearshore marine Lagoon Intertidal cliff bioerosion notch
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	f	cliff	C.01.LI.f	835	cave	Nearshore marine Lagoon Intertidal cliff cave

C	Nearshore	1 Lagoon	LI	Littoral Intertidal	f	cliff	C.01.LI.f	836	fracture	Nearshore marine Lagoon Intertidal cliff fracture
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	f	cliff	C.01.LI.f	837	rubble zone	Nearshore marine Lagoon Intertidal cliff rubble zone
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	g	mangrove swamp	C.01.LI.g	838	prop root zone	Nearshore marine Lagoon Intertidal mangrove swamp prop root zone
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	g	mangrove swamp	C.01.LI.g	839	basin swamp	Nearshore marine Lagoon Intertidal mangrove swamp basin swamp
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	g	mangrove swamp	C.01.LI.g	840	buttonwood ridge	Nearshore marine Lagoon Intertidal mangrove swamp buttonwood ridge
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	g	mangrove swamp	C.01.LI.g	841	pneumatophore zone	Nearshore marine Lagoon Intertidal mangrove swamp pneumatophore zone
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	g	mangrove swamp	C.01.LI.g	842	swamp creek	Nearshore marine Lagoon Intertidal mangrove swamp swamp creek
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	g	mangrove swamp	C.01.LI.g	843	basin pond	Nearshore marine Lagoon Intertidal mangrove swamp basin pond
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	h	beach	C.01.LI.h	844	sand beach	Nearshore marine Lagoon Intertidal sand beach
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	h	beach	C.01.LI.h	845	mixed-fine sediment beach	Nearshore marine Lagoon Intertidal mixed-fine sediment beach
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	h	beach	C.01.LI.h	846	mixed-fine sand and mud beach	Nearshore marine Lagoon Intertidal mixed-fine sand and mud beach
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	h	beach	C.01.LI.h	847	mud beach	Nearshore marine Lagoon Intertidal mud beach
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	h	beach	C.01.LI.h	848	rock and boulder beach	Nearshore marine Lagoon Intertidal rock and boulder beach
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	h	beach	C.01.LI.h	849	cobble beach	Nearshore marine Lagoon Intertidal cobble beach
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	h	beach	C.01.LI.h	850	mixed-coarse sediment beach	Nearshore marine Lagoon Intertidal mixed-coarse sediment beach
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	h	beach	C.01.LI.h	851	gravel beach	Nearshore marine Lagoon Intertidal gravel beach
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	i	rocky shore	C.01.LI.i	852	bedrock shore	Nearshore marine Lagoon Intertidal bedrock shore
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	i	rocky shore	C.01.LI.i	853	hardpan shore	Nearshore marine Lagoon Intertidal hardpan shore
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	i	rocky shore	C.01.LI.i	854	boulder shore	Nearshore marine Lagoon Intertidal boulder shore
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	i	rocky shore	C.01.LI.i	855	cobble shore	Nearshore marine Lagoon Intertidal cobble shore
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	i	rocky shore	C.01.LI.i	856	sand-bedrock shore	Nearshore marine Lagoon Intertidal sand-bedrock shore
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	j	oyster reef	C.01.LI.j	857	live oyster reef	Nearshore marine Lagoon Intertidal live oyster reef
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	j	oyster reef	C.01.LI.j	858	relict oyster reef	Nearshore marine Lagoon Intertidal relict oyster reef
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	j	oyster reef	C.01.LI.j	859	submerged oyster reef	Nearshore marine Lagoon Intertidal submerged oyster reef
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	j	oyster reef	C.01.LI.j	860	periodically emergent oyster reef	Nearshore marine Lagoon Intertidal periodically emergent oyster reef
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	j	oyster reef	C.01.LI.j	861	oyster shell midden	Nearshore marine Lagoon Intertidal oyster shell midden
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	k	mussel reef	C.01.LI.k	862	live mussel reef	Nearshore marine Lagoon Intertidal live mussel reef
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	k	mussel reef	C.01.LI.k	863	relict mussel reef	Nearshore marine Lagoon Intertidal relict mussel reef
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	l	coral reef	C.01.LI.l	864	relict coral reef	Nearshore marine Lagoon Intertidal relict coral reef
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	k	worm reef	C.01.LI.k	865	live worm reef	Nearshore marine Lagoon Intertidal live worm reef
C	Nearshore	1 Lagoon	LI	Littoral Intertidal	k	worm reef	C.01.LI.k	866	relict worm reef	Nearshore marine Lagoon Intertidal relict worm reef
C	Nearshore	2 Embayment	WU	Water Column (upper layer	a	marine water column	C.02.WU.a	867	upper water column	Nearshore marine Embayment upper water column
C	Nearshore	2 Embayment	WU	Water Column (upper layer	a	marine water column	C.02.WU.a	868	pycnocline	Nearshore marine Embayment pycnocline
C	Nearshore	2 Embayment	WU	Water Column (upper layer	a	marine water column	C.02.WU.a	869	hyperhaline water column	Nearshore marine Embayment hyperhaline water column
C	Nearshore	2 Embayment	WU	Water Column (upper layer	a	marine water column	C.02.WU.a	870	phytoplankton bloom	Nearshore marine Embayment phytoplankton bloom
C	Nearshore	2 Embayment	WU	Water Column (upper layer	a	marine water column	C.02.WU.a	871	epipelagic zone	Nearshore marine Embayment epipelagic zone
C	Nearshore	2 Embayment	WU	Water Column (upper layer	a	marine water column	C.02.WU.a	872	floating vegetation mat	Nearshore marine Embayment floating vegetation mat
C	Nearshore	2 Embayment	WU	Water Column (upper layer	a	marine water column	C.02.WU.a	873	counter current	Nearshore marine Embayment counter current

C	Nearshore	2 Embayment	WB	Water Column (bottom laye	u	undefined	C.02.WB.u	874	anoxic bottom water	Nearshore marine Embayment anoxic bottom water
C	Nearshore	2 Embayment	WB	Water Column (bottom laye	u	undefined	C.02.WB.u	875	oxic bottom water	Nearshore marine Embayment oxic bottom water
C	Nearshore	2 Embayment	WB	Water Column (bottom laye	u	undefined	C.02.WB.u	876	salt wedge	Nearshore marine Embayment salt wedge
C	Nearshore	2 Embayment	WB	Water Column (bottom laye	u	undefined	C.02.WB.u	877	groundwater seep	Nearshore marine Embayment groundwater seep
C	Nearshore	2 Embayment	WB	Water Column (bottom laye	u	undefined	C.02.WB.u	878	benthic boundary layer	Nearshore marine Embayment benthic boundary layer
C	Nearshore	11 Reef	B	Bottom	b	coral reef	C.11.B.b	879	forereef	Nearshore marine coral forereef
C	Nearshore	11 Reef	B	Bottom	b	coral reef	C.11.B.b	880	backreef	Nearshore marine coral backreef
C	Nearshore	11 Reef	B	Bottom	b	coral reef	C.11.B.b	881	spur and groove	Nearshore marine spur and groove reef
C	Nearshore	11 Reef	B	Bottom	b	coral reef	C.11.B.b	882	reef halo	Nearshore marine reef halo
C	Nearshore	11 Reef	B	Bottom	b	coral reef	C.11.B.b	883	reef remnant	Nearshore marine reef remnant
C	Nearshore	11 Reef	B	Bottom	b	coral reef	C.11.B.b	884	sand channel	Nearshore marine reef sand channel
C	Nearshore	11 Reef	B	Bottom	b	coral reef	C.11.B.b	885	patch reef	Nearshore marine patch reef
C	Nearshore	11 Reef	B	Bottom	b	coral reef	C.11.B.b	886	aggregated patch reefs	Nearshore marine aggregated patch reefs
C	Nearshore	11 Reef	B	Bottom	b	coral reef	C.11.B.b	887	pinnacle	Nearshore marine pinnacle
C	Nearshore	11 Reef	B	Bottom	b	coral reef	C.11.B.b	888	fringe reef	Nearshore marine fringe reef
C	Nearshore	11 Reef	B	Bottom	b	coral reef	C.11.B.b	889	linear reef	Nearshore marine linear reef
C	Nearshore	11 Reef	B	Bottom	b	coral reef	C.11.B.b	890	platform	Nearshore marine platform
C	Nearshore	11 Reef	B	Bottom	b	coral reef	C.11.B.b	891	individual coral head	Nearshore marine individual coral head
C	Nearshore	11 Reef	B	Bottom	b	coral reef	C.11.B.b	892	reef rubble	Nearshore marine reef rubble
D	Neritic	2 Island arc	WU	Water Column (upper layer	a	inter-island pass	D.02.WU.a	893	clear water column	Neritic Island arc Water Column (upper layer) inter-island pass clear water column
D	Neritic	2 Island arc	LI	Littoral Intertidal	a	inter-island pass	D.02.LI.a	894	turbid water column	Neritic Island arc Littoral Intertidal inter-island pass turbid water column
D	Neritic	3 Atoll	WU	Water Column (upper layer	a	interior lagoon	D.03.WU.a	895	clear water column	Neritic Atoll Water Column (upper layer) interior lagoon clear water column
D	Neritic	3 Atoll	B	Bottom	b	coral reef	D.03.B.b	896	bare carbonate sand softbottom	Neritic Atoll coral reef bare carbonate sand softbottom
D	Neritic	3 Atoll	B	Bottom	b	coral reef	D.03.B.b	897	patch reef	Neritic Atoll coral reef patch reef
D	Neritic	3 Atoll	B	Bottom	b	coral reef	D.03.B.b	898	patch reef with halo	Neritic Atoll coral reef patch reef with halo
D	Neritic	3 Atoll	B	Bottom	b	coral reef	D.03.B.b	899	forereef	Neritic Atoll coral reef forereef
D	Neritic	3 Atoll	B	Bottom	b	coral reef	D.03.B.b	900	backreef	Neritic Atoll coral reef backreef
D	Neritic	3 Atoll	B	Bottom	b	coral reef	D.03.B.b	901	spur and groove	Neritic Atoll coral reef spur and groove
D	Neritic	3 Atoll	B	Bottom	b	coral reef	D.03.B.b	902	reef halo	Neritic Atoll coral reef reef halo
D	Neritic	3 Atoll	B	Bottom	b	coral reef	D.03.B.b	903	reef remnant	Neritic Atoll coral reef remnant
D	Neritic	3 Atoll	B	Bottom	b	coral reef	D.03.B.b	904	sand channel	Neritic Atoll coral reef sand channel
D	Neritic	3 Atoll	B	Bottom	b	coral reef	D.03.B.b	905	patch reef	Neritic Atoll coral reef patch reef
D	Neritic	3 Atoll	B	Bottom	b	coral reef	D.03.B.b	906	aggregated patch reefs	Neritic Atoll coral reef aggregated patch reefs
D	Neritic	3 Atoll	B	Bottom	b	coral reef	D.03.B.b	907	pinnacle	Neritic Atoll coral reef pinnacle
D	Neritic	3 Atoll	B	Bottom	b	coral reef	D.03.B.b	908	fringe reef	Neritic Atoll coral reef fringe reef
D	Neritic	3 Atoll	B	Bottom	b	coral reef	D.03.B.b	909	linear reef	Neritic Atoll coral reef linear reef
D	Neritic	3 Atoll	B	Bottom	b	coral reef	D.03.B.b	910	platform	Neritic Atoll coral reef platform

D	Neritic	3 Atoll	B	Bottom	b	coral reef	D.03.B.b	911	individual coral head	Neritic Atoll solitary coral head
D	Neritic	3 Atoll	B	Bottom	b	coral reef	D.03.B.b	912	reef rubble	Neritic Atoll coral reef reef rubble
D	Neritic	3 Atoll	B	Bottom	c	hardbottom	D.03.B.c	913	bare limestone pavement hardbottom	Neritic Atoll bare limestone pavement
D	Neritic	3 Atoll	B	Bottom	d	unconsolidated sediments	D.03.B.d	914	carbonate sediments	Neritic Atoll carbonate sediments
D	Neritic	3 Atoll	B	Bottom	d	unconsolidated sediments	D.03.B.d	915	bare mineral mud softbottom	Neritic Atoll bare mineral mud softbottom
D	Neritic	3 Atoll	B	Bottom	d	unconsolidated sediments	D.03.B.d	916	bare organic mud softbottom	Neritic Atoll bare organic mud softbottom
D	Neritic	3 Atoll	B	Bottom	d	unconsolidated sediments	D.03.B.d	917	bare carbonate mud-shell hash softbottom	Neritic Atoll bare carbonate mud-shell hash softbottom
D	Neritic	3 Atoll	B	Bottom	d	unconsolidated sediments	D.03.B.d	918	bare carbonate mud softbottom	Neritic Atoll bare carbonate mud softbottom
D	Neritic	19 Plain	B	Bottom	a	lava field	D.19.B.a	919	compression ridge	Neritic Plain lava field compression ridge
D	Neritic	19 Plain	B	Bottom	a	lava field	D.19.B.a	920	lava tube	Neritic Plain lava field lava tube
D	Neritic	19 Plain	B	Bottom	a	lava field	D.19.B.a	921	crater	Neritic Plain lava field crater
D	Neritic	19 Plain	B	Bottom	a	lava field	D.19.B.a	922	lava flow	Neritic Plain lava field lava flow
D	Neritic	19 Plain	B	Bottom	b	sediment wave	D.19.B.b	923	organic debris	Neritic Plain sediment wave organic debris
D	Neritic	19 Plain	B	Bottom	b	sediment wave	D.19.B.b	924	mud	Neritic Plain sediment wave mud
D	Neritic	19 Plain	B	Bottom	b	sediment wave	D.19.B.b	925	sand	Neritic Plain sediment sand wave
D	Neritic	19 Plain	B	Bottom	c	bar	D.19.B.c	926	mud	Neritic Plain bar mud
D	Neritic	19 Plain	B	Bottom	c	bar	D.19.B.c	927	sand	Neritic Plain bar sand
D	Neritic	19 Plain	B	Bottom	c	bar	D.19.B.c	928	hardbottom	Neritic Plain bar hardbottom
D	Neritic	19 Plain	B	Bottom	d	moraine	D.19.B.d	929	gravel	Neritic Plain moraine gravel
D	Neritic	19 Plain	B	Bottom	d	moraine	D.19.B.d	930	pebble	Neritic Plain moraine pebble
D	Neritic	19 Plain	B	Bottom	d	moraine	D.19.B.d	931	cobble	Neritic Plain moraine cobble
D	Neritic	19 Plain	B	Bottom	d	moraine	D.19.B.d	932	boulder	Neritic Plain moraine boulder
D	Neritic	19 Plain	B	Bottom	d	moraine	D.19.B.d	933	mixed	Neritic Plain moraine mixed
D	Neritic	19 Plain	B	Bottom	d	moraine	D.19.B.d	934	bedrock	Neritic Plain moraine bedrock
D	Neritic	19 Plain	B	Bottom	e	cave	D.19.B.e	935	bedrock	Neritic Plain cave bedrock
D	Neritic	19 Plain	B	Bottom	e	cave	D.19.B.e	936	coral	Neritic Plain cave coral
D	Neritic	19 Plain	B	Bottom	e	cave	D.19.B.e	937	lava	Neritic Plain cave lava
D	Neritic	19 Plain	B	Bottom	f	crevice	D.19.B.f	938	bedrock	Neritic Plain crevice bedrock
D	Neritic	19 Plain	B	Bottom	f	crevice	D.19.B.f	939	coral	Neritic Plain crevice coral
D	Neritic	19 Plain	B	Bottom	f	crevice	D.19.B.f	940	lava	Neritic Plain crevice lava
D	Neritic	19 Plain	B	Bottom	g	sink	D.19.B.g	941	hardbottom	Neritic Plain sink hardbottom
D	Neritic	19 Plain	B	Bottom	h	debris field	D.19.B.h	942	gravel	Neritic Plain debris field gravel
D	Neritic	19 Plain	B	Bottom	h	debris field	D.19.B.h	943	pebble	Neritic Plain debris field pebble
D	Neritic	19 Plain	B	Bottom	h	debris field	D.19.B.h	944	cobble	Neritic Plain debris field cobble
D	Neritic	19 Plain	B	Bottom	h	debris field	D.19.B.h	945	boulder	Neritic Plain debris field boulder
D	Neritic	19 Plain	B	Bottom	h	debris field	D.19.B.h	946	mixed	Neritic Plain debris field mixed
D	Neritic	19 Plain	B	Bottom	h	debris field	D.19.B.h	947	bedrock	Neritic Plain debris field bedrock
D	Neritic	19 Plain	B	Bottom	i	groove	D.19.B.i	948	bedrock	Neritic Plain groove bedrock

D	Neritic	19 Plain	B	Bottom	i	groove	D.19.B.i	949	coral	Neritic Plain groove coral
D	Neritic	19 Plain	B	Bottom	j	channel	D.19.B.j	950	bedrock	Neritic Plain channel bedrock
D	Neritic	19 Plain	B	Bottom	j	channel	D.19.B.j	951	coral	Neritic Plain channel coral
D	Neritic	19 Plain	B	Bottom	k	ledge	D.19.B.k	952	bedrock	Neritic Plain ledge bedrock
D	Neritic	19 Plain	B	Bottom	k	ledge	D.19.B.k	953	coral	Neritic Plain ledge coral
D	Neritic	19 Plain	B	Bottom	l	wall	D.19.B.l	954	bedrock	Neritic Plain wall bedrock
D	Neritic	19 Plain	B	Bottom	l	wall	D.19.B.l	955	coral	Neritic Plain wall coral
D	Neritic	19 Plain	B	Bottom	m	pinnacle	D.19.B.m	956	coral	Neritic Plain pinnacle coral

E	Oceanic	3 Atoll	W	Water Column	a	interior lagoon	E.03.W.a	957	clear water column	Oceanic Atoll Water Column interior lagoon clear water column
E	Oceanic	3 Atoll	B	Bottom	b	coral reef	E.03.B.b	958	bare carbonate sand softbottom	Oceanic Atoll coral reef bare carbonate sand softbottom
E	Oceanic	3 Atoll	B	Bottom	b	coral reef	E.03.B.b	959	patch reef	Oceanic Atoll coral reef patch reef
E	Oceanic	3 Atoll	B	Bottom	b	coral reef	E.03.B.b	960	patch reef with halo	Oceanic Atoll coral reef patch reef with halo
E	Oceanic	3 Atoll	B	Bottom	b	coral reef	E.03.B.b	961	bare limestone pavement hardbottom	Oceanic Atoll coral reef bare limestone pavement hardbottom
E	Oceanic	3 Atoll	B	Bottom	b	coral reef	E.03.B.b	962	carbonate sediments	Oceanic Atoll coral reef carbonate sediments
E	Oceanic	3 Atoll	B	Bottom	b	coral reef	E.03.B.b	963	bare mineral mud softbottom	Oceanic Atoll coral reef bare mineral mud softbottom
E	Oceanic	3 Atoll	B	Bottom	b	coral reef	E.03.B.b	964	bare organic mud softbottom	Oceanic Atoll coral reef bare organic mud softbottom
E	Oceanic	3 Atoll	B	Bottom	b	coral reef	E.03.B.b	965	bare carbonate mud-shell hash softbottom	Oceanic Atoll coral reef bare carbonate mud-shell hash softbottom
E	Oceanic	3 Atoll	B	Bottom	b	coral reef	E.03.B.b	966	bare carbonate mud softbottom	Oceanic Atoll coral reef bare carbonate mud softbottom
E	Oceanic	19 Plain	B	Bottom	a	lava field	E.19.B.a	967	compression ridge	Oceanic Plain lava field compression ridge
E	Oceanic	19 Plain	B	Bottom	a	lava field	E.19.B.a	968	lava tube	Oceanic Plain lava field lava tube
E	Oceanic	19 Plain	B	Bottom	a	lava field	E.19.B.a	969	crater	Oceanic Plain lava field crater
E	Oceanic	19 Plain	B	Bottom	a	lava field	E.19.B.a	970	lava flow	Oceanic Plain lava field lava flow
E	Oceanic	19 Plain	B	Bottom	b	sediment wave	E.19.B.b	971	organic debris	Oceanic Plain sediment wave organic debris
E	Oceanic	19 Plain	B	Bottom	b	sediment wave	E.19.B.b	972	mud	Oceanic Plain sediment wave
E	Oceanic	19 Plain	B	Bottom	b	sediment wave	E.19.B.b	973	sand	Oceanic Plain sand wave sand
E	Oceanic	19 Plain	B	Bottom	c	bar	E.19.B.c	974	mud	Oceanic Plain bar mud
E	Oceanic	19 Plain	B	Bottom	c	bar	E.19.B.c	975	sand	Oceanic Plain bar sand
E	Oceanic	19 Plain	B	Bottom	c	bar	E.19.B.c	976	hardbottom	Oceanic Plain bar hardbottom
E	Oceanic	19 Plain	B	Bottom	d	moraine	E.19.B.d	977	gravel	Oceanic Plain moraine gravel
E	Oceanic	19 Plain	B	Bottom	d	moraine	E.19.B.d	978	pebble	Oceanic Plain moraine pebble
E	Oceanic	19 Plain	B	Bottom	d	moraine	E.19.B.d	979	cobble	Oceanic Plain moraine cobble
E	Oceanic	19 Plain	B	Bottom	d	moraine	E.19.B.d	980	boulder	Oceanic Plain moraine boulder
E	Oceanic	19 Plain	B	Bottom	d	moraine	E.19.B.d	981	mixed	Oceanic Plain moraine mixed
E	Oceanic	19 Plain	B	Bottom	d	moraine	E.19.B.d	982	bedrock	Oceanic Plain moraine bedrock
E	Oceanic	19 Plain	B	Bottom	e	cave	E.19.B.e	983	bedrock	Oceanic Plain cave bedrock
E	Oceanic	19 Plain	B	Bottom	e	cave	E.19.B.e	984	coral	Oceanic Plain cave coral
E	Oceanic	19 Plain	B	Bottom	e	cave	E.19.B.e	985	lava	Oceanic Plain cave lava
E	Oceanic	19 Plain	B	Bottom	e	crevice	E.19.B.e	986	bedrock	Oceanic Plain crevice bedrock

E	Oceanic	19 Plain	B	Bottom	e	crevice	E.19.B.e	987	coral	Oceanic Plain crevice coral
E	Oceanic	19 Plain	B	Bottom	e	crevice	E.19.B.e	988	lava	Oceanic Plain crevice lava
E	Oceanic	19 Plain	B	Bottom	f	sink	E.19.B.f	989	hardbottom	Oceanic Plain sink hardbottom
E	Oceanic	19 Plain	B	Bottom	g	debris field	E.19.B.g	990	gravel	Oceanic Plain debris field gravel
E	Oceanic	19 Plain	B	Bottom	g	debris field	E.19.B.g	991	pebble	Oceanic Plain debris field pebble
E	Oceanic	19 Plain	B	Bottom	g	debris field	E.19.B.g	992	cobble	Oceanic Plain debris field cobble
E	Oceanic	19 Plain	B	Bottom	g	debris field	E.19.B.g	993	boulder	Oceanic Plain debris field boulder
E	Oceanic	19 Plain	B	Bottom	g	debris field	E.19.B.g	994	mixed	Oceanic Plain debris field mixed
E	Oceanic	19 Plain	B	Bottom	g	debris field	E.19.B.g	995	bedrock	Oceanic Plain debris field bedrock
E	Oceanic	19 Plain	B	Bottom	h	groove	E.19.B.h	996	bedrock	Oceanic Plain groove bedrock
E	Oceanic	19 Plain	B	Bottom	h	groove	E.19.B.h	997	coral	Oceanic Plain groove coral
E	Oceanic	19 Plain	B	Bottom	l	channel	E.19.B.l	998	bedrock	Oceanic Plain channel bedrock
E	Oceanic	19 Plain	B	Bottom	l	channel	E.19.B.l	999	coral	Oceanic Plain channel coral
E	Oceanic	19 Plain	B	Bottom	j	ledge	E.19.B.j	1000	bedrock	Oceanic Plain ledge bedrock
E	Oceanic	19 Plain	B	Bottom	j	ledge	E.19.B.j	1001	coral	Oceanic Plain ledge coral
E	Oceanic	19 Plain	B	Bottom	k	wall	E.19.B.k	1002	bedrock	Oceanic Plain wall bedrock
E	Oceanic	19 Plain	B	Bottom	k	wall	E.19.B.k	1003	coral	Oceanic Plain coral reef wall

B. Kutcher et al. 2005, “A recommendation for a comprehensive habitat and land use classification system for the National Estuarine Research Reserve System”

Contact: Tom Kutcher, tomk@nberr.org, Narragansett Bay National Estuarine Research Reserve
 Website: <http://nerrs.noaa.gov/NarragansettBay/>

1000. Marine Habitats	2000. Estuarine Habitats	2300. Supratidal Haline	2500. Intertidal Fresh
1100. <i>Subtidal</i>	2100. <i>Subtidal Haline</i>	2310. Rock Bottom	2510. Aquatic Bed
1110. Rock Bottom	2110. Rock Bottom	2311. Bedrock	2511. Rooted Algal
1111. Bedrock	2111. Bedrock	2312. Rubble	2512. Drift Algal
1112. Rubble	2112. Rubble	2320. Unconsolidated Bottom	2513. Rooted Vascular
1120. Unconsolidated Bottom	2120. Unconsolidated Bottom	2321. Cobble	2514. Floating Vascular
1121. Cobble	2121. Cobble	2322. Gravel	2515. Aquatic Moss
1122. Gravel	2122. Gravel	2323. Sand	2520. Streambed
1123. Sand	2123. Sand	2324. Mud	2521. Bedrock
1124. Mud	2124. Mud	2325. Organic	2522. Rubble
1125. Organic	2125. Organic	2330. Aquatic Bed	2523. Cobble
1130. Aquatic Bed	2130. Aquatic Bed	2331. Rooted Algal	2524. Gravel
1131. Rooted Algal	2131. Rooted Algal	2332. Drift Algal	2525. Sand
1132. Drift Algal	2132. Drift Algal	2333. Rooted Vascular	2526. Mud
1133. Rooted Vascular	2133. Rooted Vascular	2334. Floating Vascular	2527. Organic
1134. Faunal	2134. Floating Vascular	2340. Emergent Wetland	2530. Rocky Shore
1140. Reef	2135. Faunal	2341. Persistent	2531. Bedrock
1141. Mollusk	2140. Reef	2342. Nonpersistent	2532. Rubble
1142. Coral	2141. Mollusk	2350. Scrub-Shrub Wetland	2540. Unconsolidated Shore
1143. Worm	2142. Worm	2351. BLD	2541. Cobble
1144. Artificial	2143. Artificial	2352. NLD	2542. Gravel
1200. <i>Intertidal</i>	2200. <i>Intertidal Haline</i>	2353. BLE	2543. Sand
1210. Aquatic Bed	2210. Aquatic Bed	2354. NLE	2544. Mud
1211. Rooted Algal	2211. Rooted Algal	2355. Dead	2545. Organic
1212. Drift Algal	2212. Drift Algal	2360. Forested Wetland	2550. Emergent Wetland
1213. Rooted Vascular	2213. Rooted Vascular	2361. BLD	2551. Persistent
1220. Reef	2214. Floating Vascular	2362. NLD	2552. Nonpersistent
1221. Coral	2220. Reef	2363. BLE	2560. Scrub-Shrub Wetland
1222. Worm	2221. Mollusk	2364. NLE	2561. BLD
1230. Rocky Shore	2222. Worm	2365. Mixed	2562. NLD
1231. Bedrock	2230. Streambed	2366. Dead	2563. BLE
1232. Rubble	2231. Bedrock	2400. <i>Subtidal Fresh</i>	2564. NLE
1240. Unconsolidated Shore	2232. Rubble	2410. Rock Bottom	2565. Dead
1241. Cobble	2233. Cobble	2411. Bedrock	2570. Forested Wetland
1242. Gravel	2234. Gravel	2412. Rubble	2571. BLD
1243. Sand	2235. Sand	2420. Unconsolidated Bottom	2572. NLD
1244. Mud	2236. Mud	2421. Cobble	2573. BLE
1245. Organic	2337. Organic	2422. Gravel	2574. NLE
	2240. Rocky Shore	2423. Sand	2575. Mixed
	2241. Bedrock	2424. Mud	2575. Dead
	2242. Rubble	2425. Organic	
	2250. Unconsolidated Shore	2430. Aquatic Bed	
	2251. Cobble	2431. Rooted Algal	
	2252. Gravel	2432. Drift Algal	
	2253. Sand	2433. Rooted Vascular	
	2254. Mud	2434. Floating Vascular	
	2255. Organic	2435. Aquatic Moss	
	2260. Emergent Wetland	2440. Reef	
	2261. Persistent	2441. Mollusk	
	2262. Nonpersistent		
	2270. Scrub-Shrub Wetland		
	2271. BLD		
	2272. NLD		
	2273. BLE		
	2274. NLE		
	2275. Dead		
	2280. Forested Wetland		
	2281. BLD		
	2282. NLD		
	2283. BLE		
	2284. NLE		
	2285. Mixed		
	2286. Dead		

3000. Riverine Habitats	4000. Lacustrine Habitats	5000. Palustrine Habitats	6000. Upland Habitats
3100. <i>Lower Perennial</i>	4100. <i>Limnetic</i>	5100. <i>Perennial Water</i>	6100. <i>Supratidal Upland</i>
3110. Unconsolidated Bottom	4110. Rock Bottom	5110. Rock Bottom	6110. Rocky Upland
3111. Gravel	4111. Bedrock	5111. Bedrock	6111. Bedrock
3112. Sand	4112. Rubble	5112. Rubble	6112. Rubble
3113. Mud	4120. Unconsolidated bottom	5120. Unconsolidated Bottom	6120. Unconsolidated Upland
3114. Organic	4121. Cobble	5121. Cobble	6121. Cobble
3120. Aquatic Bed	4122. Gravel	5122. Gravel	6122. Gravel
3121. Aquatic Moss	4123. Sand	5123. Sand	6123. Sand
3122. Rooted Vascular	4124. Mud	5124. Mud	6124. Clay
3123. Floating Vascular	4125. Organic	5125. Organic	6125. Loam
3130. Rocky Shore	4130. Aquatic Bed	5130. Aquatic Bed	6126. Organic
3131. Bedrock	4131. Algal	5131. Algal	6130. Herbaceous Upland
3132. Rubble	4132. Aquatic Moss	5132. Aquatic Moss	6131. Grassland
3140. Unconsolidated Shore	4133. Rooted Vascular	5133. Rooted Vascular	6132. Broad-leaved Herbs
3141. Cobble	4134. Floating Vascular	5134. Floating vascular	6140. Scrub-Shrub Upland
3142. Gravel	4200. <i>Littoral</i>	5140. Emergent Wetland	6141. BLD
3143. Sand	4210. Rock Bottom	5141. Nonpersistent	6142. NLD
3144. Mud	4211. Bedrock	5200. <i>Intermittent or Saturated</i>	6143. BLE
3145. Organic	4212. Rubble	5210. Unconsolidated Shore	6144. NLE
3150. Emergent Wetland	4220. Unconsolidated Bottom	5211. Cobble	6145. Dead
3151. Nonpersistent	4221. Cobble	5212. Gravel	6150. Forested Upland
3200. <i>Upper Perennial</i>	4222. Gravel	5213. Sand	6151. BLD
3210. Rock Bottom	4223. Sand	5214. Mud	6152. NLD
3211. Bedrock	4224. Mud	5215. Organic	6153. BLE
3212. Rubble	4225. Organic	5220. Moss-Lichen Wetland	6154. NLE
3220. Unconsolidated Bottom	4230. Aquatic Bed	5221. Moss	6155. Mixed
3221. Cobble	4231. Algal	5222. Lichen	6156. Dead
3222. Gravel	4232. Aquatic Moss	5230. Emergent Wetland	6200. <i>Inland Upland</i>
3223. Sand	4233. Rooted Vascular	5231. Nonpersistent	6210. Rocky Upland
3224. Mud	4234. Floating vascular	5232. Persistent	6211. Bedrock
3230. Aquatic Bed	4240. Rocky Shore	5240. Scrub-Shrub Wetland	6212. Rubble
3231. Algal	4241. Bedrock	5241. BLD	6220. Unconsolidated Upland
3232. Aquatic Moss	4242. Rubble	5242. NLD	6221. Cobble
3233. Rooted Vascular	4250. Unconsolidated Shore	5243. BLE	6222. Gravel
3234. Floating Vascular	4251. Cobble	5244. NLE	6223. Sand
3240. Rocky Shore	4252. Gravel	5245. Dead	6224. Clay
3241. Bedrock	4253. Sand	5250. Forested Wetland	6225. Loam
3242. Rubble	4254. Mud	5251. BLD	6226. Organic
3250. Unconsolidated Shore	4255. Organic	5252. NLD	6230. Herbaceous Upland
3251. Cobble	4260. Emergent Wetland	5253. BLE	6231. Grassland
3252. Gravel	4261. Nonpersistent	5254. NLE	6232. Broad-leaved Herbs
3253. Sand		5255. Mixed	6240. Scrub-Shrub Upland
3254. Mud		5256. Dead	6241. BLD
3255. Organic			6242. NLD
3260. Emergent Wetland			6243. BLE
3261. Nonpersistent			6244. NLE
3300. <i>Intermittent</i>			6245. Dead
3310. Streambed			6250. Forested Upland
3311. Bedrock			6251. BLD
3312. Rubble			6252. NLD
3313. Cobble			6253. BLE
3314. Gravel			6254. NLE
3315. Sand			6255. Mixed
3316. Mud			6256. Dead
3317. Organic			
3318. Vegetated			

7000. Perennial Snow and Ice Habitats	8000. Cultural Land Cover	
7100. <i>Perennial Snowfields</i>	8100. <i>Developed Upland</i>	8300. <i>Developed and Managed Wetlands and Water</i>
7200. <i>Glaciers</i>	8110. Impervious Cover	8310. Impervious Cover
	8111. Paved Lot	8311. Impervious Bottom
	8112. Paved Roadway	8312. Impervious In-water Structure
	8113. Large Building	8320. Built-up Cover
	8114. Impervious Complex	8321. Pervious In-water Structure
	8120. Built-up Cover	8322. In-water Commercial or Service Complex
	8121. Commercial or Service Complex	8323. In-water Industrial Complex
	8122. Industrial Complex	8324. Shellfish Aquaculture
	8130. Residential Cover	8325. Finfish Aquaculture
	8131. Low Density	8330. Residential Cover
	8132. Medium Density	8331. In-water Residential Complex
	8133. High Density	8340. Rocky Cover
	8140. Rocky Cover	8341. Rocky Shoreline Structure
	8141. Rocky Revetment	8342. Rocky In-water Structure
	8142. Open Quarry	8350. Unconsolidated Cover
	8150. Unconsolidated Cover	8351. Managed Unconsolidated Bottom
	8151. Cleared Land	8352. Managed Unconsolidated Shore
	8151. Dirt/gravel Lot	8360. Herbaceous Cover
	8152. Dirt/gravel Road	8361. Managed Herbaceous Wetland
	8153. Railway Corridor	8362. Agricultural Herbaceous Wetland
	8154. Mining Operation	8363. Grazed Herbaceous Wetland
	8155. Landfill Operation	8370. Shrub Cover
	8160. Herbaceous Cover	8371. Managed Wetland Shrubs
	8161. Managed Turf	8372. Agricultural Wetland Shrubs
	8162. Managed Garden	8373. Grazed Shrub Wetland
	8163. Managed Old Field	8380. Tree Cover
	8170. Shrub Cover	8381. Managed Wetland Trees
	8171. Managed Shrubs	8382. Agricultural Wetland Trees
	8180. Tree Cover	8383. Grazed Wooded Wetland
	8181. Managed Trees	
	8200. <i>Agricultural Upland</i>	
	8210. Rocky Cover	
	8211. Rocky Revetment	
	8220. Unconsolidated Cover	
	8221. Unvegetated Farmland	
	8230. Herbaceous Cover	
	8231. Turf	
	8232. Pasture	
	8233. Hay Meadow	
	8234. Crops/Cover Crops	
	8240. Shrub Cover	
	8241. Shrub Nursery	
	8242. Grazed Shrub Upland	
	8250. Tree Cover	
	8251. Tree Farm	
	8252. Orchard	
	8253. Grazed Wooded Upland	

C. Greene et. al (In Press), “Construction of digital potential marine benthic habitat maps using a coded classification scheme and their application”

Contact: Gary Greene, greeneg@mlml.calstate.edu, Moss Landing Marine Laboratories

Website: <http://geooce.mlml.calstate.edu/habcent.htm>

This code is based on the deep-water habitat characterization scheme developed by Greene et al. (1999) and modified for use in mapping habitats offshore of California (Greene et al. 2004; In Press). The code is designed so that the first character in the code, a capital letter, indicates one of nine Megahabitat types. These general Megahabitat types with suggested depth ranges in parentheses (depth estimations and can be changed to fit depth ranges known to occur for the mapping project at hand) are as follows:

A = Aprons, continental rise, deep fans and bajadas (3000-4000m)
B = Basin floors, borderland types (floors at 1000-2500m)
E = Estuary, (0-100m)
F = Flanks, continental slope, basin/island flanks (200-3000m)
I = Inland seas, fiords, and narrow inlets or passages (0-200m)
P = Plains, abyssal (4000-6000+ m)
R = Ridges and seamounts (crests at 200-2500m)
S = Shelf, continental and island shelves (0-200m)
Z = Zone of fractures (3000-5000m) or fracture zones associated with spreading ridges

The second character in the code, a lower case letter, indicates bottom induration (hardness) and consists of the following:

h = hard bottom (e.g., rock outcrop or sediment pavement)
m = mixed hard and soft bottom (e.g., local sediment cover of bedrock)
s = soft bottom, sediment cover

Sediment types (for above indurations) - Use parentheses.

(b) = boulder
(c) = cobble
(p) = pebble
(g) = gravel
(s) = sand
(m) = mud, silt, clay
(h) = halimeda sediment, carbonate

When inferred, use question mark; i.e., (m?). This part of the code is not always used so is not considered as a character in the code.

The third character in the code, another lower case letter, not always used, indicates the Meso- or Macrohabitat type (based on scale). These types consist of the following:

a = atoll
b = beach, relic (submerged)
c = canyon
d = deformed, tilted and folded bedrock
e = exposure, bedrock
f = flats, floors
g = gully, channel
i = ice-formed feature or deposit, moraine, drop-stone depression
k = karst, solution pit, sink
l = landslide
m = mound, depression; includes linear ridges
n = enclosed waters, lagoon
o = overbank deposit (levee)
p = pinnacle, cone

(Note: Pinnacles are often difficult to distinguish from boulders. Therefore, these features may be used in conjunction [as (b)/p] to designate the meso/macrophabitat).

r = rill (subterranean winnowing of sediments forming linear depressions on surface)
s = scarp, cliff, fault or slump scar
t = terrace
w = sediment waves (10 cm to <m amplitude) and dunes (10s of m in amplitude)
y = delta, fan
Z_# = zooxanthellae hosting structure, carbonate reef
Z₁ = barrier reef
Z₂ = fringing reef

Z₃ = head, bommie
 Z₄ = patch reef
 Z₅ = back reef
 Z₆ = reef flat
 Z₇ = reef crest
 Z₈ = forereef

The fourth character in the code, a subscript letter (in GIS preceded by an underline [i.e., a], is a modifier that describes the texture, bedform, biology or rock type and consists of the following:

a = anthropogenic (artificial reef/breakwall/shipwreck/disturbances)
 (a-dd) = dredge disturbances
 (a-dg) = dredge grooves or channels
 (a-dp) = dredge potholes
 (a-dm) = dredge mounds (disposal)
 (a-td) = trawl disturbances
b = bimodal (conglomeratic, mixed [includes gravel, cobbles and pebbles])
c = consolidated sediment (includes claystone, mudstone, siltstone, sandstone, breccia, or conglomerate)
d = differentially eroded
f = fracture, joint; faulted
g = granite
h = hummocky, irregular relief
i = interface, lithologic contact
k = kelp
l = limestone or carbonate
m = massive sedimentary bedrock
o = outwash
p = pavement
r = ripples (>10 cm in amplitude)
s = scour (current or ice, direction noted)
u = unconsolidated sediment
v = volcanic rock

Seafloor Slope - Use category numbers, which is the fifth character in the code. Typically calculated for survey area from x-y-z multibeam data. Numbers in parentheses are suggestions only and can be tailored to meet objectives of the habitat mapping exercise at hand.

1 Flat (0-5°)
 2 Sloping (5-30°)
 3 Steeply Sloping (30-60°)
 4 Vertical (60-90°)
 5 Overhang (> 90°)

Seafloor Complexity - Use category letters (in caps), the sixth character in the code. Typically calculated for survey area from x-y-z multibeam slope data using neighborhood statistics and reported in standard deviation units. Numbers in parentheses are suggestions only and can be tailored to meet objectives of the habitat mapping exercise at hand.

A Very Low Complexity (-1 to 0)
 B Low Complexity (0 to 1)
 C Moderate Complexity (1 to 2)
 D High Complexity (2 to 3)
 E Very High Complexity (3+)

An example of how this code for remotely sensed data can be used is given below:

Sscu4 (Q, Qsp) = Canyon head indenting shelf with smooth, soft, gentle-sloping sedimentary walls locally crop out as steep (near vertical) scarps (10-100m).

Ssfu (Q) = Flat to gently sloping shelf with soft, unconsolidated sediment (10-150m).

Fhm (Tpr) = Continental slope with sedimentary (sandstone) bedrock locally cropping out and smooth to moderately irregular relief (<1-3 m high): m means exposures often covered with sediment (200-2500m).

Geologic Unit – When possible, the associated geologic unit is identified for each habitat type and follows the habitat designation in parentheses. Examples given below:

Shp_a1D(Q/R) - Continental shelf megahabitat; flat, highly complex hard seafloor with pinnacles differentially eroded.
Geologic unit = Quaternary/Recent.

Fhd_d2C (Tmm) - Continental slope megahabitat; sloping hard seafloor of deformed (tilted, faulted, folded), differentially eroded bedrock exposure forming overhangs and caves. Geologic unit = Tertiary Miocene Monterey Formation.

Determined from video, still photos, or direct observation. Macro/Microhabitat – Preceded by an asterisk. Use parentheses for geologic attributes, brackets for biologic attributes. Based on observed small-scale seafloor features.

Geologic attributes (note percent grain sizes when possible)

- (a) = anthropogenic (e.g., cables, pipelines, disturbances)
 - (a-t) = trawl trails or grooves
 - (a-d) = dredge tracks, pits or mounds
- (b) = boulder
 - (b-d) = dropstone (kelp or ice)
- (c) = cobble
- (d) = deformed, faulted, or folded
- (e) = exposure, bedrock (sedimentary, igneous, or metamorphic)
 - (e-s) = smooth bedrock surface
 - (e-r) = rough bedrock surface
- (f) = fans or aprons
- (g) = gravel
- (h) = halimeda sediment, carbonate slabs or mound
- (i) = interface
- (j) = joints, cracks, crevices, and overhang (differentially eroded)
- (k) = knob or ridge
- (l) = limestone, carbonate deposit
- (m) = mud, silt, or clay
- (n) = notch, groove
- (p) = pebble
- (q) = coquina (shell hash)
- (r) = rubble
- (s) = sand
- (t) = flat terrace-like seafloor including sedimentary pavements
- (u) = Undulating surface, hummocky
 - (u-r) = ripples
 - (u-s) = scours
 - (u-w) = sediment wave
- (w) = wall, scarp, or cliff

Biologic attributes

- [a] = algae
 - [a-b] = red algae
 - [a-g] = green algae
 - [a-r] = red algae
- [b] = bryozoans
- [c] = corals
- [d] = detritus, drift algae
- [e] = eelgrass
- [g] = gorgonians
- [h] = holothurians
- [k] = kelp
- [n] = anemones
- [o] = other sessile organisms
 - [o-c] = crinoids
- [s] = sponges
- [t] = tracks, trails, or trace fossils (bioturbation)
 - [t-b] = burrows
 - [t-m] = mounds
- [u] = unusual organisms, or chemosynthetic communities
- [w] = worm tubes
 - [w-s] = spoon worms

Examples:

*(m)[w]1C - Flat or nearly flat mud (100%) bottom with worm tubes; moderate complexity.

*(s/c)1A - Sand bottom (>50%) with cobbles. Flat or nearly flat with very low complexity.

*(h)[c]1E - Coral reef on flat bottom with halimeda sediment. Very high complexity.

Shp_d1D(Q/R)*[m][w]1C - *Large-scale habitat type*: Continental shelf megahabitat; flat, highly complex hard seafloor with pinnacles differentially eroded. Geologic unit = Quaternary/Recent. *Small-scale habitat type*: Flat or nearly flat mud (100%) bottom with worm tubes; moderate complexity.

D. **Valentine et al. (2005), “Classification of Marine Sublittoral Habitats, with Application to the Northeastern North America Region”**

Contact: Page Valentine, pvalentine@usgs.gov, Woods Hole USGS office

Website: <http://woodshole.er.usgs.gov/project-pages/stellwagen/>

The classification includes classes, subclasses, categories, and attributes. It is designed as a template for a database. Class numbers are unique. Themes are the major subject elements of the classification and can include one to many classes. Themes are informal units and are not incorporated into the habitats database. Observations from Classes 2, 3, 11, and 24 are used to compile habitat names. Category and attribute terms in parentheses are not a complete list.

Theme 1, Topographical Setting

Class 1: Topographical setting: major seabed features and industrial structures

Subclass 1: Shallow photic (presence of macrophyte algae)

Subclass 2: Deep aphotic (absence of macrophyte algae)

Categories: Seabed slope, major physiographic and biogenic features and industrial structures

Attributes: Angle of seabed slope, types of seabed features (e.g., basin, ridge, shelf edge reef), and industrial structures (e.g., cable, oil platform)

All categories and attributes apply to Subclasses 1 and 2.

Theme 2, Seabed Dynamics and Currents

Class 2: Seabed dynamics and currents

Subclass 1: Mobile substrate

Subclass 2: Immobile substrate

Subclass 3: Intermixed mobile and immobile substrates

Categories: Types of currents (e.g., tidal, storm wave) and types of events (e.g., storms) causing sediment mobility

Attributes: Strength of currents and frequency of events (e.g., daily, monthly) causing sediment mobility

All categories and attributes apply to Subclasses 1–3.

Theme 3, Seabed Texture, Hardness, and Layering in the Upper 5–10 cm

Class 3: Seabed texture, hardness, and layering in the upper 5–10 cm

Subclass 1: Fine-grained sediment: mud, very fine (4 phi) sand, and fine (3 phi) sand

Subclass 2: Coarse-grained sediment: medium (2 phi) sand, coarse (1 phi) sand, very coarse (0 phi) sand, and gravel (gravel is composed of granules > 1 mm and < 2 mm; pebbles < 64 mm; cobbles < 256 mm; and boulders > 256 mm).

Subclass 3: Mixed fine-grained and coarse-grained sediment: mud, sand, and gravel mixtures

Subclass 4: Rock or other hard seabed (with or without mud, sand, gravel)

Categories: Descriptive sediment and hard seabed types (e.g., mud veneer on clay, gravel pavement, cobbles in muddy sand, sand veneer on rock outcrop)

Attributes: Percentage of seabed covered by sediment and hard seabed types

Categories apply to appropriate subclasses; all attributes apply to Subclasses 1–4.

Theme 4, Seabed Grain-Size Analysis

Class 4: Seabed grain sizes: general description

Subclass 1: General sediment description

Categories: Descriptive texture classification, sorting, grain size distribution, and particle shape

Attributes: Major descriptive texture classes (e.g., silty sand, gravelly mud), degree of sorting (e.g., well sorted), skewness (e.g., symmetrical), kurtosis (e.g., mesokurtic), and particle shape (e.g., rounded)

Class 22: Seabed grain sizes - major Wentworth size-classes

Subclass 1: Major Wentworth grain size-classes, weight percent

Categories: Major Wentworth grain size-classes (e.g., sand, gravel, silt, clay, mud)

Attributes: Weight percent of major Wentworth grain size-classes

Class 23: Seabed grain sizes - phi and all Wentworth size-classes

Subclass 1: Phi and all Wentworth grain size-classes, weight percent

Categories: Phi and all Wentworth size-classes (e.g., fine sand, coarse silt)

Attributes: Weight percent of phi and all Wentworth size-classes

Theme 5, Seabed Roughness

Class 5: Seabed roughness - bedforms

Subclass 1: Bedforms (physical structures)

- Categories: Bedform types (e.g., ripples, sand dunes)
 - Attributes: Percentage of seabed covered by bedform types
- Class 6: Seabed roughness - shell materials
 - Subclass 1: Shell materials (physical structures)
 - Categories: Types of shell materials and deposits (e.g., shell fragments, shell deposits)
 - Attributes: Percentage of seabed covered by shell material and deposit types
- Class 7: Seabed roughness - rough sediments and hard seabeds
 - Subclass 1: Rough sediments and hard seabeds (physical structures)
 - Categories: Associations of sediment particles, sediment type, seabed structures, and rock outcrops (e.g., cobbles in patches, piled boulders, pebbles in sand dune troughs, irregular rock outcrop)
 - Attributes: Percentage of seabed covered by rough sediment and hard seabed types
- Class 8: Seabed roughness - biogenic structures
 - Subclass 1: Biogenic structures (physical structures)
 - Categories: Types of biogenic modifications of the seabed (e.g., crab depressions, fish burrows)
 - Attributes: Percentage of seabed covered by types of biogenic structures
- Class 12: Seabed roughness - anthropogenic marks
 - Subclass 1: Anthropogenic marks (physical structures)
 - Categories: Types of marks made on the seabed by human activities (e.g., trawl marks, anchor marks)
 - Attributes: Percentage of seabed covered by types of anthropogenic marks
- Class 13: Seabed roughness - anthropogenic structures
 - Subclass 1: Anthropogenic structures (physical structures)
 - Categories: Types of minor man-made structures and equipment on the seabed (e.g., types of fishing gear)
 - Attributes: Percentage of seabed covered by types of anthropogenic structures
- Class 24: Seabed roughness - physical structures combined
 - Subclass 1: Extent of physical structures
 - Categories: Types of physical structures
 - Attributes: Percentage of seabed covered by physical structures by type and all combined
- Class 24: summarizes observations for Classes 5–8, 12, and 13.
- Class 9: Seabed roughness: attached epifauna
 - Subclass 1: Attached epifauna (biological structures)
 - Categories: Epifaunal groups attached to the seabed surface (e.g., erect sponges, tunicates, brachiopods)
 - Attributes: Percentage of seabed covered by types of attached epifauna
- Class 10: Seabed roughness - emergent epifauna
 - Subclass 1: Emergent epifauna (biological structures)
 - Categories: Epifaunal groups emergent from below the seabed surface (e.g., burrowing anemones, sea pens)
 - Attributes: Percentage of seabed covered by types of emergent epifauna
- Class 11: Seabed roughness - biological structures combined
 - Subclass 1: Extent of biological structures
 - Categories: Types of biological structures
 - Attributes: Percentage of seabed covered by biological structures by type and all combined
- Class 11: Summarizes observations for Classes 9 and 10.

Theme 6, Fauna and Flora

- Class 14: Faunal groups
 - Subclasses 1–6: Faunal groups (in several subclasses based on different methods of data collection; e.g., visual observations and/or specimens from various sampler types)
 - Categories: Faunal groups (e.g., erect sponges, burrowing anemones, sea stars, attached anemones)
 - Attributes: Presence/absence or percentage of seabed covered by individual faunal groups
- All categories and attributes apply to Subclasses 1–6.
- Class 15: Faunal species
 - Subclasses 1–6: Faunal species (in several subclasses based on different methods of data collection; e.g., visual observations and/or specimens from various sampler types)
 - Categories: Faunal species (e.g., Atlantic cod)
 - Attributes: Presence/absence or percentage of seabed covered by individual faunal species
- All categories and attributes apply to Subclasses 1–6.
- Class 16: Floral groups
 - Subclasses 1–6: Floral groups (in several subclasses based on different methods of data collection; e.g., visual observations and/or specimens from various sampler types)
 - Categories: Floral groups (e.g., calcareous algae, kelp)
 - Attributes: Presence/absence or percentage of seabed covered by individual floral groups
- All categories and attributes apply to Subclasses 1–6.
- Class 17: Floral species
 - Subclasses 1–6: Floral species (in several subclasses based on different methods of data collection; e.g., visual observations and/or specimens from various sampler types)

Categories: Floral species

Attributes: Presence/absence or percentage of seabed covered by individual floral species

All categories and attributes apply to Subclasses 1–6.

Theme 7, Habitat Association and Usage

Class 18: Fauna-habitat association: essential fish habitat (EFH)

Subclasses 1–6: Fauna-habitat association (in several subclasses based on different methods of data collection; e.g., visual observations and/or specimens from various sampler types)

Categories: Faunal species (e.g., Atlantic cod, haddock, yellowtail flounder)

Attributes: Types of fauna-habitat association by species (e.g. adult, spawning, juvenile habitat)

All categories and attributes apply to Subclasses 1–6.

Class 19: Human usage of habitat

Subclass 1: Human usage of habitat

Categories: Disturbed, undisturbed, or recovering habitat; kinds of disturbance (e.g., fishing, extraction)

Attributes: Types of disturbance activities (e.g., otter trawling, minerals mining)

Theme 8, Habitat Recovery from Disturbance

Class 20: Habitat recovery from fishing disturbance

Subclass 1: Fishing disturbance

Categories: Recovery of physical structures and biological structures (e.g., bedforms, attached epifauna)

Attributes: Time required for recovery (e.g., months, year, decades)

Class 21: Habitat recovery from natural disturbance

Subclass 1: Natural disturbance

Categories: Recovery of physical structures and biological structures (e.g., fish burrows, emergent epifauna)

Attributes: Time required for recovery (e.g., months, year, decades)

Note: Explanation of (1) habitat descriptions, (2) structural complexity of habitats, (3) three levels of descriptive habitat names, (4) probable marine sublittoral habitat types are found in Appendices 2-5 at <http://woodshole.er.usgs.gov/project-pages/stellwagen>.

APPENDIX 2: Contact information of habitat classification experts.

Contact	Organization	Email	Reasons for Collaboration
Allee, Becky	NOAA - CSC	becky.allee@noaa.gov	Technical advice and updates on CMECS
Annett, Brendan	Waquoit Bay NERR	brendan.annett@state.ma.us	Waquoit Bay implementation of the NERR classification scheme
Dethier, Megan	University of WA	mdethier@u.washington.edu	Experience with classification studies
Finkbeiner, Mark	NOAA - CSC	mark.finkbeiner@noaa.gov	Technical advice, particularly on CMECS Use of benthic data
Ford, Kathryn	MA DMF	kathryn.ford@state.ma.us	Project partner Piloting Valentine et al. (2005) scheme
Kutcher, Tom	Narragansett Bay NERR	tomk@nberr.org	Advice and updates on NERRS scheme
Madden, Chris	NatureServe	cmadden@sfwmd.gov	Technical advice and updates on CMECS
Smith, Jan	Mass Bays NEP	jan.smith@state.ma.us	US EPA/NEP efforts to use habitat classification
Valentine, Page	USGS	pvalentine@usgs.gov	Technical advice and updates on classification of Stellwagen Bank
Wilber, Pace	NOAA	pace.wilber@noaa.gov	Technical advice and updates on CMECS
Wright, Nan	University of Idaho	nwright@uidaho.edu	Technical advice and updates on CMECS Lead on Columbia River Estuary pilot project